

2005

Case-Control Study of an Acute Aflatoxicosis Outbreak, Kenya, 2004

Karen E. Giesecker

Georgia State University, kgiesecker@gsu.edu

Eduardo Azziz-Baumgartner

Kimberly Lindblade

Helen Schurz Rogers

Georgia State University, hrogers1@student.gsu.edu

Stephanie Kieszak

See next page for additional authors

Follow this and additional works at: http://scholarworks.gsu.edu/iph_facpub

 Part of the [Public Health Commons](#)

Recommended Citation

Azziz-Baumgartner, E., Lindblade, K., Giesecker, K., Schurz Rogers, H., Kieszak, S., Njapau, H., Schleicher, R., McCoy, L.F., Misore, A., DeCock, K., Rubin, C., Laurence Slutsker, L. & the Aflatoxin Investigative Group. (2005). Case-control study of an acute aflatoxicosis outbreak, Kenya, 2004. *Environmental Health Perspectives*, 113(12), 1779–1783. DOI: 10.1289/ehp.8384

This Article is brought to you for free and open access by the School of Public Health at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Public Health Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

Authors

Karen E. Gieseke, Eduardo Azziz-Baumgartner, Kimberly Lindblade, Helen Schurz Rogers, Stephanie Kieszak, Henry Njapau, Rosemary Schleicher, Leslie F. McCoy, Ambrose Misore, Kevin DeCock, Carol Rubin, Lawrence Slutsker, and Aflatoxin Investigative Group

Case–Control Study of an Acute Aflatoxicosis Outbreak, Kenya, 2004

Eduardo Azziz-Baumgartner,¹ Kimberly Lindblade,² Karen Giesecker,³ Helen Schurz Rogers,¹ Stephanie Kieszak,¹ Henry Njapau,⁴ Rosemary Schleicher,¹ Leslie F. McCoy,¹ Ambrose Misore,⁵ Kevin DeCock,⁶ Carol Rubin,¹ Laurence Slutsker,⁷ and the Aflatoxin Investigative Group*

¹National Center for Environmental Health, ²National Center for Infectious Diseases, and ³Epidemiology Program Office, Centers for Disease Control and Prevention, Atlanta, Georgia, USA; ⁴Food and Drug Administration, Washington, DC, USA; ⁵Preventive and Promotive Health Services, Kenya Ministry of Health, Nairobi, Kenya; ⁶Centers for Disease Control and Prevention, Kenya Field Office, Nairobi, Kenya; ⁷Centers for Disease Control and Prevention, Kenya Field Office, Kisumu, Kenya

OBJECTIVES: During January–June 2004, an aflatoxicosis outbreak in eastern Kenya resulted in 317 cases and 125 deaths. We conducted a case–control study to identify risk factors for contamination of implicated maize and, for the first time, quantitated biomarkers associated with acute aflatoxicosis.

DESIGN: We administered questionnaires regarding maize storage and consumption and obtained maize and blood samples from participants.

PARTICIPANTS: We recruited 40 case-patients with aflatoxicosis and 80 randomly selected controls to participate in this study.

EVALUATIONS/MEASUREMENTS: We analyzed maize for total aflatoxins and serum for aflatoxin B₁–lysine albumin adducts and hepatitis B surface antigen. We used regression and survival analyses to explore the relationship between aflatoxins, maize consumption, hepatitis B surface antigen, and case status.

RESULTS: Homegrown (not commercial) maize kernels from case households had higher concentrations of aflatoxins than did kernels from control households [geometric mean (GM) = 354.53 ppb vs. 44.14 ppb; $p = 0.04$]. Serum adduct concentrations were associated with time from jaundice to death [adjusted hazard ratio = 1.3; 95% confidence interval (CI), 1.04–1.6]. Case patients had positive hepatitis B titers [odds ratio (OR) = 9.8; 95% CI, 1.5–63.1] more often than controls. Case patients stored wet maize (OR = 3.5; 95% CI, 1.2–10.3) inside their homes (OR = 12.0; 95% CI, 1.5–95.7) rather than in granaries more often than did controls.

CONCLUSION: Aflatoxin concentrations in maize, serum aflatoxin B₁–lysine adduct concentrations, and positive hepatitis B surface antigen titers were all associated with case status.

RELEVANCE: The novel methods and risk factors described may help health officials prevent future outbreaks of aflatoxicosis.

KEY WORDS: albumin adducts, aflatoxicosis, aflatoxin, Kenya, lysine, maize. *Environ Health Perspect* 113:1779–1783 (2005). doi:10.1289/ehp.8384 available via <http://dx.doi.org> [Online 9 August 2005]

During January–June 2004, the Kenya Ministry of Health (MOH) and partners identified 317 cases of acute hepatic failure in eastern Kenya; 125 cases occurred in persons who subsequently died during the illness. Seven patients had serum samples analyzed at the Kenya Medical Research Institute (KEMRI), and all were negative for viruses known to cause hepatic disease in Kenya (e.g., yellow fever; Rift Valley fever; dengue; acute hepatitis A, B, and C; West Nile virus; and Chikungunya and Bunyamwera) (American Public Health Association 2000). Because aflatoxicosis outbreaks had occurred previously in that geographical area, the MOH suspected that the unusually high number of patients with acute hepatic failure might have acquired aflatoxicosis from eating contaminated maize (Ngindu et al. 1982). Public health officials sampled maize from the affected area and found concentrations of aflatoxin B₁ as high as 4,400 ppb, which is 220 times greater than the 20 ppb limit for food suggested by Kenyan authorities (Onsongo 2004). Although aflatoxicosis outbreaks have occurred periodically in Africa and Asia, this outbreak resulted in the largest number of fatalities ever documented (Krishnamachari et al. 1975a, 1975b; Lye et al. 1995).

Aflatoxins are produced by *Aspergillus* spp. fungi that grow on a wide variety of grains and nuts (Patten 1981). The human gastrointestinal tract rapidly absorbs aflatoxins after consumption of contaminated food, and the circulatory system transports the aflatoxins to the liver (Fung and Clark 2004). From 1 to 3% of ingested aflatoxins irreversibly bind to proteins and DNA bases to form adducts such as aflatoxin B₁–lysine in albumin (Skipper and Tannenbaum 1990). Disruption of proteins and DNA bases in hepatocytes causes liver toxicity (Tandon et al. 1978).

Early symptoms of hepatotoxicity from aflatoxicosis can manifest as anorexia, malaise, and low-grade fever. Aflatoxicosis can progress to potentially lethal acute hepatitis with vomiting, abdominal pain, hepatitis, and death (Etzel 2002). Because aflatoxin B₁–lysine adducts are not repaired, their half-life in human serum is approximately 20–60 days (i.e., similar to that of unbound albumin) (McCoy L, personal communication; Sabbioni et al. 1987).

Information about risk factors associated with outbreaks of aflatoxicosis is limited. In addition, only a few animal studies have measured aflatoxin concentrations because unbound aflatoxins remain in the blood for a

very short period of time after exposure (i.e., 13–120 min) (Unger et al. 1977; Wong and Hsieh 1978). The primary objective of our case–control study was to identify risk factors for acute aflatoxicosis. The secondary objective was to determine the concentrations of aflatoxin in maize, bound aflatoxin in serum, and hepatitis B surface antigen associated with acute aflatoxicosis.

Materials and Methods

Selection of case patients. To focus the investigation on typical cases of presumed aflatoxicosis, our case definition was restricted to acute jaundice of unknown origin (i.e., no history of cirrhosis or obstructive liver disease) leading to hospitalization, during the peak of the epidemic, in the areas most affected by the outbreak. This case definition was based on information gathered by a descriptive epidemiology investigation conducted by the MOH and partners in May

Address correspondence to E. Azziz-Baumgartner, Centers for Disease Control and Prevention, Mailstop F46, 4770 Buford Hwy NE, Atlanta, GA 30341-3717 USA. Telephone: (770) 488-3412. Fax: (770) 488-3450. E-mail: eha9@cdc.gov

*Members of the Aflatoxin Investigative Group include J. Nyamongo, C. Njuguna, E. Muchiri, J. Njau, S. Maingi, J. Njoroge, J. Mutiso, J. Onteri, A. Langat, I.K. Kilei, G. Ogana, B. Muture, J. Nyikal (Kenya Ministry of Health); P. Tukei, C. Onyango, W. Ochieng (Kenya Medical Research Institute); I. Mugoya, P. Nguku, T. Galgalo, S. Kibet, A. Many, A. Dahiye, J. Mwhia, S. Likimani, C. Tetteh (Kenya Field Epidemiology and Laboratory Training Program/Kenya Ministry of Health); J. Onsongo, A. Ngindu (World Health Organization, Kenya Country Office); P. Amornkul, D. Rosen, D. Feiken, T. Thomas (CDC Kenya); P. Mensah, N. Eseko, A. Nejjar (World Health Organization, Regional Office for Africa); M. Onsongo, F. Kessel (Foreign Agricultural Service, U.S. Department of Agriculture); D.L. Park (Center for Food Safety and Applied Nutrition, Food and Drug Administration); C. Nzioka (Office of Global Health, CDC); L. Lewis, G. Luber, L. Backer, C.D. Powers, C. Pfeiffer (National Center for Environmental Health, CDC); W. Chege, A. Bowen (Epidemiology Program Office, CDC).

The use of trade names is for identification only and does not imply endorsement by the Centers for Disease Control and Prevention, the Agency for Toxic Substances and Disease Registry, the Public Health Service, or the U.S. Department of Health and Human Services.

The authors declare they have no competing financial interests.

Received 6 June 2005; accepted 9 August 2005.

2004. The descriptive epidemiology investigation found that a large number of patients with presumed aflatoxicosis had sought treatment at Makindu Sub-District Hospital (Makueni District) during 18 May–7 June and at Mutomo Mission Hospital (Kitui Districts) during 28 May–9 June. We did not restrict cases to live case patients or to case patients from which KEMRI had obtained blood samples because we did not want to introduce bias in our assessment of risk factors associated with disease.

To select 40 patients that met our case definition, we reviewed hospital records for the relevant time period and identified 19 case patients admitted to Makindu Sub-District Hospital and 21 case patients admitted to Mutomo Mission Hospital. All of the 29 case patients were alive at the time of the investigation, and all of the families of 11 deceased case patients verbally consented to participate in the study.

Selection of controls. We randomly selected two controls from each case patient's village because the descriptive epidemiology investigation suggested that these individuals would share similar soil, microclimate, and farming practices. Because the descriptive epidemiology investigation did not find a significant association among sex, case status, and case fatality, we did not match cases and controls by sex. To choose each control, we spun a bottle in front of the village elder's home and walked to the fifth house in the direction indicated by the bottle (or to the third house in sparsely populated areas). At the selected household, we identified all residents who had slept in the house the night before, and we used a random number list to select one of these household residents. We excluded infants who were solely breast-feeding because they would not have been directly exposed to aflatoxin B₁ found in maize. If selected individuals were not at their homes, we attempted to reach them wherever they were. All controls verbally consented to participate in the study.

Survey instrument. A literature review and the descriptive epidemiology investigation allowed development of hypotheses about the relationship between aflatoxicosis and methods of handling maize. We developed a questionnaire to elicit information about maize and protein consumption, the quality of home-grown and purchased maize products, maize storage and cooking practices, and associated illness and death of family members and pet dogs. All questions related to the relevant exposure period, which was designated as 1 month before the onset of case patients' illness or 1 month before controls heard about the outbreak.

Teams piloted the questionnaire on hospitalized patients who had presumed aflatoxicosis in Thika District. Local public health officials

translated the questionnaire, which was written in English, into Kikamba and Kiswahili as needed. Teams carried measuring cups to obtain standardized information on maize food portions consumed by participants.

Food sample collection. We obtained samples of maize products from participants to quantify personal exposure to aflatoxins. We collected samples from case households if they had maize in storage from the month before individuals developed aflatoxicosis (median date of symptom onset, 20 May 2004). We collected samples from control households if they had maize in storage from the month before hearing about the outbreak (median date of first hearing about the outbreak, 19 May 2004). We used metal cups to obtain multiple samples from different areas of the maize containers. These samples were combined in a paper bag to obtain 1 kg of maize for analysis. Collected maize products were replaced with commercial maize meal.

Blood sample collection. We obtained blood samples from participants to quantify their exposure to aflatoxins in the preceding month. With the exception of six case-patients from whom KEMRI had banked blood in May, we collected approximately 5–10 mL of venous blood in a Vacutainer tube with gel separators from all participants. All blood samples were transported on ice to KEMRI for serum separation.

Laboratory analysis. We analyzed maize samples using the VICAM AflaTest (VICAM, Watertown, MA, USA) immunoaffinity fluorometric method that quantitated total aflatoxin concentrations. Ground maize (50 g) that passed through a no. 20 sieve was mixed with 100 mL of a methanol:water mixture (80:20) with 5 g sodium chloride. The twice-filtered mixture (2 mL) was then passed through the immunoaffinity column at a rate of 1–2 drops/sec. The columns were washed with water, and the aflatoxins were recovered using 1 mL methanol. The methanol extract was read using a calibrated Vicam Series-4 Fluorometer set at 360 nm excitation and 450 nm emission. This method had an aflatoxin recovery of ≥ 85% and a detection limit of 1 ppb (VICAM 2001).

The Centers for Disease Control and Prevention (CDC) analyzed the serum specimens for aflatoxin B₁-lysine albumin adducts using high-performance liquid chromatography (HPLC) and isotope dilution tandem mass spectrometry (McCoy et al. 2005). After enzymatic hydrolysis of serum albumin, aflatoxin B₁-lysine adducts were extracted using solid-phase cartridges and separated using isocratic reversed-phase chromatography. We used positive ion electrospray with selected reaction monitoring mass spectrometry to measure aflatoxin B₁-lysine adducts and its corresponding D₄-labeled internal standard.

We measured total serum albumin using a bromocresol purple binding assay and a microplate reader. The limit of detection of aflatoxin B₁-lysine albumin adducts was 0.0003 ng/mg. The CDC also analyzed all remaining sera for hepatitis B surface antigen using ETI-MAK-2 PLUS enzyme immunoassay kits from DiaSorin (DiaSorin, Stillwater, MN).

Data management and analysis. Data were analyzed using SAS, version 8.02 (SAS Institute, Cary, NC). We used conditional logistic regression to calculate odds ratios (ORs) between case status and participants' methods of harvesting, storing, and preparing maize. We also used conditional logistic regression models to explore the relationship between case status, maize and protein consumption, aflatoxin concentrations in maize, aflatoxin B₁-lysine adduct concentrations, and hepatitis B surface antigen titers in serum. We restricted mixed linear regression models to controls because we wanted to investigate the relationship between serum aflatoxin concentrations and methods of harvesting, storing, and preparing maize, daily maize and protein consumption, and total aflatoxin concentrations in maize using a sample that more closely resembled the general population. We also used Cox proportional hazards models to explore the relationship between the number of days case patients survived after the onset of jaundice and aflatoxin concentrations in maize, aflatoxin B₁-lysine adducts concentrations in serum, hepatitis B surface antigen titers, and reported maize and protein consumption. Calculations were adjusted for age, sex, and participant's district.

Results

Demographic information. With few exceptions, case patients ($n = 40$) and controls ($n = 80$) had similar demographic characteristics (Table 1). Half of the participants lived in the Makueni District and the other half lived in the Kitui District. The mean age of case patients was similar to that of controls [22.5 years (range, 1.3–80.0 years) vs. 26 years (range, 0.5–75.0 years), respectively]. When

Table 1. Demographic characteristics [n (%)] of jaundiced case patients ($n = 40$) and village controls ($n = 80$), Eastern Province, Kenya, 2004.

Characteristic	Case patients	Controls	p -Value ^a
District			
Kitui	21 (52.5)	42 (52.5)	1.00
Makueni	19 (47.5)	38 (47.5)	
Mean age (years)	22.5	26.0	0.37 ^b
Age < 15 years	22 (55.0)	31 (38.8)	0.09
Male	25 (62.5)	27 (33.8)	0.003
Family with jaundice	15 (37.5)	3 (3.8)	< 0.001
Heard of outbreak	34 (85.0)	72 (90.0)	0.33

^aValues calculated using chi-square test unless indicated otherwise. ^bStudent's t -test used for comparison of means.

compared with controls, more of the case patients were male (62.5% vs. 33.8%, respectively; $p = 0.003$). Case patients were also more likely than controls to report having family members with acute jaundice during the 2 months before the study (37.5% vs. 3.8%; $p < 0.001$). As of 9 August, 18 of the 40 case patients (7 additional case patients since completion of our study) had died of acute liver failure.

Food consumption and maize aflatoxin analysis. Eating contaminated homegrown maize kernels was the primary risk factor for developing aflatoxicosis. On average, maize samples were collected 33 days (range, 8–112 days) after case-patients' onset of symptoms. Homegrown maize kernels from case households had significantly higher aflatoxin concentrations than kernels sampled from control households [geometric mean (GM) = 354.5 ppb vs. 44.1 ppb, respectively; $p = 0.04$; Figure 1]. Eating homegrown maize kernels was significantly associated with case status (adjusted OR = 3.0; 95% confidence interval (CI), 1.01–8.8). Owning "bad" homegrown maize kernels (maize with colored flecks, discoloration, unusual odor, or signs of mold) was found to be a risk factor for aflatoxicosis (adjusted OR = 5.9; 95% CI, 1.9–18.2). Case patients who fed their dogs household food reported dog deaths more often (43%) than controls (15%; adjusted OR = 15.2; 95% CI, 1.8–127.4). We did not find an association between case status and the number of portions of maize, beans, or meat participants consumed on a weekly basis.

Serum aflatoxin B₁-lysine adduct analysis. On average, serum samples were collected 33 days after case-patients' onset of symptoms. Using conditional logistic regression, we found that having aflatoxin B₁-lysine adduct concentrations at or above the median (0.25 ng/mg) was a risk factor for developing aflatoxicosis (adjusted OR = 14.8; 95% CI, 3.0–72.2). Case patients who provided serum samples ($n = 29$) had higher aflatoxin B₁-lysine adduct concentrations in their serum than did controls ($n = 62$; GM = 1.2 ng/mg of albumin vs. 0.15 ng/mg of albumin; $p < 0.001$; Figure 2). We found a

positive association between concentrations of aflatoxins in homegrown maize and aflatoxin B₁-lysine adduct concentrations in serum mixed linear regression adjusted for age, sex, village, and district ($p < 0.05$). For each milligram increase in the maize aflatoxin concentration, there was a 0.5 pg/mg increase in the logarithm of the serum aflatoxin B₁-lysine adduct concentration.

Serum hepatitis B surface antigen analysis. There was sufficient serum to analyze 72 (60%) samples for hepatitis B surface antigen. The mean age of participants with positive titers was 33 years, and most of them were female (58%). Eight (44%) of 18 cases had positive titers, while only 4 (7%) of 54 controls had positive titers (Table 2). Using conditional logistic regression, we found that having positive hepatitis B surface antigen titers was a risk factor for acute hepatic failure (adjusted OR = 9.8; 95% CI, 1.5–63.1). When we restricted the conditional logistic regression to participants with negative hepatitis B titers, we found that having aflatoxin B₁-lysine adduct concentrations at or above the median for this subgroup (0.2 ng/mg) was a risk factor for developing aflatoxicosis (95% CI, 2.1–∞, $p = 0.004$).

Risk associated with toxin. Case patients with known dates of death who had provided blood samples ($n = 8$) had higher aflatoxin B₁-lysine adduct concentrations in their serum than did case patients who survived ($n = 17$; 3.2 ng/mg vs. 0.5 ng/mg; $p = 0.07$) after adjusting for age, sex, and district. In our survival analysis, we found a significant association between time from jaundice to death and serum aflatoxin B₁-lysine adduct concentration (adjusted hazard ratio = 1.3; 95% CI, 1.04–1.6; $p = 0.02$).

Risk associated with food preparation and storage. Storing maize that was not completely dry and storing maize inside the home rather than in a granary were both independently associated with development of aflatoxicosis (OR = 3.5; 95% CI, 1.2–10.3; OR = 12.0; 95% CI, 1.5–95.7, respectively; Table 3). Participants who reported storing their maize mixed with ash had lower concentrations of aflatoxins in their maize than those who did

not (GM = 17.4 ppb vs. 142.2 ppb; $p = 0.05$). We did not find an association between case status, the type of container used to store maize (plastic burlap, plastic bucket, woven basket, clay pot, gourd, or sisal), the use of soda and pesticides in the storage area, or the culling of maize kernels that appeared moldy.

Discussion

Food consumption and aflatoxin analyses. This is the first investigation to quantify the association among environmental contamination, a history of exposure, biomarker concentrations, and acute aflatoxicosis. The results of our case-control study suggest that consumption of contaminated maize kernels placed people in this region of Kenya at risk for life-threatening aflatoxicosis (case-fatality rate of 39%). Through systematic sampling of maize and serum from participants, we found a strong association between aflatoxin concentrations in homegrown maize, serum B₁-albumin adducts, hepatitis B surface antigen titers, and case status.

The aflatoxin concentrations measured from the maize of case patients was comparable with those measured in other acute aflatoxicosis outbreaks. The aflatoxin B₁-lysine adduct concentrations measured from the serum of case patients are the highest ever reported. This is the first study to quantify aflatoxin B₁-lysine adduct concentrations in the serum of case patients during an outbreak of acute aflatoxicosis; a critical step in the elucidation of the clinically relevant action levels for aflatoxin exposure. We associated these serum aflatoxin B₁-lysine adduct concentrations with the risk for life-threatening acute aflatoxicosis.

Table 2. Serum aflatoxin B₁-lysine albumin adduct concentration and hepatitis B surface antigen titers (μg/mg of albumin) in cases and controls [GM (n)].

	Adduct concentration	
	Cases	Controls
Hepatitis B positive	0.17 (8)	0.08 (4)
Hepatitis B negative	3.55 (10)	0.16 (50)

Table 3. Risk factors [n (%)] for jaundice among case patients ($n = 28$) and controls ($n = 43$) who ate maize kernels grown on their own farms, Kenya, 2004.

Characteristic	Case patients	Controls	OR (95% CI)
Initial dryness of stored maize			
Wet	15 (53.6)	11 (25.6)	3.5 (1.2–10.3)
Dry	13 (46.4)	32 (74.4)	1.0
Storage location			
House	22 (81.5)	23 (53.5)	12.0 (1.5–95.7)
Granary	5 (18.5)	20 (46.5)	1.0
Preservatives added to storage			
Ash	6 (15.4)	13 (17.6)	1.6 (0.4–5.6)
Insecticide	9 (23.1)	21 (28.1)	0.6 (0.2–1.8)




Figure 1. Frequency of total maize aflatoxin concentrations for participants.




Figure 2. Frequency of serum aflatoxin B₁-lysine albumin adduct concentrations for participants.

We found an association between aflatoxin concentrations in maize and aflatoxin B₁-lysine adduct concentrations in serum from controls. The GM aflatoxin B₁-lysine adducts concentration in serum from controls is higher than the majority of concentrations documented in population-based studies from countries with a high incidence of liver cancer (Wild et al. 1990). It is unclear why some controls with high aflatoxin B₁-lysine adduct concentrations did not manifest symptoms of acute hepatitis during the time of the investigation. The concentrations found in controls were not associated with acute symptoms and may have represented chronic exposure to aflatoxins. Chronic exposure to aflatoxins is associated with impaired immunity, malnutrition, and liver cancer (the third most common cause of death from cancer in Africa) (Parkin et al. 2003; Williams et al. 2004). People chronically exposed to elevated concentrations of aflatoxins are three times more likely to develop hepatocellular carcinoma.

We also found an independent association between hepatitis B surface antigen titers and case status. Although people with hepatitis B (which is endemic in Kenya) who are chronically exposed to aflatoxins may be more likely to develop hepatocellular carcinoma, this is the first study to quantify the association between hepatitis B, aflatoxin adducts, and acute hepatitis (Keenlyside et al. 1977; Qian et al. 1994). Further research is needed to determine if the high incidence of liver cancer in eastern Kenya is attributable to chronic asymptomatic exposure to aflatoxins. In addition, clinicians working in areas where aflatoxicosis is endemic should consider obtaining a dietary history for aflatoxin exposure from cases patients with symptoms of acute hepatitis and positive hepatitis B titers.

Risk factors. Our case-control study quantified ORs for suspected risk factors described in previous aflatoxicosis outbreaks. As in a 1974 outbreak in India (Krishnamachari et al. 1975b), we found that males were more likely to die from aflatoxicosis, in spite of eating similar quantities of maize as females. We found that acute aflatoxicosis manifests in family clusters, as reported in a 1988 outbreak in Malaysia (Lye et al. 1995). Sharing contaminated food and genetic polymorphisms of cytochrome P₄₅₀ enzymes may place families at risk for aflatoxicosis (Chen et al. 2000). As reported by Ngindu (1982) in a 1981 outbreak in Kenya, we found that, more often than controls, case patients reported dog deaths before developing aflatoxicosis. In the future, reports of deaths in dogs may warn public health officials of a potential aflatoxin contamination of the food supply.

Food preparation and storage analysis. Although maize is traditionally stored in granaries, storage inside homes occurs during

periods of food shortage; this may have facilitated the contamination of maize with aflatoxins. The rainy season (from March through May) accounts for 80% of annual food production [Food and Agriculture Organization (FAO) 2000]. In 2004, an early and insufficient rainy season caused a food shortage of 156,000 metric tons of maize (Associated Press 2004). Some participants reported storing maize inside their homes to ensure it would not be stolen during the food shortage. Drought conditions stress maize plants and render them susceptible to contamination by *Aspergillus* spp. (Wilson and Payne 1994). The warm environment inside these windowless homes and storage of maize on the dirt floor may have promoted fungal growth in wet maize kernels.

Our case-control study suggests that traditional methods of drying and storing maize in elevated granaries were protective against aflatoxicosis. Traditional granaries are raised structures that are well ventilated, and they promote the drying of grain (FAO 1998). The granaries' elevated platforms isolate the maize from spores and insects on the ground. We also found that storing maize mixed with ash was associated with lower concentrations of aflatoxin than storing maize without ash. Ash acts as a physical barrier against insects and helps keep maize dry.

Limitations. Our case-control study was limited by its retrospective design. It is possible that case patients (or the family members of deceased case patients) may have recalled the amount, source, and quality of maize that was consumed differently than did controls. The aflatoxin concentrations measured in sampled maize may have differed from those consumed by case patients before they became ill with aflatoxicosis. We may not have found an association between the number of portions of maize consumed and case status due to the limited accuracy of the food questionnaires. In addition, it is possible that some case patients developed jaundice as a result of undiagnosed medical conditions unrelated to aflatoxicosis. This potential misclassification would have weakened any demonstrable associations.

Conclusion

Aflatoxins and other mycotoxins contaminate 25% of agricultural crops worldwide and are a source of morbidity and mortality throughout Africa, Asia, and Latin America (Smith et al. 1994). To prevent future aflatoxicosis outbreaks, it is necessary to explore public health interventions that promote effective production, storage, and processing of homegrown and commercial maize. In addition, surveillance that monitors aflatoxin concentrations in food and incidence of acute jaundice in humans may prevent widespread outbreaks of acute aflatoxicosis (Trucksess and Wood 1994). In the future, serum aflatoxin B₁ albumin

adducts may be used to diagnose acute aflatoxicosis and monitor interventions aimed at reducing aflatoxin exposure (Kensler et al. 1999). Although short-term interventions such as food replacement mitigate the loss of life during outbreaks, it is necessary to develop long-term, culturally appropriate strategies to prevent aflatoxicosis.

REFERENCES

- American Public Health Association. 2000. Arthropod-borne viral diseases (arboviral diseases). In: *Control of Communicable Diseases Manual* (Chin J, ed). 17th ed. Washington, DC: American Public Health Association, 30-31.
- Chen SY, Chen CJ, Tsai WY, Ahsan H, Liu TY, Lin JT, et al. 2000. Associations of plasma aflatoxin B₁-albumin adduct concentration with plasma selenium concentration and genetic polymorphisms of glutathione S-transferase M1 and T1. *Nutr Cancer* 38:179-185.
- Etzel R. 2002. Mycotoxins. *JAMA* 287:425-427.
- FAO. 1998. African Experience in the Improvement of Post-harvest Techniques. Rome: Food and Agriculture Organization of the United Nations. Available: http://www.fao.org/documents/show_cdr.asp?url_file=/docrep/W1544E/W1544E00.htm [accessed 22 February 2005].
- FAO (Food and Agriculture Organization of the United Nations). 2000. Drought in Kenya: Food Situation Is Deteriorating Rapidly - First Starvation-Related Deaths. Press Release 00/40. Available: http://www.fao.org/WAICENT/OIS/PRESS_NE/PRESSENG/2000/pren0040.htm [accessed 22 February 2005].
- Fung F, Clark R. 2004. Health effects of mycotoxins: a toxicological overview. *J Toxicol* 42:217-234.
- International Federation of Red Cross and Red Crescent Societies. 2004. Kenya: Drought. Appeal no. 18/04. Available: <http://www.ifrc.org/docs/appeals/04/1804.pdf> [accessed 24 October 2005].
- Keenlyside RA, Smith DH, Hirst D, Zuckerman AJ, Preece J. 1977. The distribution and significance of hepatitis B surface antigen in a rural population in Kenya. *Ann Trop Med Parasitol* 71:167-177.
- Kensler TW, Groopman JD, Sutter TR, Curphey TJ, Roebuck BD. 1999. Development of cancer chemopreventive agents: oltipraz as a paradigm. *Chem Res Toxicol* 12:113-126.
- Krishnamachari KA, Bhat RV, Nagarajan V, Tilak TB. 1975a. Hepatitis due to aflatoxicosis: an outbreak in western India. *Lancet* 1:1061-1063.
- Krishnamachari KA, Bhat RV, Nagarajan V, Tilak TB. 1975b. Investigations into an outbreak of hepatitis in parts of western India. *Indian J Med* 63:1036-1049.
- Lye MS, Ghazali AA, Mohan J, Alwin N, Nair RC. 1995. An outbreak of acute hepatic encephalopathy due to severe aflatoxicosis in Malaysia. *Am J Trop Med Hyg* 53:68-72.
- McCoy LF, Scholl PF, Schleicher RL, Groopman JD, Powers CD, Pfeiffer CM. 2005. Analysis of aflatoxin B₁ lysine adduct in serum using isotope-dilution liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom* 19:2203-2210.
- Ngindu A, Kenya PR, Ocheng DM, Omondi TM, Ngare W, Gatei D, et al. 1982. Outbreak of acute hepatitis caused by aflatoxin poisoning in Kenya. *Lancet* 1:1346-1348.
- Onsongo J. 2004. Outbreak of aflatoxin poisoning in Kenya. *EPI/IDS Bull* 5:3-4. Available: <http://www.afro.who.int/csr/ids/bulletins/eastern/jun2004.pdf> [accessed 20 October 2005].
- Parkin DM, Ferlay J, Hamdi-Chérif M, et al. 2003. Liver Cancer. *IARC Sci Publ* 153:299-314.
- Patten RC. 1981. Aflatoxins and disease. *Am J Trop Med Hyg* 30:422-425.
- Qian GS, Ross RK, Yu MC, Yuan JM, Gao YT, Henderson BE, et al. 1994. A follow-up study of urinary markers of aflatoxin exposure and liver cancer risk in Shanghai, People's Republic of China. *Cancer Epidemiol Biomarkers Prev* 3:3-10.
- Sabbioni G, Skipper PL, Buchi G, Tannenbaum SR. 1987. Isolation and characterization of the major serum albumin adduct formed by aflatoxin B₁ in vivo in rats. *Carcinogenesis* 8:819-824.
- Smith JE, Solomons GL, Lewis CW, Anderson JG. 1994. Mycotoxins in Human Health. Brussels: European Commission.
- Skipper PL, Tannenbaum SR. 1990. Protein adducts in the

- molecular dosimetry of chemical carcinogens. *Carcinogenesis* 11:507–518.
- Tandon H, Tandon B, Ramalingaswami V. 1978. Epidemic of toxic hepatitis in India of possible mycotoxic origin. *Arch Pathol Lab Med* 102:372–376.
- Trucksess MW, Wood GE. 1994. Recent methods of analysis for aflatoxins in foods and feeds. In: *The Toxicology of Aflatoxins, Human Health, Veterinary, and Agricultural Significance* (Eaton DL, Groopman JD, eds). San Diego, CA:Academic Press, 409–431.
- Unger PD, Mehendale HM, Hayes AW. 1977. Hepatic uptake and disposition of aflatoxin B₁ in isolated perfused rat liver. *Toxicol Applied Pharmacol* 41:523–534.
- VICAM. 2001. About Aflatest®. Available: <http://www.vicam.com/products/aflatest.html> [accessed 18 February 2005].
- Wild CP, Jiang YZ, Allen SJ, Jansen LA, Hall AJ, Montesano R. 1990. Aflatoxin-albumin adducts in human sera from different regions of the world. *Carcinogenesis* 11:2271–2274.
- Williams J, Phillips TD, Jolly PE, Stiles JK, Jolly CM, Aggarwal D. 2004. Human aflatoxicosis in developing countries: a review of toxicology, exposure, potential health consequences, and interventions. *Am J Clin Nutr* 80:1106–1122.
- Wilson DM, Payne GA. 1994. Factors affecting *Aspergillus flavus* group infection and aflatoxin contamination of crops. In: *The Toxicology of Aflatoxins: Human Health, Veterinary, and Agricultural Significance* (Eaton DL, Groopman JD, eds). San Diego, CA:Academic Press, 309–315.
- Wong ZA, Hsieh DPH. 1978. Aflatoxicol: major aflatoxin B₁ metabolite in rat plasma. *Science* 200:325–327.

The American Plastics Council respectfully requests that *EHP* address the misinformation that appeared in these articles and which is available on the *EHP* website.

The author is employed by the American Chemistry Council/American Plastics Council.

Patricia A. Enneking
American Plastics Council
Arlington, Virginia
E-mail: patty_enneking@plastics.org

REFERENCE

- Hood E. 2005. Are EDCs blurring issues of gender? *Environ Health Perspect* 113:A670–A677.
Phillips ML. 2005. Children's centers study kids and chemicals. *Environ Health Perspect* 113:A664–A668.
SC Johnson. 2006. Bogus E-Mail Attacking Plastic Wraps Refuted. Available: http://www.scjohnson.com/family/fam_pre_pre_news.asp?art_id=64 [accessed 12 January 2006].

Editor's note: The following erratum was published in the January 2006 issue (Environ Health Perspect 114:A21):

In the October articles "Children's Centers Study Kids and Chemicals" [Environ Health Perspect 113:A664–A668 (2005)] and "Are EDCs Blurring Issues of Gender?" [Environ Health Perspect 113:A670–A677 (2005)], photographs and their captions erroneously imply that plastic drink bottles contain ortho-phthalates. Plastic drink bottles sold in the United States are made from polyethylene terephthalate and do not contain ortho-phthalates. Also, at the end of the EDCs article, references are made to plastic wrap and Saran Wrap. For clarification, neither plastic wrap nor Saran Wrap contains ortho-phthalates. EHP regrets these errors.

ERRATA

Azziz-Baumgartner et al. noticed two errors in "Case-Control Study of an Acute Aflatoxicosis Outbreak—Kenya?" [*Environ Health Perspect* 113:1779–1783]. The units in Figure 2 and Table 2 should be nanograms per milligram instead of micrograms per milligram. The errors were introduced when new figures and tables were generated during the final revision of the paper. The authors apologize for these errors.

In the article by Feist et al. [*Environ Health Perspect* 113:1675–1682], the units were incorrect in several figures and tables: "Lipid ($\mu\text{g/g}$)" should be " $\mu\text{g/g}$ lipid" in Tables 1 and 2 and in the y -axes of Figures 2 and 3A–C. Also, on the y -axes in Figure 5A–D, "dL" should be "mL." *EHP* regrets these errors.

The photograph on page A29 of the January 2006 NIEHS News section should have been credited to Jennifer Gorenstein/UTMDACC COEP. The photographs on page A30 should have been credited to Tom Van Biersel/Louisiana Geological Survey (left) and Bryan Parras/UTMB (right). Additionally, Parras's photograph depicts residents of Pointe-aux-Chenes, not LaRose, and includes no COEP staff.

In the Beyond the Bench article in this same section, "COEPs Contribute to Hurricane Relief" [*Environ Health Perspect* 114:A30–A31 (2006)], Peter Thorne was incorrectly identified as director of the University of Iowa COEP; he is in fact director of the University of Iowa Environmental Health Sciences Research Center as well as head of the NIEHS Working Group on Mold, Microbial Agents, and Respiratory Diseases. It was the latter group that "collected air and surface samples from water-damaged homes in New Orleans" as our article stated. Finally, the aid teams that traveled throughout Louisiana included members from the UTMDACC COEP as well as the UTMB COEP.

EHP regrets the errors.

In the January Focus article "In Katrina's Wake" [*Environ Health Perspect* 114:A32–A39 (2005)], Hurricane Katrina was identified as a Category 4 storm, reflecting statements from the National Hurricane Center as of press time. The National Hurricane Center has since reported that Katrina was actually a Category 3 storm at the time of landfall.