

2015

# Inter and Intra Buffer Variability: A Case Study Using Scale

Dawn Haynie

Georgia State University, [Shaynie@gsu.edu](mailto:Shaynie@gsu.edu)

Follow this and additional works at: [http://scholarworks.gsu.edu/art\\_design\\_facpub](http://scholarworks.gsu.edu/art_design_facpub)



Part of the [Architecture Commons](#), and the [Art and Design Commons](#)

---

## Recommended Citation

Haynie, Dawn. "Inter and intra buffer variability: A case study using scale," 10th International Space Syntax Symposium Proceedings, London, England July 2015. [http://www.sss10.bartlett.ucl.ac.uk/wp-content/uploads/2015/07/SSS10\\_Proceedings\\_050.pdf](http://www.sss10.bartlett.ucl.ac.uk/wp-content/uploads/2015/07/SSS10_Proceedings_050.pdf).

This Conference Proceeding is brought to you for free and open access by the Ernest G. Welch School of Art and Design at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Art and Design Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

050

## Inter and intra buffer variability:

### A case study using scale

---

**S Dawn Haynie**Georgia State University  
shaynie@gsu.edu

---

**Abstract**

*Many morphological studies select sample sets to explore neighbourhoods of interest, particularly in terms of their structural street properties, measures of scale or density, and proximity to the metropolitan center (Cervero and Gorham, 1995, Crane and Crepeau, 1998, French and Scoppa, 2007, Handy et al., 2003, Jacobs, 1993, Peponis et al., 2007, Southworth and Owens, 1993); yet beyond the established and distinctive structures of these neighborhoods, few have analyzed, in depth, the variability in their measures. This study randomly samples 4,321 localities from the 24 largest American metropolitan areas and describes a method using the measures of length and area to evaluate the variability both between and within these localities. Calculated as the standard deviation of mean scale, Inter Buffer Variability is introduced to describe the variation between these localities while Intra Buffer Variability describes the variation, or consistency, within these localities. How varied then are the measures of scale, and are the measures for some MSAs more varied than others? As will be shown, the MSA Inter Buffer Variability for both length and area are broad, which is expected given both the urban and suburban localities captured across each MSA; and yet, the MSA Intra Buffer Variability is also broad suggesting more variation within these localities than originally suggested by the samples illustrated within the literature. Comparatively for each measure of length and area, both Inter and Intra Buffer Variability are graphed one in relation to the other with their associated means used to delineate those trending higher or lower than average. Interestingly, four quadrants emerge distinctively delineating the measures of scale for these MSAs.*

**Keywords**

*Urban morphology, urban design, variability, neighbourhood scale.*

**1. Introduction**

Beyond the differences demonstrated between neighborhoods and their associated measures of scale, density, or connectivity, few have been able to analyze, in depth, variability in the sprawling landscape of the metropolitan city. Methodologically, many studies have extracted representative samples from larger metropolitan areas to illustrate extremes between various neighborhoods or localities of particular interest (Cervero and Gorham, 1995; Crane and Crepeau, 1998; Doxiadis, 1968; Frank et al., 2007; Handy et al., 2003; Hess, 1997; Jacobs, 1993; Jo, 1998; Ozbil and Peponis, 2007; Peponis et al., 2007; Siskna, 1997; Southworth and Owens, 1993). These neighborhoods were often chosen for their purity of structural type, planning history, demographics, or primary mode of transportation. Statistical inferences were drawn, and frequently, these results were generalized to describe the consistency, density, and pedestrian-oriented context of the city center in juxtaposition to the varied, sprawling, auto-oriented context of the peripheral suburb.

In a global context, Doxiadis (1968) illustrated variations in scale across different metropolitan areas to describe patterns of growth, as did Abler and Adams (1976), Passonneau and Wurman (1966), and Adams in his discussion of the New York Regional Plan (1929). In a local context, Cervero and Gorham (1995) illustrated differences in scale between neighborhoods with prevailing modes of choice for transit; Handy (2003) illustrated differences between neighborhoods of specific structural interest; Jacobs (1993) illustrated differences in the measures between neighborhoods of significant historical interest; and Peponis et al. (2007) demonstrated differences between localities influenced by various planning policies and urban design initiatives.

In each of these cases, the measures of road segments and blocks were analyzed to offer a fundamental sense of the scale or size of urban elements that combine to form the texture of the urban fabric; and yet, given the method of selective sampling, these neighborhoods are not necessarily representative of the city in its entirety, essentially ignoring the variability experienced between or within these illustrated extremes.

If neighborhoods, or localities, were sampled randomly, with equitable probability, and for a population size that yields statistically significant results, would the inferences in the measures, as established in the literature, persist? How varied are the average measures between localities, how varied are the measures within each locality, and perhaps more fundamental, how can the variability in scale be described analytically?

## 2. Defining Measures of Variability

To discuss variability, two distinctions are suggested. First, a measure to describe differences *between* the average measures of each locality is defined to capture, as an example, differences between the average lengths of road segments for a city center in comparison to the average for a remote suburb. Second, an alternate measure to describe differences in the measures *within* each locality is defined to capture, as an example, the consistency in the lengths of road segments for city centers with a strong planning initiative.

### *Inter Buffer Variability:*

describes differences between the average measures of neighborhoods; thus, it is calculated as the standard deviation for a set of neighborhood means. For example, the Inter Buffer Variability of Length is calculated by taking the standard deviation of mean road length for a sampled set of neighborhoods.

### *Intra Buffer Variability:*

describes differences within the measures of any particular neighborhood, thus, it is calculated as the mean for a set of neighborhood standard deviations, which is calculated for any set of elements within that neighborhood. For example, the Intra Buffer Variability of Length is calculated as the mean of the standard deviation of road length for roads within a sampled set of neighborhoods.

## 3. Method for constructing a Randomly Sampled Set

To ensure an equitable distribution of sampled areas, a framework was established. From a defined point of center, rings radiated outward at a distance relevant to the scale of the maps studied; and a coordinate system, fixed by the point of center, was superimposed and rotated 45 degrees to define rather than divide the quadrants of North, South, East and West. From each section of this established framework, x and y coordinates were randomly selected at a particular distance and degree from the designated point of center. Included was a provision for eliminating the potential of overlapping areas such that all selected buffers were complete and distinct from one another. These randomly selected

coordinates, along with each point of center, were imported into ESRI software to create circular buffers at a radius relevant to the variables being studied and then used to extract spatial information to describe smaller, more local areas within the maps.

#### 4. Case Study analyzing the variability of scale in American Cities


To study the urban and suburban conditions found across the United States, 24 of the largest, most populated cities were selected for random sampling. These MSAs included: Atlanta, Baltimore, Boston, Chicago, Cincinnati, Cleveland, Dallas, Denver, Detroit, Houston, Los Angeles, Miami, Minneapolis, New York City, Philadelphia, Phoenix, Pittsburgh, Portland, San Diego, San Francisco, Seattle, St. Louis, Tampa, and Washington D.C. For each MSA, GIS based vector data was compiled from the Street Map and County databases released by ESRI in their ArcMap software. Data was decompressed, exported, converted into various shape files, dissolved, clipped and then eventually merged into a single shape file representative of each MSA.

Initially, each city was defined simply by the legal boundary of its larger Metropolitan Statistical Area (MSA);<sup>1</sup> yet in several cases, the overall density and development of the city was continuous across the landscape from one MSA to another. In these cases, the two MSAs were combined into a single area for analysis to complete the overall morphology of the city and to reduce any possible distortions in the measures due to 'edge effects.' These combinations include the union of Cleveland with Akron, Denver with Boulder, Los Angeles with Riverside and Ventura, Philadelphia with Trenton, and San Francisco with San Jose. In comparison to the metropolitan areas originally evaluated by Abler and Adams (1976), the MSAs selected here captured 5% of the total land area held within the contiguous U.S., and they represented 49.5% of the population.

In this selected set of MSAs, the point of center was established by the position of the original City Hall and/or a similarly associated, politically significant building in the MSA. Rings radiated outward from the point of center at 5, 15, 30, and 60 mile intervals (Figure 1). From each section of this established framework, coordinates were randomly selected using a script programmed in Java. From these randomly selected coordinates, along with each point of center, circular buffers measuring 2 miles in diameter were established.


---

<sup>1</sup> The Metropolitan Statistical Areas were defined per the MSA Boundary Map, as referenced on May 2006: [http://ftp2.census.gov/geo/maps/metroarea/us\\_wall/jun2003/cbsa\\_us\\_0603\\_rev.pdf](http://ftp2.census.gov/geo/maps/metroarea/us_wall/jun2003/cbsa_us_0603_rev.pdf)



**Figure 1:** Illustration of the Framework, overlaid Atlanta as an example, to ensure an equitable distribution for the Randomly Sampled Buffers

Intending to capture 10% of the total land area, 363 coordinate pairs were identified for each of the selected MSAs. Included was a provision for eliminating the potential of overlapping areas; and thus, all selected buffers were complete and distinct from one another. If all coordinate pairs captured development, the Randomly Sampled Set would have contained 8,712 buffers for study; but inevitably, many fell either outside the political boundary of the MSA or in rural, undeveloped areas (Figure 2).



**Figure 2:** Illustrations of the various areas (shown in a darker grey) captured by the Randomly Sampled Buffers (shown in black)

To test this methodology and to access the effectiveness of the proposed definitions for variability, the following measures were extracted from the randomly sampled set:

*Length*

:the distance (feet) between two choice intersections, with road segments drawn as street centerlines

*Buffer Length*

:the sum length of road segments (feet) in a defined buffer divided by the number of road segments captured by that buffer; i.e. the mean length of road segments for a defined buffer

*Area*

:the landmass (acres) of a block bounded by a continuous set of road segments, with road segments drawn as street centerlines

*Buffer Area*

:the sum area of blocks (acres) in a defined buffer divided by the number of blocks captured by that buffer; i.e. the mean area of blocks for a defined buffer

In addition to those buffers capturing undeveloped areas, road segments and blocks of extreme scale, both large and small, were identified and excluded. To prevent potential distortion from extremes within the database, the work of Thomas Jefferson and his influence on the Land Ordinance of 1785 (Rashid, 1996) was assessed in conjunction with the work of Doxiadis (1965) and Leon Krier (1976) to set parameters for pragmatically defining and removing extremes. Blocks more than 640 acres in area or less than 0.12 acres were excluded. Similarly, road segments more than 1 mile in length or less than 72 feet were excluded.

## 5. Limitations of the Database


While there were no significant 'edge effects', the measures were subject to a number of other limitations. Area calculations were accurate, subject to the projections necessary for GIS to represent a spherical model in a single plane of two dimensions.<sup>2</sup> Block area included not only the sum area of its parcels but a portion of its defining streets as well because the vectors were drawn to represent the centerline of each street without consideration of street width. In addition, the calculations of available street length were accurate, subject to the reliability of the available GIS data.<sup>3</sup>

In consequence to the defined method of sampling, several complications were encountered, which limit the inferences that can be made of the resulting Randomly Sampled Set. First, the automated process for sampling allowed for the creation of a larger set of samples than would otherwise be possible; however, the size of the Randomly Sample Buffers then made it impossible to examine the data to correct errors.

<sup>2</sup> The area calculation for each MSA was calculated from the .shp files of the ESRI Database in a NAD 83 projection.

<sup>3</sup> According to ESRI, their databases were created and compiled from individual TIGER files and subsequently corrected in house to remove traditional issues of alignment and varying coordinate systems when connecting larger networks. The accuracy of this data remains subject to the individual error declared by ESRI. The individual line segments of each road network have not been verified or corrected here within this study, but the error within each overall network system chosen for analysis is presumed to be minimal given the scale of the analysis and thus, should have little impact on the results.

As a result, the Randomly Sampled Set of Buffers was only as good as the original set from which the data was drawn, without interventions to correct inaccuracies. Second, many of the Randomly Sampled Buffers captured significantly more area or length than was initially intended, given that the blocks and road segments intersecting each buffer were included along with those completely contained within (Figure 3). As a result, the Randomly Sampled Sets of blocks and road segments were not complete, congruent and comprehensive sets. For this reason, statistical correlations between the measures of the Randomly Sampled Buffers should be considered carefully.



**Figure 3:** Illustrations of the associated blocks (shown in a darker grey) and road segments (shown in black) captured within each of the Randomly Sampled Buffers

## 6. Analyzing Inter and Intra Buffer Variability

Measures for the Randomly Sampled Buffers were studied to assess the variations between them. Calculated as the standard deviation of Buffer Length or Buffer Area, the Inter Buffer Variability described the variability or consistency found between the measures of Buffer Length or Buffer Area for the Randomly Sampled Set. Subsequently, the MSA Inter Buffer Variability of Length or Area described the average variation or consistency found between the measures of buffer scale within a particular MSA.

In addition to analyzing the variation in the measures of scale between buffers, or the differences in Buffer Scale for these Randomly Sampled Buffers, variations in the measures of Length and Area were also considered within each buffer. Calculated as a mean for the standard deviation of Buffer Scale, Intra Buffer Variability of Length and Area described the variability or consistency found within the measures of Length or Area for the Randomly Sampled Buffers. Subsequently, the MSA Intra Buffer Variability of Length and Area described the average variation or consistency found among the measures of the Randomly Sampled Road Segments or Blocks within each buffer.

For the Randomly Sampled Buffers, Inter Buffer Variability of Length measures 855.12 feet (Table 1). When calculated for each metropolitan area, the MSA Inter Buffer Variability of Length varies from 458.71 in Boston to 1070.88 feet in Phoenix. Likewise, Inter Buffer Variability of Area measures 144.36 acres (Table 2), and when calculated for each metropolitan area, the MSA Inter Buffer Variability of Area varies from 67.67 acres in Boston to 191.55 acres in Minneapolis – St. Paul.

Notably, Atlanta, Boston, New York City, and Washington D.C. illustrate a lower MSA Inter Buffer Variability of Length and Area, which suggests greater consistency between the measures of their buffers. Los Angeles – Riverside – Ventura, Phoenix, San Diego, San Francisco – San Jose, and Seattle exhibit a higher MSA Inter Buffer Variability of Length with a lower MSA Inter Buffer Variability of Area, which suggests relative consistency between the measures of Buffer Area despite variability between the

measures of Buffer Length. Lastly, Cincinnati, Dallas, and St. Louis exhibit a lower MSA Inter Buffer Variability of Length with a higher MSA Inter Buffer Variability of Area, which suggests relative consistency between the measures of Buffer Length despite variability between the measures of Buffer Area.

**Table 1:** Statistical Measures of Buffer Length for the Randomly Sampled Buffers

	MSA	(n) Randomly Sampled Buffers	MSA Mean of Buffer Length (feet)	MSA Median of Buffer Length (feet)	MSA Intra Buffer Variability Mean [Std Dev of Buffer Length] (feet)	MSA Std Deviation [Buffer Length] (feet)
all MSAs		4321	1382.6562	1158.4521	968.4598	855.12
Atlanta		291	1279.3100	999.0812	992.8728	639.97
Baltimore		90	1145.9438	927.0117	874.3102	627.74
Boston		125	921.0272	674.9106	785.2839	458.71
Chicago		213	1328.2620	1115.2535	884.1587	1023.93
Cincinnati		155	1516.6831	1259.1298	1094.8248	669.08
Cleveland - Akron		101	1528.7196	1292.0128	1036.9714	985.69
Dallas		258	1450.6091	1200.6352	1038.7708	734.94
Denver - Boulder		190	1790.6753	1578.7486	1117.2205	1058.15
Detroit		134	1323.7026	1098.7564	934.3410	853.93
Houston		280	1423.4125	1186.7527	998.2042	856.64
Los Angeles – Riverside - Ventura		212	1195.4960	1006.4543	822.9983	856.85
Miami		68	926.6310	744.7042	756.9758	693.21
Minneapolis - St. Paul		225	1671.2101	1441.4431	1086.7686	965.85
New York City		209	894.8752	683.1129	724.4346	566.01
Philadelphia - Trenton		188	1171.5391	931.5364	895.3749	700.21
Phoenix		209	1751.4890	1594.6980	1048.7428	1070.88
Pittsburgh		189	1259.9701	987.0297	1017.4978	662.22
Portland		211	1532.9129	1365.6079	1036.3200	800.23
San Diego		114	1365.8487	1165.2267	969.0424	918.59
San Francisco - San Jose		181	1526.1984	1303.6005	1000.4448	1042.18
Seattle		150	1569.4133	1390.3612	971.0437	930.79
St. Louis		266	1568.2355	1332.2371	1081.8066	739.21
Tampa - St. Petersburg		96	1186.9352	964.4266	840.5445	769.50
Washington D.C.		166	1176.1295	940.7750	890.5396	651.96

Alternately, for Length within these Randomly Sampled Buffers, Intra Buffer Variability of Length measures 968.46 feet (Table 1). When calculated for each metropolitan area, the MSA Intra Buffer Variability of Length varies from 724.43 feet in New York City to 1117.22 feet in Denver – Boulder. Likewise for Area, the Intra Buffer Variability of Area measures 118.89 feet (Table 2). When calculated for each metropolitan area, the MSA Intra Buffer Variability of Area varies from 71.31 feet in Tampa – St. Petersburg to 156.99 feet in Pittsburgh.




**Table 2:** Statistical Measures of Buffer Area for the Randomly Sampled Buffers

MSA	Randomly Sampled Buffers (n)	MSA Mean of Buffer Area (acre)	MSA Median of Buffer Area (acre)	MSA Intra Buffer Variability MSA Mean [Std Dev of Buffer Area] (acre)	MSA Inter Buffer Variability MSA Std Deviation [Buffer Area] (acre)
all MSAs	3749	134.2432	108.7585	118.8869	144.36
Atlanta	271	152.5512	112.9346	154.5027	116.89
Baltimore	88	142.3519	124.0517	130.2395	138.91
Boston	122	81.7104	37.9541	116.9234	67.67
Chicago	195	133.1570	101.4659	119.4062	157.06
Cincinnati	119	204.4641	185.5560	140.9650	179.77
Cleveland - Akron	94	164.3451	135.3040	142.5841	144.36
Dallas	224	151.0491	126.5676	128.2411	153.40
Denver - Boulder	135	137.9317	124.7992	95.9796	165.78
Detroit	130	177.0784	147.3925	140.6987	166.60
Houston	234	126.3528	106.4669	104.4205	146.66
Los Angeles – Riverside - Ventura	190	73.3082	49.1912	82.9921	88.23
Miami	64	66.4412	44.3581	76.2510	95.51
Minneapolis - St. Paul	209	231.6870	212.8346	148.8624	191.55
New York City	206	75.4649	45.0933	99.8590	92.21
Philadelphia - Trenton	181	119.7133	88.3221	123.9231	107.35
Phoenix	173	125.3527	102.7028	104.7815	129.45
Pittsburgh	178	175.7874	145.3813	156.9866	139.83
Portland	153	137.9359	112.9229	120.7735	134.80
San Diego	97	86.2608	65.9065	83.1923	115.35
San Francisco - San Jose	131	111.9393	96.3576	88.9875	138.62
Seattle	110	104.3815	82.4772	100.4655	116.03
St. Louis	208	168.6837	149.5219	121.9097	175.49
Tampa - St. Petersburg	90	84.0708	64.6226	71.3103	126.60
Washington D.C.	147	92.4470	67.0328	112.8528	113.46

Like many of the neighborhoods sampled and studied by Jacobs (1993), the Intra Buffer Variability of both length and area is also broad in this case study. Miami, Los Angeles – Riverside – Ventura, New York City, and Tampa – St. Petersburg illustrate a lower MSA Intra Buffer Variability of Length and Area, which suggests greater consistency between the measures within their buffers. Denver - Boulder, Phoenix, San Diego, and San Francisco – San Jose exhibit a higher MSA Intra Buffer Variability of Length with a lower MSA Intra Buffer Variability of Area, which suggests relative consistency among the measures of area despite variability between the measures of length. Boston, Baltimore, Detroit, and Philadelphia - Trenton exhibit a lower MSA Intra Buffer Variability of Length with a higher MSA Intra Buffer Variability of Area, which suggests relative consistency among the measures of length despite variability between the measures of area.

When the measures of scale are averaged by buffer and the variability between them is evaluated, the measures differ considerably. Given that buffers are sampled from the metropolitan center as well as the periphery of each MSA, not surprisingly, the Inter Buffer Variability of Scale, or the variability in the average scale between buffers, is broad for both Buffer Length and Buffer Area; but unexpectedly, the Intra Buffer Variability, or the variability in the scale of road segments and blocks within each of the buffers, is also broad.

As a comparison, Inter Buffer Variability is graphed in relation to the Intra Buffer Variability of Scale, with their associated means used to delineate those trending higher or lower than average.



**Figure 4:** Graph of MSA Mean of Inter Buffer Variability for Scale in relation to MSA Mean of Intra Buffer Variability for Scale, for the Randomly Sampled Set

In considering the variability of length, both between and within each buffer, the graph illustrates four distinct quadrants for these selected MSAs (Figure 4A). Denver – Boulder exhibits a higher MSA Inter Buffer Variability of Length and a higher MSA Intra Buffer Variability of Length, with Cleveland – Akron, Minneapolis – St. Paul, Phoenix, San Diego, San Francisco – San Jose, and Seattle similar. Contrastingly, Boston exhibits a lower MSA Inter Buffer Variability of Length with a lower MSA Intra Buffer Variability of Length with Baltimore, Miami, New York City, Philadelphia – Trenton, Tampa – St. Petersburg, and Washington D.C. similar. Atlanta, Cincinnati, Dallas, Houston, Pittsburgh, Portland, and St. Louis exhibit a lower Inter Buffer Variability of Length with a higher Intra Buffer Variability of Length. Interestingly, only Chicago exhibits high variability in the average measures of length between buffers with more consistency of length within its buffers.

In considering the variability of Area, both between and within each buffer, the graph again illustrates four distinct quadrants (Figure 4B). For these selected MSAs, Chicago, Cincinnati, Cleveland – Akron, Dallas, Detroit, Minneapolis – St. Paul, and St. Louis exhibit a higher MSA Inter Buffer Variability of Area and a higher MSA Intra Buffer Variability of Area, illustrating variability both between and within the measures of the buffers; in contrast, Los Angeles – Riverside – Ventura, Miami, New York City, Phoenix, San Diego, San Francisco – San Jose, Seattle, Tampa – St. Petersburg, and Washington D.C. exhibit a lower MSA Inter Buffer Variability of Area and a lower MSA Intra Buffer Variability, illustrating greater consistency both between and within the measures of the buffers. Atlanta, Baltimore, Boston, Philadelphia – Trenton, Pittsburgh, and Portland exhibit a higher MSA Intra Buffer Variability with a lower

Inter Buffer Variability of Area. Interestingly, only Denver – Boulder and Houston exhibit high variability in the average measures of area between buffers with more consistency of area within them.

## 7. Statistical Inferences

Measures for the Inter Buffer Variability, as calculated from these randomly sampled neighborhoods, capture the expected variation within this selected set of MSAs. Miami exhibits consistency among the average measures of its neighborhoods while Denver – Boulder exhibits variation, and neither is surprising given the gridded street structure of Miami juxtaposed to the varied geography of Denver – Boulder.

Measures for the Intra Buffer Variability capture far more variation and unpredictability in the measures of scale than originally anticipated, particularly given the examples often analytically studied as extremes in the literature. Chicago exhibits consistency in the length of road segments within its neighborhoods though there is greater variation in its block size despite its regularized plan. Contrastingly, Minneapolis – St. Paul exhibits variation in its road segment length and its block area.

When Inter Buffer Variability is plotted against Intra Buffer Variability for both length and area, several probabilities are suggested. First, the measures for length and area operate independently in several MSAs, despite their strong correlation (Peponis et al. 2007). Second, only a few of the 24 MSAs demonstrate a higher Inter Buffer Variability with a lower Intra Buffer Variability so the chance of encountering greater variability in the measures of scale between buffers and yet not within them is quite low. Lastly, almost twice as many MSAs exhibit low variability between their buffers while the variability within each is relatively evenly distributed around the mean. In summary, many buffers within these MSAs behave as originally perceived, despite perhaps the extraordinary variation amid the scale of the road segments and blocks within them.

## 8. Implications

As introduced, Inter and Intra Buffer Variability can be used to capture the consistency and/or variation in any number of measures. Inter Buffer Variability describes the differences *between* the average measures of each locality while Intra Buffer Variability describes the differences in the measures *within* each locality, regardless of the variable analyzed. Furthermore, the method introduced for sampling localities randomly ensures statistical significance in the results and can be utilized in other analyses.

## References

- Abler, R. and Adams, J. (1976), *A Comparative atlas of America's great cities: twenty metropolitan regions*, Washington DC, Association of American Geographers.
- Cervero, R. and Gorham, R. (1995), 'Commuting in transit versus automobile neighborhoods'. In *Journal of American Planning Association*, Vol. 61.
- Crane, R. and Crepeau, R. (1998), 'Does neighborhood design influence travel?: a behavioral analysis of travel diary and GIS data.' In *Transportation Research Part D*, Vol. 3, p.225-238.
- Doxiadis, C. (1968), *Ekistics: an Introduction to the science of human settlements*, New York, Oxford University Press.
- Frank, L., Kerr, J., Chapman, J. and Sallis, J. (2007), 'Urban form relationships with walk trip frequency and distance among youth'. In *American Journal of Health Promotion*, Vol. 21 (4 Suppl), p.1-8.
- Handy, S., Paterson, R. and Butler, K. (2003), *Planning for Street Connectivity: getting from here to there*, American Planning Association, Planning Advisory Service.
- Hess, P. (1997), 'Measures of Connectivity'. In *Places*, Vol. 11, p.58-65.
- Jacobs, A. (1993), *Great streets*, Cambridge, MIT Press.
- Jo, S. (1998), 'Spatial Configuration and Built Form', In *Journal of Urban Design*, Vol. 3, p.285-302.
- Krier, L. (1976), 'Projects on the City'. In *Lotus International*, Vol. 11, p.73-93.
- New York, N.Y. (1929), *New York and its Environs: a graphic regional plan* (Vol.1). New York: Regional Plan of New York.
- Ozbil, A. and Peponis, J. (2007), 'Modeling street connectivity and pedestrian movement according to standard GIS street network representations'. In: Kubat, A. S., Ertekin, O., Guney, Y. I. and Eyuboglu, E. (eds), *Proceedings of the Sixth International Space Syntax Symposium*, Istanbul: ITU Faculty of Architecture.
- Passonneau, J. and Wurman, R. (1966), *Urban atlas: 20 American cities; a communication study notating selected urban data at a scale of 1:48,000*, Cambridge, MIT Press.
- Peponis, J., Allen, D., Haynie, S., Scoppa, M., and Zhang, Z. (2007), 'Measuring the Configuration of Street Networks'. In: Kubat, A. S., Ertekin, O., Guney, Y. I. and Eyuboglu, E. (eds), *Proceedings of the Sixth International Space Syntax Symposium*, Istanbul: ITU Faculty of Architecture.
- Rashid, M. (1996), 'The Plan Is the Program: Thomas Jefferson's Plan for the Rectilinear Survey of 1784.' 84th ACSA Annual Meeting. *Regional Papers*, 615-619.
- Siksna, A. (1997), 'The Effects of Block Size and Form in North American and Australian City Centres'. In *Urban Morphology*, Vol. 1, p.19-33.
- Southworth, M. and Owens, P. (1993), 'The Evolving Metropolis: Studies of Community, Neighborhood, and Street Form at the Urban Edge'. In *Journal of the American Planning Association*, Vol. 59(3).