

2012

Relationship of Body Mass Index and Activity Level with Sleep Quality Among College Women

Caroline Wuertz
Georgia State University

Yesmira Chia
Georgia State University

Jiwon Lee
Georgia State University

Follow this and additional works at: <http://scholarworks.gsu.edu/discovery>

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Wuertz, Caroline; Chia, Yesmira; and Lee, Jiwon (2012) "Relationship of Body Mass Index and Activity Level with Sleep Quality Among College Women," *DISCOVERY: Georgia State Honors College Undergraduate Research Journal*: Vol. 1, Article 14.
Available at: <http://scholarworks.gsu.edu/discovery/vol1/iss1/14>

This Article is brought to you for free and open access by ScholarWorks @ Georgia State University. It has been accepted for inclusion in *DISCOVERY: Georgia State Honors College Undergraduate Research Journal* by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

Relationship of Body Mass Index and Activity Level with Sleep Quality among College Women

Byrdine F. Lewis School of Nursing and Health Professions
 Caroline Wuertz, Yesmira Chia, Jiwon Lee
 Faculty: Shih-Yu (Sylvia) Lee, RNC, PhD



BACKGROUND

- The effect of obesity & sedentary lifestyle on health has been thoroughly researched.
- College women tend to prioritize studying over healthy sleeping habits.
- Little is known regarding Body Mass Index (BMI) and activity level in relation to sleep quality among college women.
- Determining these relationships would be beneficial in educating college women about the implications of their behaviors.

PURPOSE

- To explore the relationship between:
 - 1) BMI and sleep quality
 - 2) Physical activity and sleep quality

METHODS

- This cross-sectional correlation study uses partial data from phase I of the Psychoneuroimmunological (PNI) Biomarkers and Health Outcomes in College Women study.
- A convenience sample of 68 Georgia State University female students (mean age=26.4, SD= 7.5) completed a 7-day sleep diary including questions about physical activity.
- Physical activity level was calculated by using an average of self-reported 7-day, 5-point physical activity scale.
- Average daily activity time was obtained from a single-item self-report.
- Sleep quality measured by the Pittsburgh Sleep Quality Index (PSQI).
- BMI was calculated based on self-reported height and weight.

RESULTS

- 67.6 % of the study participants are poor sleepers (PSQI > 5).
- 23.5 % of the study participants are overweight or obese (BMI ≥ 25).
- Most participants (91.2%) self-reported their overall health as good to excellent.
- Although not statistically significant, the mean activity level was negatively correlated with BMI ($r = -.10$).
- Participants with high mean activity level fall asleep quicker than lower mean activity level ($r = -.28, p < .05$).
- Participants with normal BMI had better sleep efficiency ($r = .29, p < .05$).
- The t-tests revealed that poor sleepers reported significantly lower physical activity levels and daily physical activity time than good sleepers but not a significant difference in BMI.

Table 1. Sample Characteristics (N=68)

Variables	Mean (SD)	Frequencies (%)
Age	26.4 (7.5)	
Ethnicity		
White		28 (41.2)
African American		15 (22.1)
Asian		12 (17.6)
Hispanic		8 (11.8)
Other		5 (7.4)
Health Condition		
Fair		6 (8.8)
Good		25 (36.8)
Very Good		29 (42.6)
Excellent		8 (11.8)
BMI*		
Underweight (<18.5)		1 (1.5)
Normal (18.5-24.9)		47 (69.1)
Overweight (25-29.9)		7 (10.3)
Obese (≥30)		9 (13.2)
Activity (min) (n=63)	44.50 (20.02)	
Activity level (n=58)		
Basic		27 (39.7)
Mild		34 (50)
Moderate		7 (10.3)
PSQI	7.2 (2.8)	
PSQI >5		58 (67.6)

* n = 64

Table 2. Correlations between Sleep, BMI, and Activity Level

	PSQI Global	PSQI SL	PSQI SE	BMI	Activity Level	Activity (min)
PSQI Global	--					
SL ^a	.64**	--				
SE ^b	-.66**	-.46**	--			
BMI	.07	-.02	.29*	--		
Activity Level	-.24*	-.28*	-.13	-.10	--	
Activity (min)	-.27*	-.05	-.25	-.14	.28*	--

^a Sleep Latency, ^b Sleep Efficiency

** Correlation is significant at the 0.01 level

* Correlation is significant at the 0.05 level

Table 3. T-test for PSQI vs. Activity Level and BMI

	Poor Sleeper Mean (SD)	Good Sleeper Mean (SD)	T-test
Activity Level ^a (n = 68)	3.03 (.55)	3.36 (.71)	-2.028*
Activity (minutes) (n = 60)	41.09 (19.98)	51.84 (18.51)	1.98*
BMI (n = 64)	24.16 (4.3)	23.47 (6.2)	.445

*p < .05

^a 1 = no activity; 2 = basic activity; 3 = mild activity (e.g. easy walking);

4 = moderate activity (e.g. easy aerobics); 5 = strenuous activity (e.g. running)

CONCLUSIONS

- The results indicated a high prevalence of sleep problems in college women.
- Our findings demonstrated that poor sleep was significantly associated with low physical activity but not associated with BMI.
- The disconnection between BMI and sleep quality could be explained by both metabolic resiliency and small sample size.
- Given the high percentage of students with poor sleep quality (67.6%), further investigation with larger sample size is necessary to examine possible factors influencing poor sleep quality in this population.