Georgia State University ScholarWorks @ Georgia State University

Mathematics Theses

Department of Mathematics and Statistics

5-3-2007

Omnibus Tests for Comparison of Competing Risks with Covariate Effects via Additive Risk Model

Duytrac Vu Nguyen

Follow this and additional works at: http://scholarworks.gsu.edu/math theses

Recommended Citation

Nguyen, Duytrac Vu, "Omnibus Tests for Comparison of Competing Risks with Covariate Effects via Additive Risk Model" (2007). *Mathematics Theses.* Paper 25.

This Thesis is brought to you for free and open access by the Department of Mathematics and Statistics at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Mathematics Theses by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

OMNIBUS TESTS FOR COMPARISON OF COMPETING RISKS WITH COVARIATE

EFFECTS VIA ADDITIVE RISK MODEL

by

DUYTRAC VU NGUYEN

Under the direction of Yichuan Zhao

ABSTRACT

It is of interest that researchers study competing risks in which subjects may fail from any

one of K causes. Comparing any two competing risks with covariate effects is very important in

medical studies. This thesis develops omnibus tests for comparing cause-specific hazard rates

and cumulative incidence functions at specified covariate levels. In the thesis, the omnibus tests

are derived under the additive risk model, that is an alternative to the proportional hazard model,

with by a weighted difference of estimates of cumulative cause-specific hazard rates.

Simultaneous confidence bands for the difference of two conditional cumulative incidence

functions are also constructed. A simulation procedure is used to sample from the null

distribution of the test process in which the graphical and numerical techniques are used to detect

the significant difference in the risks. A melanoma data set is used for the purpose of illustration.

INDEX WORDS:

Cause-specific hazard rates, Additive risk model, Cumulative

incidence function, Confidence bands, Competing risks.

OMNIBUS TESTS FOR COMPARISON OF COMPETING RISKS WITH COVARIATE EFFECTS VIA ADDITIVE RISK MODEL

by

DUYTRAC VU NGUYEN

A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of

Master of Science

in the College of Arts and Sciences

Georgia State University

Copyright by
Duytrac Vu Nguyen
2007

OMNIBUS TESTS FOR COMPARISON OF COMPETING RISKS WITH COVARIATE EFFECTS VIA ADDITIVE RISK MODEL

by

DUYTRAC VU NGUYEN

Major Professor: Dr. Yichuan Zhao Dr. Yu-Sheng Hsu Committee: Dr. Xu Zhang

Dr. Gengsheng (Jeff) Qin

Electronic Version Approval:

Office of Graduate Studies College of Arts and Sciences Georgia State University May 2007

ACKNOWLEDGEMENTS

I would like to acknowledge those who have helped me in the completion of this thesis. First of all, I must thank Dr. Zhao who had been an extraordinary advisor, a great professor, and a great supporter. This thesis cannot be finished without him. It has been a great learning experience for me to work and take his classes for almost two years.

I would like to acknowledge the other members of my committee, Dr. Yu-Sheng Hsu, Dr. Gengsheng (Jeff) Qin, and Dr. Xu Zhang for taking the time to read this thesis and provide useful comments.

I'd also like to thank all other professors in the Mathematics & Statistics Department at Georgia State University for their supports and encouragement. They have been a great helper and very generous. I thank you for everything.

Lastly, I want to acknowledge the support of my parents and my fiancé who endured this long process with me, always offering support and love.

TABLE OF CONTENTS

Acknowledgements	iv
List of Tables	vi
List of Figures	vii
List of Abbreviation	viii
Chapter One: Introduction	1
Chapter Two: Test Procedure	5
2.1 Cumulative cause-specific hazard function	5
2.2 Test Process	7
2.3 Confidence Bands	11
Chapter Three: Simulation Study	13
3.1 Simulation example (1)	15
3.2 Simulation example (2)	17
3.3 Simulation example (3)	19
Chapter Four: Application	
Chapter Five: Conclusion	32
References	34
Appendix	36

List of Tables

Table 3.1:	Powers of test for equality of conditional cause-specific hazard rates based on D_2 and D_3 at nominal level 0.05 with $z_0 = (0.5, 0.5)$, CR = 0.30	16
Table 3.2:	Powers of test for equality of conditional cause-specific hazard rates based on D_2 and D_3 at nominal level 0.05 with $z_0 = (0.5,0.5)$, CR =0.15	.18
Table 3.3:	Powers of test for equality of conditional cause-specific hazard rates based on D_2 and D_3 at nominal level 0.05 with $z_0 = (0.3,0.3)$, CR = 0.30	.19
	Statistics test under additive risk model for failure caused by disease and failure caused by other causes. H_2 and H_3 as alternative hypotheses	.21
	Statistics test under Cox regression model for failure caused by disease and failure caused by other causes. H_2 and H_3 as alternative hypotheses	.22
Table 4.3:	Risk factors in Additive risk model for failure time to death occurring from different causes of 205 malignant melanoma patients	.25
Table 4.4:	Risk factors in Cox regression model for failure time to death occurring from different causes of 205 malignant melanoma patients	.25

List of Figures

Figure 4.1a: Cumulative incidence functions plotting	23
Figure 4.1b: Hazard rate plotting	23
Figure 4.2: The test process $L(t)$ using W_I plotting	27
Figure 4.3: The test process $L(t)$ using W_2 plotting	28
Figure 4.4: The test process $L(t)$ using W_3 plotting	29
Figure 4.5: The test process $L(t)$ using W_4 plotting	30
Figure 4.6: The simultaneous confidence bands for the difference of cumulative incidence functions	31

List of Abbreviation

MM: Malignant Melanoma

CR: Censor Rate

Chapter One: Introduction

Competing risks has been arising in many applications in medicine nowadays. In competing risks people can study the cause of risk that each subject may fail in. It is important to discover which cause affects the death of the subject because it will help researchers have a good way of treatment to patients or sometimes they can design a new vaccine for a disease. Each subject is observed with a time to failure and an indicator that tells which of the competing risks causes the failure. The classic example of competing risks is competing causes of death, such as cancer, heart disease, AIDS, etc. For example, in cancer clinical trials, patients being followed up for relapse may die before relapse occurs, so relapse and death without antecedent relapse are associated as competing failure types. Another example, in clinical study that involves 205 patients operated for malignant melanoma, 71 patients died in which 57 patients were recorded as having died of the disease and 14 patients who died from causes unrelated to the disease during the follow-up (Andersen et al., 1993). This thesis develops a method that is called omnibus tests in which competing risks is compared via additive risk model with using nonnegative weight functions. Comparison of cause-specific hazard rates and cumulative incidence functions at specified covariate levels are applied in this method.

As we consider a competing risks framework in which subjects are at risk from k types of failure and covariate measurements on each subject are available. We will propose a testing that provides graphical and numerical methods for comparing any two of the k cause-specific hazard rates, or cumulative incidence functions, at a specified covariate level. Suppose each subject has an underlying failure time X that may be subject to censoring. We assume there is an existence of the latent failure time T_i corresponding to each failure

type j=1,...,k. Then the observed time of failure is given by $X=\min_j T_j$, which may be right censored. When X is uncensored, the cause of failure $\delta \in \{1,...,k\}$ is observed along with $p-vector\ Z$ representing the covariate information. We are interested in focusing on the conditional cause-specific hazard rate $\lambda_j(t\mid z_0)=\lim_{\Delta t\to 0}P(t\leq X\leq t+\Delta t,\delta=j\mid X\geq t,Z=z_0)/\Delta t$. In other words, $\lambda_j(t\mid z_0)$ is the instantaneous of failure of type j given z_0 and in the presence of the other failure types. In the presence of dependent competing risks, the conditional cumulative incidence function is defined by $F_j(t\mid z_0)=P(X\leq t,\delta=j\mid Z=z_0)$.

According to McKeague, Gilbert, and Kanki (2001), a proposal of omnibus tests for comparison of competing risks with covariate effects has been developed under Cox proportional hazards model (Cox, 1972). Also, nonparametric tests that allow censoring and dependent competing risks but no covariate have been considered by Aly, Kochar, and McKeague (1994), Lam (1998), among others. Although the Cox model has been considered as a major tool for the regression analysis of censored survival data, the proportional hazards assumption may not be appropriate to some data analyses. Therefore, it is of interest for us to investigate an alternative approaches to modeling the association between failure times and risk factors through hazard functions. Additive risk model in various forms which has been developed and successfully utilized by numerous authors (Aalen, 1980, Cox and Oakes, 1984; Lin and Ying, 1994) is one such alternative that has been used to describe different aspects of these associations, and it is suitable in many applications.

Under the additive risk model, we specify each $\lambda_j(t \mid z_o)$ and assume that the censoring is conditionally independent of the latent failure times given Z. From these

assumptions, we develop a graphical method, along with a formal procedure, for testing the null hypothesis

$$H_0: \lambda_1(t \mid z_0) = \lambda_2(t \mid z_0), \ 0 \le t \le \tau,$$

where z_0 is a specified *p*-vector of covariate levels and $[0, \tau]$ is the time interval of interest. Then, the following hypotheses will be considered:

$$H_1: \lambda_1(t \mid z_0) \neq \lambda_2(t \mid z_0), \ 0 \le t \le \tau$$

$$H_2: F_1(t \mid z_0) \le F_2(t \mid z_0), \ 0 \le t \le \tau$$

$$H_3: \lambda_1(t \mid z_0) \le \lambda_2(t \mid z_0), \ 0 \le t \le \tau$$

with strict inequality for some $t \in [0, \tau]$ in H_2 and H_3 . The hypothesis H_0 is equality of the corresponding cumulative incidence functions over $[0, \tau]$, because $F_j(t \mid z_o) = \int_0^t S_X(u \mid z_o) \lambda_j(u \mid z_o) du$ (Kalbfleisch and Prentice, 1980; Gay, 1988), where $S_X(t \mid z_o) = P(X > t \mid Z = z_o)$ is the conditional survival function of X. The hypotheses H_2 and H_3 are alternative expressing that cause 2 is more serious than cause 1, with H_3 being the more restrictive alternative. H_2 is appropriate for a comparison in terms of absolute risk and H_3 in terms of risks intensity. We develop omnibus tests that are consistent against all departures from H_0 in the directions of H_1 , H_2 , and H_3 .

The rest of this thesis is organized as follows. Chapter two includes a brief introduction to the test procedure in which a formula will be proposed for using comparison based on a comparison of the cumulative cause-specific hazard estimators at the specified covariate level z_0 . A sampling from the null distribution of the test process that is used to simulate and detect departure from the null hypothesis is discussed in this chapter. Choice of

weight process that will help the test leads to graphical procedure is also covered. Confidence band for difference of two cumulative incidence functions is constructed in Chapter two. Simulation results, comparing cause-specific hazard rates and cumulative incidence functions are presented in Chapter three. In Chapter four, we apply the proposed test to the real malignant melanoma data for comparing the rate of death. The thesis concludes with a summary of results and potential research.

Chapter Two: Test Procedure

2.1 Cumulative cause-specific hazard function

As shown in Lin and Ying (1994), the regression coefficient estimator in the semiparametric additive risk model has an explicit form, which leads to a simpler mathematical solution in comparison to that of the Cox model. In contrast to the Cox model, the additive risk model specifies the hazard function for a covariate vector Z in the form of

$$\lambda_i(t \mid Z) = \lambda_{0i}(t) + \beta_i^T Z(t), \tag{2.1}$$

where $\lambda_{0j}(.)$ is an unspecified baseline hazard function and β_j is an unknown p-vector of regression parameter for the jth cause of failure. Here, Z(t) is a p-vector of possibly timevarying covariates and to be time independent. Lin and Ying (1994) applied the standard counting process martingale techniques and showed that the estimator is asymptotically normal with mean β_j and a variance-covariance matrix.

Let C denote the censoring time. The competing risks model data are assumed to be given by n independent replicated of $(\widetilde{X}, \widetilde{\delta}, Z)$, where $\widetilde{X} = \min(X, C)$, $\widetilde{\delta} = \delta I(X \le C)$, and I(.) is indicator function. We also assume that C is conditionally independent of $T_l, ..., T_k$ given covariates Z(t). The latent failure time T_j do not have to be independent, but we do require that $P(T_j = T_l) = 0$ for $j \ne l$. The cause of failure time $\delta = j$ when $X = T_j$. Let $\Delta_{ji} = I(\widetilde{\delta}_i = j)$, and $N_{ji}(t) = \Delta_{ji}I(\widetilde{X}_i \le t)$ (i = 1, ..., n) be a counting process for indicating whether an event of type j has been observed for ith subject by time t. Let $Y_i(t) = I(\widetilde{X}_i \ge t)$ denote the predictable process indicating whether or not the ith subject is at risk just before

time t. Let τ satisfy $P(X_i > \tau) > 0$. Lin and Ying (1994) proposed the following estimation function

$$U_{j}(\beta) = \sum_{i=1}^{n} \int_{0}^{\tau} Z_{i}(t) \{ dN_{ji}(t) - Y_{i}(t) d\hat{\Lambda}_{0j}(\beta_{j}, t) - Y_{i}(t) \beta_{j}^{T} Z_{i}(t) dt \},$$
 (2.2)

where

$$\hat{\Lambda}_{0j}(\hat{\beta},t) = \int_{0}^{\tau} \frac{\sum_{i=1}^{n} \left\{ dN_{ji}(u) - Y_{i}(u) \hat{\beta}_{j}^{T} Z_{i}(u) du \right\}}{\sum_{i=1}^{n} Y_{j}(u)}.$$

(2.2) is equivalent to

$$U_{j}(\beta) = \sum_{i=1}^{n} \int_{0}^{\tau} \{Z_{i}(t) - \overline{Z}(t)\} \{dN_{ji}(t) - Y_{i}(t)\beta_{j}^{T} Z_{i}(t)dt\},$$
 (2.3)

where

$$\overline{Z}(t) = \frac{\sum_{j=1}^{n} Y_{j}(t) Z_{j}(t)}{\sum_{j=1}^{n} Y_{j}(t)}.$$

The regression coefficients are estimated by solving the equation $U_{j}(\hat{\beta}) = 0$. That is

$$\left[\sum_{i=1}^{n} \int_{0}^{\tau} Y_{i}(t) \{Z_{i}(t) - \overline{Z}(t)\}^{\otimes 2} dt\right] \hat{\beta} = \left[\sum_{i=1}^{n} \int_{0}^{\tau} \{Z_{i}(t) - \overline{Z}(t)\} dN_{ji}(t)\right]. \tag{2.4}$$

The resulting estimator of β is

$$\hat{\beta}_{j} = \left[\sum_{i=1}^{n} \int_{0}^{\tau} Y_{i}(t) \{ Z_{i}(t) - \overline{Z}(t) \}^{\otimes 2} dt \right]^{-1} \left[\sum_{i=1}^{n} \int_{0}^{\tau} \{ Z_{i}(t) - \overline{Z}(t) \} dN_{ji}(t) \right], \tag{2.5}$$

where $a^{\otimes 2} = aa'$. The confidence region for the regression parameter β_j has been developed by (Zhao and Hsu, 2005) in which the empirical likelihood method under the semiparametric additive risk model with right censoring had been used.

Under model (2.1), for each j, the counting process $N_{ji}(.)$ can be decomposed uniquely so that for every ith subject at t, $N_{ji}(t) = M_{ji}(t) + \int_0^t Y_i(u) d\Lambda_j(u; Z_i)$, where $M_{ji}(\cdot)$ is a local square integrable martingale (Lin and Ying, 1994). The jth cumulative cause-specific hazard function for T_j given $Z_i = z_0$ takes the form

$$\Lambda_{j}(t \mid z_{0}) = \Lambda_{oj}(t) + \int_{0}^{t} \beta_{j}^{T} z_{0} du, \qquad (2.6)$$

where

$$\hat{\Lambda}_{oj}(t) = \frac{1}{n} \sum_{i=1}^{n} \int_{0}^{t} \frac{1}{Y(u)} dN_{ji}(u) - \int_{0}^{t} \hat{\beta}_{j}^{T} \overline{Z}(u) du , \qquad (2.7)$$

where β_j values are computed from (2.5) and $\overline{Y}(t) = \frac{1}{n} \sum_{j=1}^{n} Y_j(t)$.

2.2 Test process

We use an approach test that is based on a comparison of the cumulative causespecific hazard estimators at the specified covariate level z_0 , using general predictable locally bounded nonnegative weight processes W(.),

$$L(t) = \int_{0}^{t} W(u) \left(\hat{\Lambda}_{2}(du \mid z_{0}) - \hat{\Lambda}_{1}(du \mid z_{0}) \right).$$
 (2.8)

The weight process provides a flexible way to control the relative importance attached to differences to the cause-specific hazards at different times and is useful for controlling instability in the tails. Thus, it is important to choose the weight function so that the function (2.8) gives a better and nominal level. Aly, Kochar, and McKeague (1994) used weight function for comparing cumulative incidence functions and cause-specific hazard rates, but

the covariate was absent. Then, McKeague, Gilbert, and Kanki (2001) applied weight functions for comparing cause-specific hazard rates and cumulative incidence functions at specified covariate levels. Note that the various choice of weight function was under Cox hazards proportional model.

There are many ways to choose the weight process. First, the simplest choice is $W_1(t) = n^{1/2}$, which reduces the test process to the normalized difference of the estimated cumulative cause-specific hazard functions,

$$L(t) = \sqrt{n} \left(\hat{\Lambda}_2(t \mid z_0) - \hat{\Lambda}_1(t \mid z_0) \right). \tag{2.9}$$

This choice is good and easy to interpret, but the variance of L(t) increases with t, so it is preferable to use a decreasing weight process that gives less weight to the tail, such as

$$W_2(t) = \sum_{i=1}^{n} I(\tilde{X}_i \ge t) / \sqrt{n} , \qquad (2.10)$$

where I(.) = 1 when $(X_i \wedge C_i \ge t)$, otherwise I(.) = 0.

Another choice that gives more sophisticated weight process and has a similar effect under the additive risk model is

$$W_3(t) = \hat{S}_X (t - |z_0|^{1/2} \left(\frac{2}{S^{(o)}(t)}\right)^{-1/2}, \qquad (2.11)$$

where

$$S^{(o)}(t) = \sum_{i=1}^{n} Y_i(t),$$

and $\hat{S}_X(t \mid z_0) = \exp\left\{-\sum_{j=1}^k \hat{\Lambda}_j(t \mid z_0)\right\}$ is the conditional survival function of X. In this case,

the asymptotic distribution of L(t) is the additive hazards model based on the estimator of a relatively simple form, and the variance function of V(t) simplifies to the conditional

cumulative incidence function $F(t | z_0)$. For k = 2 and no covariates, $W_3(t)$ reduces to a weight process considered by (Aly et al., 1994) and makes the test statistics asymptotically distribution free.

The last relatively simple choice is $W_4(t) = n^{1/2} \hat{S}_X(t - |z_0)$, which gives the normalized difference of the estimated cumulative incidence functions,

$$L(t) = \sqrt{n} \left(\hat{F}_2(t \mid z_0) - \hat{F}_1(t \mid z_0) \right), \tag{2.12}$$

where
$$F_{j}(t \mid z_{o}) = \int_{0}^{t} \hat{S}_{X}(u - |z_{0}) \hat{\Lambda}_{j}(du \mid z_{0}).$$

We use a plotting method to demonstrate (2.8) in looking for possible departure from H_0 , with a tendency for large absolute values under H_1 , large positive values under H_2 , and an increasing H_3 . This be identity trend under from the can seen $F_j(t \mid z_o) = \int_0^t S_X(u \mid z_o) \lambda_j(u \mid z_o) du$. However, these plots can be difficult to interpret when the test process that occur even under the hypothesis of equal cause-specific hazards. McKeague, Gilbert, and Kanki (2001) showed that L(t) from (2.8) converges in distribution under H_0 to a zero mean Gaussian process provided $W(t)/n^{1/2}$ converges uniformly in probability over $[0,\tau]$ to a bounded function. Moreover, the limiting covariance is complicated. Therefore, we develop a simulation method for approximately sampling from the null distribution of L(t) under the additive risk model. This procedure was presented in Song, Jeong, and Song (1997) for finding confidence bands for survival curve without using weight function; in Shen and Cheng (1999) for confidence bands for cumulative incidence curves.

In order to sample the null distribution of the test process, we derive the limiting distribution of the cumulative hazard estimator.

Let

$$L_{j}(t) = n^{1/2} \left\{ \hat{\Lambda}_{j}(t \mid z_{0}) - \Lambda_{j}(t \mid z_{0}) \right\},$$

and $\tau = \inf\{t \ge 0; H(t) = 1\},$

where H(.) is the distribution function of the observed failure time \widetilde{X}_i .

Let

$$G(t,z_0) = \int_0^t W(u) \{z_0(u) - \overline{Z}(u)\} du, \qquad \overline{Y}(t) = \frac{1}{n} \sum_{j=1}^n Y_j(t),$$

$$C^{-1} = \frac{1}{n} \sum_{i=1}^{n} \int_{0}^{\infty} \left\{ Z_{i}(u) - \overline{Z}(u) \right\} Y_{i}(u) Z_{i}(u) du ,$$

where
$$\overline{Z}(t) = \frac{\sum_{i=1}^{n} Y_i(t) Z_i(t)}{\sum_{j=1}^{n} Y_j(t)}$$
.

Under H_0 , the form (2.8) is also equivalent to:

$$L(t) = \int_{0}^{t} W(u) (\hat{\Lambda}_{2}(du \mid z_{0}) - \hat{\Lambda}_{1}(du \mid z_{0})) =$$

$$\int_{0}^{t} W(u) (\hat{\Lambda}_{2}(du \mid z_{0}) - \Lambda(du \mid z_{0})) - \int_{0}^{t} W(u) (\hat{\Lambda}_{1}(du \mid z_{0}) - \Lambda(du \mid z_{0}))$$

Then, the process L(t) is asymptotically equivalent to the process $L_2(t) - L_1(t)$, where

$$L_{j}(t) = \frac{G'(t, z_{0})C'}{n} \sum_{i=1}^{n} \int_{0}^{\infty} \left\{ Z_{i}(u) - \overline{Z}(u) \right\} dM_{ji}(u) + \frac{1}{n} \sum_{i=1}^{n} \int_{0}^{t} \frac{W(u)}{\overline{Y}(u)} dM_{ji}(u), \qquad (2.13)$$

where
$$M_{ji}(t) = N_{ji}(t) - \int_{0}^{t} Y_{i}(u) d\Lambda_{j}(u, Z_{i})$$
.

The process to simulate L(t) is defined by $L^*(t)$ in which we replace $M_{ji}(u)$ in the first and second terms of $L_j(t)$ by $G_{ji}N_{ji}(u)$ and $\widetilde{G}_{ji}N_{ji}(u)$, respectively, where $\left\{G_{ji},\widetilde{G}_{ji}:j=1,2;i=1,...,n\right\}$ are independent standard normal variables. Realizations of $L^*(t)$ are approximate draws from the null distribution of the test process. In other words, under H_0 , the conditional distribution of $L^*(t)$ given the observed data is the same in the limit as the unconditional distribution of L(t) (McKeague, Gilbert, and Kanki, 2001). The method works essentially because $M_{ji}(t)$ has mean zero and variance $E\left\{N_{ji}(t)\right\}$.

With the sampling from the null distribution and the proposal of the test, we have formal procedures that detect departures from H_0 in the direction of H_1 , H_2 , and H_3 :

$$D_1 = \sup_{0 \le t \le \tau} \big| L(t) \big|, \qquad \qquad D_2 = \sup_{0 \le t \le \tau} L(t), \qquad \qquad D_3 = \sup_{0 \le s \le t \le \tau} \big\{ L(t) - L(s) \big\}, \text{ respectively.}$$

2.3 Confidence Bands

In this section, we construct confidence bands for the difference of two conditional cumulative incidence functions since confidence intervals are used for inference or for making statistical statements about estimates. Gray (1988), Shen and Cheng (1999), developed the confidence bands for cumulative incidence function for each type of failure. However, McKeague, Gilbert, and Kanki, (2001) constructed confidence bands for the difference in two conditional cumulative incidence functions using the weight function W_4 under Cox regression model. Consider the difference $\delta(t) = F_2(t \mid z_0) - F_1(t \mid z_0)$ can be obtained from the L(t) (2.12) in which the weight function W_4 is used. Since the process $n^{1/2}(\hat{\delta}(t) - \delta(t))$ converges in distribution to the null limiting distribution of L(t), we can use

 $\hat{\delta}(t) \equiv \hat{F}_2(t \mid z_0) - \hat{F}_1(t \mid z_0) = L(t)/n^{1/2}$ to estimate the difference of two conditional cumulative incidence functions. The earlier Monte Carlo procedure based on $L^*(t)$ can be used to estimate an upper $\alpha/2$ – quantile, $c_{\alpha/2}(t)$, of the limiting distribution of |L(t)|. An approximate $100(1-\alpha)\%$ pointwise confidence band for $\delta(t)$ is given by $\hat{\delta}(t) \pm c_{\alpha/2}(t)n^{-\frac{1}{2}}$. A simultaneous band for $\delta(t)$ can be found by suitably scaling the pointwise band. For example, we can use $\hat{\delta}(t) \pm ac_{\alpha/2}(t)n^{-\frac{1}{2}}$, $t \in [0,\tau]$,

where $a \ge 1$ and is the critical value of $\sup_{0 \le t \le \tau} \left| \frac{\sqrt{n} \left(\hat{\mathcal{S}}(t) - \mathcal{S}(t) \right)}{c_{\alpha/2}(t)} \right| = \sup_{0 \le t \le \tau} \left| \frac{L^*(t)}{c_{\alpha/2}(t)} \right|$. Monte Carlo procedure is used to adjust the constant a to furnish the desired $100(1-\alpha)\%$ simultaneous confidence level.

Chapter Three: Simulation Study

In this chapter, we designed simulation to study the performance of the proposed test procedure. We consider if the nominal size accurately matches observed levels at moderate sample size and which weight function that gives the best performance in terms of power. We also want to know whether covariate level and censoring rate have effect on the powers of test.

First, we consider a simple additive risk model with the baseline hazard $\lambda_0(t) = \alpha_j$, j = 1,...,k where α_j is a various choices of constant, and set $\beta_1 = \beta_2 = ... = \beta_p = 1$, then model (2.1) becomes

$$\lambda_{j}(t \mid z) = \alpha_{j} + \left[Z_{1}(t) + Z_{2}(t) +, \dots, + Z_{p}(t) \right]. \tag{3.1}$$

Simply we denote it as

$$\lambda_{i}(t \mid z) = \alpha_{i} + W, \qquad (3.2)$$

where $W = Z\beta$, $Z = (Z_1, Z_2, ..., Z_p)$ covariate vector, and $\beta = (1, 1, ..., 1)^T$.

Let *T* denote failure times, *C* denote the censoring times. We simulate competing risk data sets with right censoring survival time as following steps.

1. Simulating covariates $Z_{pi}(t)$, i = 1,...,n.

Z's are drawn from uniform distribution U(0,1). Since $Z_1,Z_2,...,Z_p$ are random independent uniform variables, we randomly choose a matrix with n rows and p columns from uniform distribution as the simulated competing risk data.

2. Simulating failure time T_i , j = 1,...,k

To simulate failure time T, we assume the conditional distribution of T given Z in model (3.1) is exponential distribution with parameter $\alpha_j + W$. The survival function related with exponential distribution is derived as $S(x) = e^{-(\alpha_j + W)x}$. Suppose the hazard function for failure time T is known, say $\lambda_j(t \mid z) = \alpha_j + W$, we can derive the survival function S(t) and the distribution function F(t) for T. Since $\lambda(t) = \frac{-d \ln(S(t))}{dt} = -\frac{S'(t)}{S(t)}$, we have

$$S(t \mid z) = e^{-\int_{0}^{t} \lambda(s|z) ds} = e^{-\int_{0}^{t} (\alpha_{j} + W) ds} = e^{-(\alpha_{j} + W)t}, \qquad (3.2)$$

$$F(t) = 1 - S(t) = 1 - e^{-(\alpha_j + W)t}, \tag{3.3}$$

$$t = -\frac{\ln(1 - F(t))}{W + \alpha_j}.$$
(3.4)

Given a distribution function F(t), we can simulate failure time samples T distributed with such a cumulative function F(t). Suppose U is drawn from uniform distribution U(0,1), then the failure time T is obtained from

$$T_{j} = -\frac{\ln(1-U)}{W + \alpha_{j}} \tag{3.5}$$

3. Simulating censoring time C_i , i = 1,...,n

Censoring time C_i 's are drawn from exponential distribution with decay parameter adjusted so 15-30% of the observations were censored in $[0,\tau]$.

Finally, the simulated observations from the additive hazard risk model (3.2) are the triples $\operatorname{as}\left(\widetilde{X}_{i},\widetilde{\delta}_{i},W_{p}\right),\ i=1,\ldots,n$, where $\widetilde{X}=\min(X,C)$, $\widetilde{\delta}=\delta I(X\leq C)$ and $\delta=j$ when $X=T_{j}$.

3.1 Simulation Example (1)

In this simulation example, we generate a competing risk data set with censoring rate (CR) 0.30, a p = 2 dimensional covariate Z. We set covariate level $z_0 = (0.5,0.5)$, and the sample size n = 50, 100, 200, and 400, respectively. The size and power of test based on D_2 and D_3 at the nominal 0.05 level were estimated from 1000 independent samples, with critical values obtained in each sample from 1000 realizations of $L^*(t)$.

Since we assumed the alternative hypotheses H_2 and H_3 from the proposal test that are expressing the notion of cause (risk) 2 more serious than cause 1, we choose $\alpha_1 = 1$ and various choices of $\alpha_2(1, 1.5, 2.0, 2.5, 3.0)$ in order to generate T from (3.5). Then we test all weight functions, W_1 , W_2 , W_3 , and W_4 , respectively.

Table 3.1

Observed levels and powers of test for equality of conditional cause-specific hazard rates based on D_2 and D_3 at nominal level 0.05, $z_0 = (0.5,0.5)$, CR = 0.30.

n	$\alpha_{_j}$		D_2			D_3				
	$\alpha_{\scriptscriptstyle 1}$	α_2	W_{I}	W_2	W_3	W_4	W_{I}	W_2	W_3	W_4
50	1	1	0.047	0.047	0.046	0.047	0.050	0.052	0.054	0.052
	1	1.5	0.116	0.123	0.132	0.120	0.131	0.139	0.137	0.133
	1	2.0	0.257	0.273	0.271	0.263	0.261	0.268	0.261	0.258
	1	2.5	0.368	0.387	0.393	0.323	0.377	0.383	0.377	0.330
	1	3.0	0.539	0.564	0.557	0.543	0.509	0.525	0.551	0.522
	1	1	0.057	0.060	0.064	0.059	0.054	0.061	0.061	0.060
100	1	1.5	0.206	0.215	0.216	0.207	0.201	0.209	0.208	0.206
100	1	2.0	0.417	0.430	0.441	0.414	0.382	0.397	0.393	0.371
	1	2.5	0.558	0.568	0.588	0.544	0.549	0.557	0.563	0.539
	1	3.0	0.798	0.884	0.814	0.804	0.747	0.786	0.784	0.740
	1	1	0.049	0.057	0.057	0.053	0.050	0.052	0.052	0.052
	1	1.5	0.283	0.282	0.295	0.233	0.286	0.288	0.296	0.269
200	1	2.0	0.614	0.622	0.631	0.615	0.575	0.600	0.589	0.562
	1	2.5	0.865	0.866	0.875	0.866	0.837	0.876	0.866	0.858
	1	3.0	0.953	0.966	0.968	0.957	0.926	0.935	0.938	0.916
			•						•	
400	1	1	0.057	0.057	0.059	0.051	0.059	0.057	0.059	0.047
	1	1.5	0.427	0.439	0.438	0.419	0.392	0.426	0.406	0.402
	1	2.0	0.860	0.867	0.862	0.856	0.839	0.853	0.853	0.851
	1	2.5	0.991	1.000	0.996	0.994	0.986	1.000	0.988	0.970

The results in Table 3.1 show that the observed levels for D_2 and D_3 quite accurately match the 0.05 nominal level of the test. Since we simulate failure times from (3.5) and give value α_1 and α_2 that represent risk-1 and risk-2, respectively, the failure time of risk-2 is always bigger than the failure time of risk-1. Moreover, if the value of α_2 is much bigger

than the value of α_1 , then the powers of test is larger. That means the test statistics is much significant.

In the simulation, we find that sample size and weight functions also have appreciable effect on accuracy. For example, if the sample size is increasing, the powers of test are also increasing. For example, the powers of test of sample size n = 200 are bigger than the powers of test of sample size n = 100 (cf., Table 3.1). Especially, when sample size n = 400 and $\alpha_2 = 2.5$, the power of test tends to be one. The weight functions W_2 and W_3 gave better performance overall than W_1 and W_4 in terms of power. This makes sense because both W_1 and W_4 do not down weight observations in the tail, where there tends to be sharp increase in the variance of the cumulative baseline hazard estimate.

3.2 Simulation Example (2)

In this simulation example we want to consider if censoring rate have effect on powers of test. Doing the same as simulation in example (1), we generate a competing risk data set with setting censoring rate 0.15, two covariates, covariate level $z_0 = (0.5,0.5)$, and the sample size n = 50, 100, 200, respectively.

The results, reported in Table 3.2 show that the censoring rate has effect on powers of test. As the censoring rate decreasing, the powers of test seem to be increasing, but appears not so much difference.

Table 3.2 Observed levels and powers of test for equality of conditional cause-specific hazard rates based on D_2 and D_3 at nominal level 0.05, $z_0 = (0.5,0.5)$, CR = 0.15.

n	CR	a	\mathcal{E}_j		1	O_2		D_3			
		$\alpha_{\scriptscriptstyle 1}$	α_2	W_1	W_2	W_3	W_4	W_{I}	W_2	W_3	W_4
		1	1	0.047	0.047	0.046	0.047	0.050	0.052	0.054	0.052
		1	1.5	0.116	0.123	0.132	0.120	0.131	0.139	0.137	0.133
	0.30	1	2.0	0.257	0.273	0.271	0.263	0.261	0.268	0.261	0.258
		1	2.5	0.368	0.387	0.393	0.323	0.377	0.383	0.377	0.330
50		1	3.0	0.539	0.564	0.557	0.543	0.509	0.525	0.551	0.522
50											
		1	1	0.059	0.054	0.060	0.057	0.057	0.056	0.054	0.059
		1	1.5	0.142	0.149	0.144	0.137	0.136	0.144	0.147	0.143
	0.15	1	2.0	0.285	0.290	0.293	0.288	0.269	0.274	0.273	0.265
		1	2.5	0.412	0.427	0.420	0.418	0.391	0.416	0.419	0.412
		1	3.0	0.540	0.575	0.578	0.562	0.516	0.565	0.560	0.554
		1	1	0.057	0.060	0.064	0.059	0.054	0.061	0.061	0.060
		1	1.5	0.206	0.215	0.216	0.207	0.201	0.209	0.208	0.206
	0.30	1	2.0	0.417	0.430	0.441	0.414	0.382	0.397	0.393	0.371
		1	2.5	0.558	0.568	0.588	0.544	0.549	0.557	0.563	0.539
100		1	3.0	0.798	0.884	0.814	0.804	0.747	0.786	0.784	0.740
100			П	Т	Т		T			Т	
		1	1	0.061	0.059	0.061	0.059	0.068	0.064	0.064	0.064
		1	1.5	0.210	0.222	0.217	0.209	0.208	0.215	0.214	0.208
	0.15	1	2.0	0.440	0.445	0.444	0.443	0.376	0.415	0.396	0.408
		1	2.5	0.627	0.639	0.652	0.643	0.588	0.620	0.624	0.616
		1	3.0	0.802	0.891	0.821	0.810	0.750	0.788	0.790	0.784
				0.040	0.0==	0.055	0.053	0.050	0.050	0.050	0.050
		1	1	0.049	0.057	0.057	0.053	0.050	0.052	0.052	0.052
	0.20	1	1.5	0.283	0.282	0.295	0.233	0.286	0.288	0.296	0.269
	0.30	1	2.0	0.614	0.622	0.631	0.615	0.575	0.600	0.589	0.562
		1	2.5	0.865	0.866	0.875	0.866	0.837	0.876	0.866	0.858
200		1	3.0	0.953	0.966	0.968	0.957	0.926	0.935	0.938	0.916
200		1	1	0.040	0.045	0.05	0.052	0.054	0.042	0.044	0.052
		1	1	0.049	0.045	0.05	0.053	0.054	0.043	0.044	0.052
	0.15	1	1.5	0.266	0.284	0.298	0.240	0.292	0.282	0.286	0.232
	0.13	1	2.0	0.617	0.626	0.649	0.623	0.578	0.593	0.594	0.573
		1	2.5	0.871	0.872	0.883	0.873	0.843	0.880	0.885	0.867
		1	3.0	0.961	0.986	0.977	0.967	0.942	0.956	0.961	0.950

3.3 Simulation Example (3)

Now we consider the effect of covariate level. We want to adjust the covariate level by setting $z_0 = (0.3,0.3)$; two covariates; censoring rate 0.30; the sample size n = 50 and 100. Then, we observe the result of powers of test as shown in Table 3.3. The results show the powers of test seem to be changed a little.

Table 3.3Observed levels and powers of test for equality of conditional cause-specific hazard rates based on D_2 and D_3 at nominal level 0.05, $z_0 = (0.3,0.3)$, CR = 0.30.

n				D_2			D_3				
n	Zo	α	\mathcal{L}_j			J ₂				<i>y</i> ₃	
		$\alpha_{\scriptscriptstyle 1}$	α_2	W_{I}	W_2	W_3	W_4	W_{I}	W_2	W_3	W_4
		1	1	0.047	0.047	0.046	0.047	0.050	0.052	0.054	0.052
	(0.5)	1	1.5	0.116	0.123	0.132	0.120	0.131	0.139	0.137	0.133
		1	2.0	0.257	0.273	0.271	0.263	0.261	0.268	0.261	0.258
		1	2.5	0.368	0.387	0.393	0.323	0.377	0.383	0.377	0.330
50		1	3.0	0.539	0.564	0.557	0.543	0.509	0.525	0.551	0.522
50											
		1	1	0.071	0.063	0.059	0.065	0.060	0.067	0.063	0.066
		1	1.5	0.134	0.136	0.132	0.130	0.133	0.143	0.142	0.140
	(0.3)	1	2.0	0.273	0.259	0.265	0.263	0.245	0.253	0.256	0.252
		1	2.5	0.414	0.423	0.422	0.420	0.422	0.431	0.429	0.419
		1	3.0	0.528	0.530	0.546	0.518	0.518	0.526	0.533	0.526
		1	1	0.057	0.060	0.064	0.059	0.054	0.061	0.061	0.060
		1	1.5	0.206	0.215	0.216	0.207	0.201	0.209	0.208	0.206
	(0.5)	1	2.0	0.417	0.430	0.441	0.414	0.382	0.397	0.393	0.371
		1	2.5	0.558	0.568	0.588	0.544	0.549	0.557	0.563	0.539
		1	3.0	0.798	0.884	0.814	0.804	0.747	0.786	0.784	0.740
100											
		1	1	0.053	0.055	0.052	0.050	0.051	0.051	0.054	0.054
		1	1.5	0.230	0.234	0.233	0.231	0.219	0.228	0.224	0.217
	(0.3)	1	2.0	0.410	0.412	0.408	0.404	0.401	0.408	0.404	0.402
		1	2.5	0.593	0.594	0.600	0.592	0.558	0.579	0.583	0.574
		1	3.0	0.802	0.891	0.821	0.810	0.750	0.788	0.790	0.784

Chapter Four: Application

The data consist of measurements made on patients with malignant melanoma. Each patient had their tumors removed by surgery at the Department of Plastic Surgery, University Hospital of Odense, Denmark during the period 1962 to 1977. The surgery consisted of complete removal of the tumors together with about 2.5cm of the surrounding skin. Among the measurements taken were the thickness of the tumors and whether it was ulcerated or not. These are thought to be important prognostic variables in that patients with a thick and/or ulcerated tumors have an increased chance of death from melanoma. Patients were followed until the end of 1977. There were 205 patients involving operated for malignant melanoma, 71 patients died, of whom 57 were recorded as dies of the disease and 14 from causes unrelated to the disease during the follow-up. The remaining 134 patients were censored at the closure of the study since they were still alive at that point in time (Andersen et al., 1993). For time-varying covariate effects to cause of risk were also collected, including sex, age, year entry study, the size of tumor, and ulceration status at the time of operation. The latter faction is dichotomous and is scored as "present" if the surface of the melanoma shows signs of ulcers and as "absent" otherwise. Failure time is defined as the time since operation.

For comparison, we use the proposed test to compare the rate of death by the disease and the rate of death from other causes. Let the failure caused by the disease, melanoma be risk-2 and the failure caused by other factors be risk-1. We assume risk-2 is more serious than risk-1. The follow-up time for the test is ten years. Then we have hypotheses as follow:

$$H_0: \lambda_1(t \mid z_0) = \lambda_2(t \mid z_0), \ 0 \le t \le 10,$$

where z_0 is a specified covariate levels of a four-vector covariate, i.e., sex, age, the size of tumor, and ulceration status. The alternative hypotheses will be considered:

$$H_2: F_1(t \mid z_0) \le F_2(t \mid z_0), \ 0 \le t \le 10$$

$$H_3: \lambda_1(t \mid z_0) \le \lambda_2(t \mid z_0), \ 0 \le t \le 10$$

In our application, the covariate age has mean 52 years and standard deviation 17 years, and tumor thickness covariate has mean 2.9 mm and standard deviation 3 mm. Since the prediction of the cumulative incidence based on the standardize covariates has reduced standard errors, we specified the z_0 levels as z_0 = {ulceration, 2.9 mm, male, 52 years} for the covariates ulceration, tumor size, sex, and age, respectively. Table 4.1 summarizes the result of test statistics and its p-value in which we used the simulation method for sampling from the null distribution of L(t). We also used the four weight functions, W_1 , W_2 , W_3 , and W_4 as described in Section 2.2 to detect departures from H_0 .

Table 4.1

Statistical test under additive risk model for failure caused by disease and failure resulting by other causes. H₂ and H₃ as alternative hypotheses.

		W_{I}	W_2	W_3	W_4
H_2	Test statistics	4.291	2.685	2.350	3.345
2	<i>P</i> -value	< 0.001	0	0	<0.001
H_3	Test statistics	4.552	2.941	2.531	3.601
	<i>P</i> -value	< 0.001	0	0	< 0.001

Table 4.2 Statistical test under Cox regression model for failure caused by disease and failure resulting by other causes. H_2 and H_3 as alternative hypotheses.

		W_{I}	W_2	W_3	W_4
H_2	Test statistics	5.758	3.564	3.731	4.467
	<i>P</i> -value	0.046	0.014	0.019	0.032
H_3	Test statistics	5.921	3.726	3.915	4.630
	<i>P</i> -value	0.045	0.013	0.015	0.027

From Tables 4.1 and 4.2, the p-values of W_1 , W_2 , W_3 , and W_4 in both alternative hypotheses, H_2 and H_3 show the test is very consistent and significant under both additive risk model and Cox regression model. That means the failure caused by the disease is overall higher than the failure resulting by other causes unrelated to the disease.

The graphical method is also applied in these alternative hypotheses test. Applying estimation of the cumulative incidence probability directly by the subdistribution approach (Gay, 1988; Fine and Gray, 1999) to estimate the cumulative incidence function of risk-1 and risk-2, and nonparametric Epanechnikov kernel estimates hazard rates. As shown in Figures 4.1a and 4.1b, in which the cumulative incidence of risk-2 (malignant melanoma) overally exceeded the cumulative incidence of risk-1 and the hazard rate of risk-2 is also higher than the hazard rate of risk-1. Thus, the test is satisfied for both alternative hypotheses, H_2 and H_3 .




Figure 4.1a. Malignant melanoma cohort study. No adjustment for covariates. Estimated the disease and other causes cumulative incidence functions with 95% pointwise confidence limits.




Figure 4.1b. Nonparametric Epanechnikov kernel estimates of the disease and other causes hazard rates based on the smoothed hazard rate estimate with 95% pointwise confidence limits calculated by transforming the symmetric confidence limits.

According to Andersen et al., (1993), the number death patients who involving operated for malignant melanoma is higher than the number death patients from causes unrelated to the disease during the follow-up. The results in Tables 4.1,4.2 and Figures 4.1a, 4.1b show the test satisfies the alternative hypotheses and our assumption. We also want to consider the effect of risk factors on the rate of death under the additive risk models. At the 0.05 significance level, univariable additive risk models identified tumor thickness and ulceration status are highly significant risk factors for death caused by malignant melanoma, whereas both covariates have little effect on the intensity of dying from other causes (Tables 4.3,4.4). In fact, the covariate age has the most significant influence on death caused by other factors. In addition, it is worth mentioning that certain transformations of the continuous covariates may improve the efficiency of estimators under the additive risk model (Shen and Cheng, 1999). In our application, the covariates age (mean: 52 years; standard deviation: 17 years) and tumor thickness (mean: 2.92 mm; standard deviation: 3 mm) are standardized as given in Tables 4.3 and 4.4. We also fit the cause-specific intensity function by the Cox proportional hazards model (Cox, 1972) in order to compare its results with the additive risk model. Although the magnitude of each parameter estimate appears to differ between the two model fittings, it is notable that there is good agreement between the test statistics for subject covariate effects under both models.

Table 4.3
Estimation results of risk factors in additive risk model for failure time to death occurring from different causes of 205 malignant melanoma patients.

	Covariate	Estimate	SE	Estimate/SE	P-value
	ulceration	0.05074	0.01577	3.217	0.001
Failure caused	(thickness - 2.92)/3	0.02511	0.01195	2.1014	0.0356
by melanoma	sex	0.0196	0.01457	1.3453	0.1785
	(age -52)/17	0.0105	0.00772	1.3600	0.1738
	ulceration	0.002205	0.007022	0.3141	0.7535
Failure	(thickness - 2.92)/3	0.003441	0.00524	0.6573	0.511
resulting from other causes	sex	0.004487	0.007093	0.6326	0.527
	(age -52)/17	0.0110	0.00356	3.0906	0.002


Table 4.4
Estimation results of risk factors in Cox regression model for failure time to death occurring from different causes of 205 malignant melanoma patients.

	Covariate	Estimate	SE	Estimate/SE	<i>P</i> -value
	ulceration	1.141	0.312	3.657	<0.001
Failure caused by melanoma	(thickness - 2.92)/3	0.3651	0.0925	3.946	<0.001
	sex	0.400	0.269	1.487	0.137
	(age -52)/17	0.2213	0.1411	1.567	0.117
	ulceration	0.1239	0.5842	0.212	0.832
Failure resulting from	(thickness - 2.92)/3	0.1646	0.2384	0.690	0.490
other causes	sex	0.3735	0.5455	0.685	0.493
	(age -52)/17	1.136	0.343	3.311	0.001

We applied the test to compare cause-specific hazard rates for the full model in which four covariates, ulceration status, tumor thickness, sex, and age have been included, and another model in which only the significant risk factors covariates, ulceration status, tumor thickness, and age (sex was excluded because it has little effect on the intensity of dying the disease and other causes (cf., Tables 4.3, 4.4).

Test of H_0 versus H_2 and H_3 applied at covariate level z_0 = {ulceration, 2.9 mm, male, 52 years} for the full model and z_0 = {ulceration, 2.9 mm, 52 years} for other one, indicated a significantly greater type 1 hazard rate and cumulative incidence function, and the weight functions W_2 and W_3 performed better than the weight function W_1 and W_4 as shown in Tables 4.1, 4.2 and Figures 4.2 – 4.5 in both models. Figures 4.2-4.5 also show the model without covariate sex performed better than the full model.

It is of interest to estimate the difference between the risk-1 and risk-2 cumulative incidence functions for various covariate subgroups. The confidence bands displayed in Figure 4.6 were computed via the procedure described in Section 2.3 for the same covariate subgroups as in Figures 4.2-4.5, conditions a and b, respectively. The constant value a is estimated by Monte Carlo procedure in the condition with covariate sex is 1.4, and the condition without covariate sex is 1.9 Based on the 90% simultaneous bands, we find the probability of risk-2 (malignant melanoma) is lower than the probability of risk-1 in the first two years. However, the probability of risk-2 is overall higher than the probability of risk-1 during the follow-up.







Figure 4.2. The test process L(t) (solid line) and 10 realizations of $L^*(t)$ (dash lines) for the weight process $W_I(t)$. **a.** Conditions with covariate {ulceration status, tumor thickness, sex, and age}. **b.** Conditions with covariate {ulceration status, tumor thickness, and age}.







Figure 4.3. The test process L(t) (solid line) and 10 realizations of $L^*(t)$ (dash lines) for the weight process $W_2(t)$. **a**. Conditions with covariate {ulceration status, tumor thickness, sex, and age}. **b**. Conditions with covariate {ulceration status, tumor thickness, and age}.

•







Figure 4.4. The test process L(t) (solid line) and 10 realizations of $L^*(t)$ (dash lines) for the weight process $W_3(t)$. **a.** Conditions with covariate {ulceration status, tumor thickness, sex, and age}. **b.** Conditions with covariate {ulceration status, tumor thickness, and age}.







Figure 4.5. The test process L(t) (solid line) and 10 realizations of $L^*(t)$ (dash lines) for the weight process $W_4(t)$. **a.** Conditions with covariate {ulceration status, tumor thickness, sex, and age}. **b.** Conditions with covariate {ulceration status, tumor thickness, and age}.






Figure 4.6. Ninety-five percent pointwise and 90% simultaneous confidence bands for the difference in conditional rate of death cumulative incidence functions. **a**. Conditions with covariate {ulceration status, tumor thickness, sex, and age} at level $z_0 = \{\text{ulceration, 2.9 mm, male, 52 years}\}$. **b**. Conditions with covariate {ulceration status, tumor thickness, and age} at level $z_0 = \{\text{ulceration, 2.9 mm, 52 years}\}$.

Chapter Five: Conclusion

The main purpose of this thesis was to develop omnibus tests for comparison of competing risks with covariate effects via semiparametric additive risk model that is used for alternative of the Cox regression model. It has shown how useful it is to compare cause-specific hazard rate and cumulative incidence in competing risks data. The graphical method is also applied for illustration of the comparison.

The simulation results and the applied problem verify that the additive risk model is to be fit with competing risk data. The developing of the simulation method for approximately sampling the null distribution of L(t) under the additive risk models to detect the hypotheses is consisted. Beside formal omnibus tests that are consistent against all departures from H_0 in the directions of H_1 , H_2 , and H_3 , we used graphical method as shown in Figures 4.1a - 4.1b and Figures 4.2 - 4.6 to illustrate the comparison risk-1 and risk-2. An adjustment of covariate level and censoring rate are also used in simulation of this thesis. It is very important to know which covariate has effect on the intensity of dying because it will help the researcher to determine the cause of death and its rate of death, or in the designing of a new vaccine. For example, McKeague, Gilbert, and Kanki, (2001) have mentioned about designing a new vaccine for the human immunodeficiency virus (HIV) in which they needed to compare the infection rate HIV type1 and HIV type2 with adjustment for covariate effects because in order for an HIV vaccine to be efficacious in a particular geographic region, it may be necessary to match the genotypes of the HIV antigens contained in the vaccine to the local HIV genotype that pose the greatest risk of HIV infection. This thesis has shown it to be a reliable and useful tool for comparison of competing risk data where covariate has possibly time-varying.

Future research could extend this to comparing of more than two competing risks under the additive risk model or the Cox regression model. Sun (2001) studied the nonparametric test procedures for comparing multiple cause-specific hazard rates. We will investigate the challenging problem in the future. Another area of the research could be to develop a goodness-of-fit test for the model in which a comparison between the additive risk model and the Cox regression model is considered.

References

- Aalen, O.O. (1980). A Model for Non-parametric Regression Analysis of Counting Processes. In Lecture Notes on Mathematical Statistics and Probability, 2, W. Klonecki, A. Kozek, and J. Rosiski, eds. New York: Springer-Verlag, pp. 1-25.
- Andersen, P. K., Borgan, O., Gill, R. D., and Keiding, N. (1993). *Statistical Models Based on Counting Processes*. New York: Springer-Verlag.
- Aly, E.-E., Kochar, S. C., and McKeague, I. W. (1994). Some tests for comparing cumulative incidence functions and cause-specific hazard rates. Journal of the American Statistical Association 89, 994-999.
- Cheng, M., Hall, P., and Tu, D. (2006). *Confidence bands for hazard rates under random censorship*. Biometrika 93, 357-366.
- Cox, D. R (1972). Regression models and life tables (with discussion). Journal of the Royal Statistical Society, Series B, 34, 187-220.
- Cox, D. R and Oakes, D. (1984). Analysis of Survival Data. London: Chapman & Hall.
- Fine, J. P. and Gray, R. J. (1999). A Proportional Hazards Model for the Subdistribution of a Competing Risk. JASA. 94, 496-509.
- Gray, R. J. (1988). A class of k-sample tests for comparing the cumulative incidence of a competing risk. Annals of Statistics 16, 1140-1154.
- Kalbfleisch, J. D. and Prentice, R. L. (1980). *The Statistical Analysis of Failure Time Data*. New York: Wiley.
- Klein, J.P., and Moesehberger, M.L. (2003). *Survival analysis techniques for censored and truncated data*. New York: Spring Verlag, 2nd Edition.
- Lam, K. F. (1998). A class of tests for the equality of k cause-specific hazard rates in a competing risks model. Biometrika, 85, 179-188.
- Lin, D.Y. and Ying, Z.L. (1994). Semiparametric analysis of the additive model. Biometrika, 81, 61-71.
- Lin, D.Y. and Ying, Z.L. (1997). *Additive hazards regression models for survival data*. In: proceeding of the First Seattle Symposium in biostatistics: Survival Analysis.
- Martinussen, T. and Scheike, T. H. (2006). *Dynamic Regression Models for Survival Data*. Springer.

- McKeague, I.W., Gilbert, P.B., and Kanki, P.J. (2001). Omnibus test for comparison of competing risks with adjustment for covariate effects. Biometrics 57, 818-828.
- Shen, Y. and Cheng, S.C. (1999). *Confidence bands for cumulative incidence curves under the additive risk model*. Biometrics 55, 1093-1100.
- Song, M., Jeong, D., and Song, J. (1997). *Confidence bands for survival curve under the additive risk model.* Journal of the Korean, Statistical Society, Vol. 26, No. 4.
- Sun, Y. (2001). Generalized Nonparametric Test Procedures for Comparing Multiple Cause-Specific Hazard Rates. Journal of Nonparametric Statistics, 13, 171—207.
- Zhao, Y. and Hsu, Y.S. (2005). Semi-parametric analysis for additive risk model via empirical likelihood. Communications in Statistics---Simulation and Computation, 34, 135-143.

Appendix A

FORTRAN Code for comparison competing risk in simulation

```
c
    This program is to implement tests of ordering
c
    of two conditional cause-specific hazard functions
c
    and two cumulative incidence functions.
c
   Simulated data example.
c
c
    Test H_0: g_1(t|z)=g_2(t|z) on [t_1,t_2]
c
        where z, t1 and t2 need to be specified
c
    versus
c
c
       H_1: F_1(t|z) \le F_2(t|z) on [t1,t2]
c
c
       H_2: g_1(t|z) \le g_2(t|z) \text{ on } [t_1,t_2]
     (with strict inequality for some t)
c Note: H_1 and H_2 are denoted H_2 and H_3 in the paper.
   program pvalues
   parameter(mxnsmp=1000,mxncov=2)
c
    ncov: number of covariates
c
c
    set mxncov=ncov in the subroutine ESTP as well!!
c
c
    nsamp: sample size
c
    time(i): failure time for subject i
c
    censor(i): cause-of-failure = 0 (censored), 1 or 2
c
    covar(j,i,k): covariate j for subject i at k-th event time
c
c
    clevel(j): specified level of covariate j
    nboot: number of simulated test processes used
c
         to find P-value (nboot=1000 gives good results)
c
    nsim: number of runs (only needed for a simulation study
c
        to check accuracy of the tests;
c
c
   integer ncov,nsamp,indx(mxnsmp),seeda,seedb,seedc,seedd,
         nsim,nboot
   real censor(mxnsmp),covar(mxncov,mxnsmp,mxnsmp),
   $ time(mxnsmp),dat(14),ccensor(mxnsmp),cov(mxncov,mxnsmp),
   $ ctime(mxnsmp), s0(mxnsmp),
   $ U(mxncov),copy(mxnsmp),nrej,nreja,
  $ U2(mxncov),s1(mxncov,mxnsmp),s2(mxncov,mxncov),
   $ var(mxncov,mxncov),beta(mxncov),base2(2,0:mxnsmp),
   $ beta2(2,mxncov),s02(2,mxnsmp),clevel(mxncov),phi(0:mxnsmp),
   $ stt(0:mxnsmp),betav1,betav2,cbeta,clev,cc,
```

```
$ s12(2,mxncov,mxnsmp),w(mxnsmp),censor1(mxnsmp),
  $ censor2(mxnsmp),aa, bb(0:mxnsmp), phiboot(mxnsmp),
  $ v1(mxncov,mxncov),v2(mxncov,mxncov),
  $ uu(mxncov,0:mxnsmp),zt(mxnsmp,mxncov),
  $ g(mxncov,0:mxnsmp),gt(1,mxncov,mxnsmp),Yi,sumYi,
  $ cvalue(mxncov,mxncov),ctemp(mxncov,mxnsmp),
  $ cinv(mxncov,mxncov),ct(mxncov,mxncov),
  $ gcvalue(1,mxncov,mxnsmp),Zbar(mxncov,mxnsmp),
  $ temp1(mxncov,mxnsmp),temp2(mxncov,mxnsmp),
  $ value11, value22, value2(2,0:mxnsmp),
  $ value3(2,0:mxnsmp), cumulative(2,0:mxnsmp),delta(mxnsmp),
  $ uu1(mxncov),uu2(mxncov),b1(0:mxnsmp),b2(0:mxnsmp)
   *** Give some values to simulate data ***
   Use these input values for the numerical example:
     42 37 45 51 1. 1. .3 400 .5 1000 0. 0.15
c
   seeda=42.
   seedb=37.
   seedc=45.
   seedd=51.
   betav1=1.
   betav2=1.
   cbeta=.3
   nsamp=200
   clev=.5
   nboot=1000
   t1=0.
   t2=0.15
   nsim=1000
   call rstart(seeda,seedb,seedc,seedd)
   ncov=2
   do 5 j=1,ncov
    clevel(j)=clev
  continue
   rcounta=0.
   rcount=0.
   do 1000 isim=1,nsim
   do 10 i=1,nsamp
    zz=uni()
     zzz=uni()
    cov(1,i)=zz
     cov(2,i)=zzz
```

```
tim1=-alog(uni())/(betav1*zz+betav1*zzz+1)
    tim2=-alog(uni())/(betav2*zz+betav2*zzz+2.5)
    cc=-alog(uni())*exp(-cbeta*zz-cbeta*zzz)
    if ((tim1.lt.tim2).and.(tim1.lt.cc)) then
      censor(i)=1.
      time(i)=tim1
    endif
    if ((tim2.lt.tim1).and.(tim2.lt.cc)) then
      censor(i)=2.
      time(i)=tim2
    endif
    if ((tim1.ge.cc).and.(tim2.ge.cc)) then
      censor(i)=0.
      time(i)=cc
    endif
10 continue
   print *, ' '
print*,'data simulated'
    *** order the data by time ***
call indexx(nsamp,time,indx)
  do 20 i=1,nsamp
    copy(i)=time(i)
20 continue
  do 40 i=1,nsamp
    time(i)=copy(indx(i))
40 continue
   do 60 i=1,nsamp
    copy(i)=censor(i)
60 continue
   do 80 i=1,nsamp
    censor(i)=copy(indx(i))
80 continue
   do 95 j=1,ncov
    do 90 i=1,nsamp
      copy(i) = cov(j,i)
     continue
    do 95 i=1,nsamp
```

```
cov(j,i) = copy(indx(i))
      do 94 ii=1,nsamp
      covar(j,i,ii) = cov(j,i)
94
      continue
95
      continue
c Find indices iii and jjj corr to time > t1 and
    first failure time <= t2)
   iii=1
   jjj=1
   do 100 i=1,nsamp
     if (time(i).lt.t1) then
      iii=i+1
     endif
     if (time(i).le.t2) then
      jjj=i
     endif
100 continue
   do 105 i=1,nsamp
     copy(i)=censor(i)
     if (censor(i).gt.1.5) then
       censor(i)=0.
     endif
     censor1(i)=censor(i)
      print*, i, censor(i)
105 continue
c -----
    do 200 j=1,ncov
     beta(j)=0.
200 continue
c *** CALL SUBROUTINE estp to get beta hat value ***
    call estp(nsamp,ncov,time,covar,censor,s0,s1,s2,beta,
  $ CL,var,NDEAD,NMIS)
   print *, 'beta1=', (beta(j),j=1,ncov)
   do 205 j=1,ncov
```

```
beta2(1,j)=beta(j)
205 continue
   print*, ''
c Find the first cumulative function
c for first cause-of-failure
   do 210 i=1,nsamp
        do 208 j=1,ncov
          base2(j,i)=0.
          cumulative(j,i)=0.0
       value2(j,i)=0.0
       value3(j,i)=0.0
208
      continue
210 continue
   start=0.0
   sumYi=nsamp
   value11=0.0
   bb1=0.0
   tempvalue1=0
   do 215 j=1,ncov
     bb1=bb1+beta2(1,j)*clevel(j)
215 continue
   do 225 i=1,nsamp
     s02(1,i)=s0(i)
     value11= value11 +censor1(i)/sumYi
     diff=time(i)-start
     do 220 j=1,ncov
       s12(1,j,i)=s1(j,i)
       Zbar(j,i) = S1(j,i)/S0(i)
       tempValue1=tempValue1 +beta2(1,j)*Zbar(j,i)*diff
220
      continue
     value2(1,i)=tempValue1
     value3(1,i)=value3(1,i) + time(i)*bb1
     base2(1,i)=base2(1,i-1)+value11-value2(1,i)
        cumulative(1,i)=cumulative(1,i)+value11 -value2(1,i)+value3(1,i)
```

```
start=time(i)
     sumYi=nsamp -i
     do 225 k=1,ncov
      s12(1,k,i)=s1(k,i)
225 continue
c Now consider second cause of failure:
   do 300 j=1,ncov
     beta(j)=0.
300 continue
   do 305 i=1,nsamp
     if (copy(i).ge.2.) then
      censor(i)=1.
     else
      censor(i)=0.
     endif
     censor2(i)=censor(i)
305 continue
c *** CALL SUBROUTINE estp to get second cause of failure ***
   call estp(nsamp,ncov,time,covar,censor,s0,s1,s2,beta,
  $ CL,var,NDEAD,NMIS)
   print *, 'beta2=', (beta(j),j=1,ncov)
   do 310 j=1,ncov
     beta2(2,j)=beta(j)
```

```
310 continue
 Find the second cumulative function ***
c -----
   value22=0
   start=0.0
   sumYi=nsamp
   bb2=0.0
   tempvalue2=0
   do 315 j=1,ncov
    bb2=bb2+beta2(2,j)*clevel(j)
315 continue
   do 330 i=1,nsamp
    s02(2,i)=s0(i)
    value22= value22 +censor2(i)/sumYi
    diff=time(i)-start
    do 320 j=1,ncov
      s12(2,j,i)=s1(j,i)
      Zbar(j,i) = S1(j,i)/S0(i)
      tempValue2=tempValue2 + beta2(2,j)*Zbar(j,i)*diff
320
      continue
    value2(2,i)=tempValue2
    value3(2,i)=value3(2,i) + time(i)*bb2
    base2(2,i)=base2(2,i-1)+value22-(value2(2,i))
       cumulative(2,i)=cumulative(2,i)+value22 -value2(2,i)+value3(2,i)
    start=time(i)
       sumYi=nsamp-i
    do 330 k=1,ncov
      s12(2,j,i)=s1(j,i)
330 continue
```

```
c
    st=1.
    st1=1.
    st2=1.
    st0=1.
    phi(0)=0.0
    do 350 i=nsamp
     phi(i)=0
350 continue
    bb1=0.
    bb2=0.
    do 400 j=1,ncov
     bb1=bb1+beta2(1,j)*clevel(j)
     bb2=bb2+beta2(2,j)*clevel(j)
400 continue
    sumYi=nsamp
    do 450 i=1,jjj
     st0=st
     st = st*(exp(-(cumulative(1,i)-cumulative(1,i-1) +
                        cumulative(2,i)-cumulative(2,i-1))))
     st1 = st1*(exp(-(cumulative(1,i)-cumulative(1,i-1))))
     st2 = st2*(exp(-(cumulative(2,i)-cumulative(2,i-1))))
     if (i.ge.iii) then
c for constant weight function W_1 use:
     w(i)=sqrt(float(nsamp))
c for weight function W_2 use
      w(i)=float(nsamp-i+1)/sqrt(float(nsamp))
   for weight function W_3 use
c
      w(i)=1./sqrt(2./sumYi)
c for weight function W_4 use
      w(i)=sqrt(float(nsamp))*st0
c
      phi(i)=sqrt(float(nsamp))*(st1-st2)
c
     phi(i)=phi(i-1)+((cumulative(2,i)-cumulative(2,i-1))-
             (cumulative(1,i)-cumulative(1,i-1)))*w(i)
     else
        phi(i)=phi(i-1)
```

```
endif
    sumYi=nsamp-i
     print*, time(i), cumulative(1,i),cumulative(2,i),st
450 continue
c **********************
  **** compute the test statistics ****
stata=phi(iii)
   stat=phi(iii)
   do 460 i=iii,jjj-1
    xx=phi(i)
    if (xx.gt.stata) then
     stata=xx
    endif
    do 460 j=i+1,jjj
    xxx=phi(j)-phi(i)
    if (xxx.gt.stat) then
     stat=xxx
    endif
460 continue
   N.B. the sqrt(nsamp) in front of the test statistic
      is merged into the weight function;
c
c
      this avoids having to normalize s^{(0)}
      by nsamp in various places.
c
   print*, ''
   print*, 'The output of sample size ',nsamp
   print *,'Test statistic (H_0 vs H_1) =', stata
   print *, 'Test statistic (H_0 vs H_2) =', stat
   print*,''
c **** simulation to determine P-value ****
c -----
   b1(iii-1)=0
   b2(iii-1)=0
   nreja=0.
   nrej=0.
   do 900 iboot=1,nboot
   sumYi=nsamp
   do 510 i=nsamp
    do 505 j=1,ncov
      uu1(i)=0
      uu2(j)=0
```

```
ctemp(j,i)=0
       g(j,i)=0
505
      continue
510 continue
c Find the second term of L(t)
c *** culculate G' value ***
   start=0.
   g(1,0)=0
   do 520 i=1, jjj
        diffvalue=time(i)- start
        do 515 j=1,ncov
          g(j,i)=g(j,i-1)+w(i)*(clevel(j)-zbar(j,i))*diffvalue
515 continue
     start=time(i)
520 continue
c -----
   start2=0
   do 580 i=1,nsamp
    diffvalue2=time(i)- start2
       do 525 j=1,ncov
          gt(1,j,i)=g(j,i)
525 continue
     print*, i, (gt(1,j,i), j=1,2)
c *** transpose of Z value ***
   do 530 j=1,ncov
          zt(i,j)=covar(j,i,i)
530 continue
   do 535 j=1,ncov
     ctemp(j,i)=ctemp(j,i)+(covar(j,i,i)-zbar(j,i))*diffvalue2
535 continue
   start2=time(i)
   do 550 K=1,ncov
     do 545 L=1,ncov
       temp10=zero
       do 540 M=1,1
```

```
temp10 = temp10 + ctemp(L,i)*zt(i,K)/float(nsamp)
540
       continue
     cvalue(L,K)=cvalue(L,K)+temp10
545
     continue
550 continue
580 continue
c -----
c *** get C inverse ***
c -----
  call MATINV(cvalue,ncov,ncov,IFLAG)
  do 620 \text{ K} = 1, ncov
    do 610 J = 1, ncov
     cinv(K,J)=cvalue(K,J)
610
    continue
620 continue
c -----
c *** get C transpose ***
c -----
  do 640 k=1,ncov
      do 630 L=1,ncov
     ct(K,L)=cinv(L,K)
630 continue
640 continue
   print*, ct(1,1), ct(1,2)
   print*, ct(2,1), ct(2,2)
  do 680 i=1,nsamp
      do 660 k=1,ncov
     temp15=zero
        do 650 j=1,ncov
       temp15 = temp15 + gt(1,j,i)*ct(j,k)/float(nsamp)
650
       continue
     gcvalue(1,k,i)=temp15
660
     continue
680 continue
c -----
  Find phiboot(i) then p-value
c -----
  do 750 i=iii,jjj
    b1(i)=b1(i-1)+w(i)*rnor()*censor1(i)/(sumYi)
    b2(i)=b2(i-1)+w(i)*rnor()*censor2(i)/(sumYi)
    a1 = 0
    a2 = 0
```

```
yy1=rnor()*censor1(i)
     yy2=rnor()*censor2(i)
     do 710 j=1,ncov
       uu1(j)=uu1(j)+yy1*(covar(j,i,i)-zbar(j,i))
       uu2(j)=uu2(j)+yy2*(covar(j,i,i)-zbar(j,i))
       a1=a1 + gcvalue(1,j,i)*uu1(j)
       a2=a2 + gcvalue(1,j,i)*uu2(j)
710
       continue
        sumYi=nsamp-i
     phiboot(i)=b1(i)+b2(i) + (a1+a2)/float(nsamp)
750 continue
c -----
   bstata=phiboot(iii)
   bstat=phiboot(iii)
   do 800 i=iii,jjj-1
   xx=phiboot(i)
   if (xx.gt.stata) then
     bstata=xx
   endif
   do 800 j=i+1,jjj
    xxx=phiboot(j)-phiboot(i)
     if (xxx.gt.bstat) then
      bstat=xxx
     endif
800 continue
   if (bstata.gt.stata) then
     nreja=nreja+1.
   endif
   if (bstat.gt.stat) then
    nrej=nrej+1.
   endif
900 continue
   print*, ''
   print*, 'P-value (vs H_1) = ', nreja/float(nboot)
   print*, 'P-value (vs H_2) = ', nrej/float(nboot)
   print *,'(H_1 and H_2 are denoted H_2 and H_3 in the paper!)'
   print*,'Completed simulation',isim, ', runs with nboot = ', nboot
   if (nreja/float(nboot).lt.0.05) then
     rcounta=rcounta+1.
```