Georgia State University ScholarWorks @ Georgia State University

Biology Theses

Department of Biology

8-27-2011

Modulation of NALP3 Inflammasome Genes by Estrogen

Anton P. Chesnokov Mr. Georgia State University

Follow this and additional works at: http://scholarworks.gsu.edu/biology_theses

Recommended Citation

Chesnokov, Anton P. Mr., "Modulation of NALP3 Inflammasome Genes by Estrogen." Thesis, Georgia State University, 2011. http://scholarworks.gsu.edu/biology_theses/33

This Thesis is brought to you for free and open access by the Department of Biology at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Biology Theses by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

MODULATION OF NALP3 INFLAMMASOME GENES BY ESTROGEN

by

ANTON CHESNOKOV

Under the Direction of Dr. Roberta Attanasio

ABSTRACT

Immunity is known to be sexually dimorphic. This dimorphism may be attributed to the action of different hormones. Aluminum is a component of several vaccines and acts as an adjuvant of immunogenicity. The activation of the Nalp3 inflammasome plays a role in aluminum's adjuvancy. Estrogen affects immune cells by regulating the expression of genes involved in immune-related mechanisms; such as the modulation of cytokine secretion. We hypothesized that estrogen modulates the aluminuminduced secretion of IL-1 β and IL-18. Using an *ex vivo* mouse macrophage model this study examined: (i) the effects of estrogen on Nalp3 inflammasome genes expression and (ii) the estrogen receptor involved in the modulation of these genes. Our results indicate that estrogen up-regulates Nalp3 gene expression via ER α/β heterodimerization, and caspase-1 activity may be indirectly modulated due to the upregulation of SPI-6 via ER β .

INDEX WORDS: Nalp3 inflammasome, Estrogen, IL-1beta, Aluminum hydroxide

MODULATION OF NALP3 INFLAMMASOME GENES BY ESTROGEN

by

ANTON CHESNOKOV

A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of

Master of Science

in the College of Arts and Sciences

Georgia State University

2011

Copyright by Anton Pavelovich Chesnokov 2011

MODULATION OF NALP3 INFLAMMASOME GENES BY ESTROGEN

by

ANTON CHESNOKOV

Committee Chair: Dr. Roberta Attanasio

Committee: Dr. Franco Scinicariello

Dr. Andrew Clancy

Electronic Version Approved:

Office of Graduate Studies

College of Arts and Sciences

Georgia State University

August 2011

ACKNOWLEDGEMENTS

I thank my committee for allowing me to present this document to them. I thank my advisor, Dr. Roberta Attanasio, for giving me the opportunity to develop a scientific mind, conduct research, and allow me to create a project that I became passionate about. And thanks to the many colleagues that have helped me in developing, and establishing, a critical and scientific thought process.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS iv					
LIST OF FIGURES vii					
LI	ST OF /	ABBREVIATIONSvii			
1	INTE	RODUCTION			
	1.1	Purpose of the Study7			
2	EXP	ERIMENTS			
	2.1	Animals			
	2.2	Ovariectomy			
	2.3	Isolation of Bone Marrow Derived Macrophages (BMDM)8			
	2.4	Cell Culture Stimulation9			
	2.5	Supernatant and RNA collection9			
	2.6	ELISA			
	2.7	Real-Time PCR			
	2.8	Flow Cytometry			
	2.9	Statistical Analysis			
3	RES	ULTS			
	3.1	BMDM are F4/80 and CD11(b) Positive11			
	3.2	17β-estradiol Increases Nalp3 Expression12			
	3.3	Stimulation of BMDM Decreases the Gene Expression of ASC14			

	3.4	Estrogen Receptor Agonists Modulate Caspase-115
	3.5	Serpinb9 is Up-regulated in Stimulated Macrophages via ERβ16
	3.6	BMDM Constitutively Express IL-1 eta mRNA, but at Much Lower Levels then when
Stimula	ated	
	3.7	17 eta -Estradiol does not Modulate IL-1 eta Secretion in BMDM Stimulated with Aluminum
and LP	S	
4	DI	SCUSSION19
5	СС	DNCLUSIONS
6	RE	FERENCES

LIST OF FIGURES

Figure 1.1 Murine BMDM were F4/80 Positive11
Figure 1.2 Murine BMDM were CD11(b) Positive11
Figure 2.1 17β-estradiol Increased Endogenous Nalp3 Gene Expression
Figure 2.2 17β-estradiol Increased Nalp3 Gene Expression in Stimulated Macrophages
Figure 3.1 Stimulation of BMDM with Aluminum and LPS Down-regulated ASC Gene Expression
Figure 4.1 Estrogen Receptor Beta Increased Caspase-1 Gene Expression15
Figure 5.1 Estrogen Receptor Beta Increased Serpinb9 Expression in Stimulated Macrophages
Figure 6.1 BMDM Increased Production of IL-1β mRNA after Stimulated17
Figure 7.1 Estrogen Receptor Agonists Did Not Modulate IL-1 β Secretion in BMDM Stimulated with
Aluminum and LPS

LIST OF ABBREVIATIONS

AlAluminum	Hydroxide
------------	-----------

- LPS.....Lipopolysaccharide
- Мф.....Масrophage
- BMDM......Bone Marrow-Derived Macrophage
- E2.....17β-estradiol
- PPT.....Propylpyrazoletriphenol
- DPN.....Diarylpropionitrile
- Nalp3.....Nucleotide-binding site and leucine-rich repeats with N-terminal pyrin domain containing 3
- ASC.....Apoptosis-associated speck-like protein containing a caspase recruitment domain

IL-....Interleukin

SPI-6.....Serine protease inhibitor-6

1 INTRODUCTION

Since Edward Jenner developed a smallpox vaccine at the turn of the 19th century, vaccinations have been an efficient and common means of protecting the population from infectious diseases. Today, a large number of vaccines are administered for many different kinds of diseases. An adjuvant of a vaccine allows for a more pronounced stimulation of the immune system, i.e. resulting in higher antibody titers. Aluminum has been administered in vaccines as an adjuvant since the 1930's. Rhoads (1930) found an aluminum precipitated antigen to provide immunity against polio in Rhesus macaques. In 1932, Park and Schroder reported a higher percentage of children obtained immunity when vaccinated with an aluminum precipitated diphtheria toxin. Eventually a large study by Volk and Bunney (1939) included 1,800 children and found that the best response was induced in patients who took the aluminum precipitated toxoid. The aluminum adjuvant is known to increase the effectiveness of vaccines by increasing antibody titers (Volk and Bunney 1941; Gupta 1998). The initial findings suggested that absorption of the toxoid/antigen onto the aluminum resulted in the precipitation of the toxoid/antigen from it, resulting in a steady release of antigen (Carlinfanti 1949; Gupta 1998). In theory, this steady release would maintain stimulation of the immune response longer and is a result of the adjuvant's action.

It has been recently established that a protein complex known as the 'Nalp3 inflammasome,' found within, but not limited to, monocytes, macrophages, and dendritic cells (Eisenbarth, Colegio et al. 2008), allows for aluminum's adjuvancy. The complex is composed of several proteins: Nalp3, ASC, pro-Caspase-1, and possibly CARDINAL. The protein nucleotidebinding site (a.k.a. nucleotide-binding oligomerization domain) and leucine-rich repeats with Nterminal pyrin domain containing 3 (Nalp3), seems to be the first cellular proteins to be involved. This protein is constitutively expressed, but without appropriate signals remains in an inactive conformation with interactions with HSP90 and SGT1. (Mayor, Martinon et al. 2007). Upon stimulation with danger associated molecular patterns, such as ATP or pathogen associated molecular pattern, Nalp3 is presumed to detach from HSP90 and SGT1 and exposes the pyrin domain (Duncan, Bergstralh et al. 2007; Mayor, Martinon et al. 2007). The activation of Nalp3 is followed by the binding of apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC) to Nalp3's pyrin domain (Mariathasan, Newton et al. 2004; Franchi and Gabriel 2008). This model suggests that ASC (as well as CARDINAL, in humans), once bound to Nalp3, allows for the binding of pro-caspase-1, resulting in caspase-1 activation (Agostini, Martinon et al. 2004; Mariathasan, Newton et al. 2004; Petrilli, Dostert et al. 2007; Church, Cook et al. 2008). The activated caspase-1 can now cleave pro-cytokines IL-1 β and IL-18, turning them into active cytokines (Thornberry 1992; Gu, Kuida et al. 1997; Eisenbarth, Colegio et al. 2008; Franchi and Gabriel 2008). Aluminum salts activate the Nalp3 protein, either directly or indirectly and aluminum's adjuvancy manifests in the secretion of the IL-1 β and IL-18 cytokines (Li, Nookala et al. 2007; Eisenbarth, Colegio et al. 2008; Gregorio, Tritto et al. 2008; Li, Willingham et al. 2008).

The cytokines IL-1 β and IL-18 are included in the IL-1 gene family, but produce different effects on the cells they target. IL-1 β is an endogenous pyrogen, or inflammatory inducer, and acts on many different types of tissues; it induces fever, dilation of vascular endothelium, and increasing expression of cell adhesion molecules (Parham 2009). IL-1 β can also activate B-cells

and T-cells. IL-18 contributes to inflammation and can be produced by different cell types, including macrophages (Stoll, Jonuleit et al. 1998; Murakami, Otsuki et al. 2005; Dinarello 2006), and, together with IL-2 or IL-12, co-stimulates the production of interferon- γ and other cytokines from T-cells and macrophages, among others (Dinarello 1996, 2006, 2009). This costimulation aids in the development of type 1 T helper cells, via interferon- γ (Dao, Ohashi et al. 1996; Stoll, Jonuleit et al. 1998), which function to activate macrophages (Parham 2009).

Studies have shown that after having been stimulated with appropriate factors, mice deficient in Nalp3, ASC, or Caspase-1 are unable to secrete cytokines (Mariathasan, Newton et al. 2004; Franchi and Gabriel 2008; Kool, Petrilli et al. 2008; Li, Willingham et al. 2008). Aluminum does not stimulate the production of cytokines, but rather only activates the Nalp3 protein; the synthesis of cytokines involves a separate pathway, including Toll-like-receptor ligands such as lipopolysaccharides (LPS) (Li, Nookala et al. 2007; Li, Willingham et al. 2008). In addition to the Nalp3 protein's importance in the secretion of these cytokines, higher antibody titers have also been attributed to Nalp3 gene expression; titers against a specific antigen are decreased in Nalp3 knock-out mice (Eisenbarth, Colegio et al. 2008; Li, Willingham et al. 2008).

Varying effectiveness of vaccines between the sexes has been reported (Giron-Gonzalez, Moral et al. 2000; Card, Carey et al. 2006; Cook 2008). All the factors for this sexual dimorphism are not known, but it has been shown that estrogen plays a role in immunity and/or on immune cells (Hu, Mitcho et al. 1988; Deshpande, Khalili et al. 1997; Kramer and Wray 2002; Masseoud 2008; Nguyen, Masseoud et al. 2011). Estrogen has been found to affect IL-1 production of monocytes/macrophages, seemingly increasing IL-1 production (Yoshida, Nakamura et al. 1996), with negative feedback at higher concentrations (Hu, Mitcho et al. 1988; Polan, Daniele et al. 1988; Polan, Loukides et al. 1989). However, a more recent study found that human peripheral blood monocytes increase IL-1β secretion at high levels of estradiol (100 nM) (Pioli, Weaver et al. 2006). In correlation with this, 17β-estradiol (E2) up-regulates IL-1β mRNA expression (Harris, Feldberg et al. 2000; Pioli, Weaver et al. 2006). Estrogen-estrogen receptor complex is known to enhance the IL1-β promoter (Ruh, Bi et al. 1998). Also, an estrogen responsive B box protein has been identified to interact with proIL-1β, enhancing secretion (Munding, Keller et al. 2006). Further, the IL-1β transcription factor, NF- κ B, was shown to interact with estrogen receptor α (Ghisletti, Meda et al. 2005; Gionet, Jansson et al. 2009), but the effects are not clear.

However, a contradictory report suggests a down regulation of IL-1 secretion monocytes/macrophages in response to E2 (Deshpande, Khalili et al. 1997; Morishita, Miyagi, and Iwamoto 1999). Another study showed that sera from male mice express higher responses to LPS stimulation with respect to cytokine secretion (Marriott, Bost et al. 2006). This same study showed that male macrophages secrete more IL-1 β *in vitro*, than females. Further, a trend of lower IL-1 β levels was observed from cells of postmenopausal women undergoing estrogen replacement therapy than those not undergoing the therapy (Rogers and Eastell 1998). Similarly, postmenopausal women were found to have increased IL-18 levels than fertile women (Cioffi, Esposito et al. 2002). In addition to contrary data, human monocyte-derived macrophages were shown not to alter IL-1 β gene expression levels in the presence of E2 at low or high concentrations (Corcoran, Meydani et al. 2010). Also, a study by Lewis et al. (2008) shows no effect on IL-1 β secretion in microglia when treated with 17 β -estradiol, but a reduction when treated with DPN, an ER β agonist. Estrogen has also been shown to impact IL-18 mRNA production, which was inhibited in the mouse uteri (Murakami, Otsuki et al. 2005).

In addition to the direct modulation of IL-1 β and IL-18 gene expression, estrogen receptors can up-regulate the expression of a caspase-1 inhibitor, serine protease inhibitor 6 (SPI-6), which also inhibits granzyme B, a cytotoxic enzyme secreted by cytotoxic T-cells and natural killer cells. SPI-6 gene expression is induced in response to both LPS and IL-1 β in some immunoprivileged cells, suggesting a negative feedback mechanism for controlling inflammation and/or protection from granzyme B degradation (Kannan-Thulasiraman and Shapiro 2002; Bots, de Bruin et al. 2007; Andrew, Simkins et al. 2008). The SPI-6 gene, Serpinb9, contains an estrogen responsive unit downstream of the transcription site and is responsible for estrogen induced expression (Krieg, Krieg, and Shapiro 2001; Krieg, Krieg et al. 2004). Regardless of the influence that estrogen may have on the expression of SPI-6, Stout-Delgado et al. (2007) found no difference in expression between genders in hepatocytes from mice challenged with a viral infection. The expression of SPI-6 is well established in the dendritic cell model, an antigen presenting cell of the myeloid lineage (Andrew, Simkins et al. 2008); immunohistological data of multiple tissues showed negative staining for SPI-6 in macrophages (Bladergroen, Strik et al. 2001). However, preliminary data from our lab suggests SPI-6 expression in the peritoneal macrophage cell line PMJ2, but estrogenic modulation was not present (unpublished). The expression in macrophages and the potential estrogenic modulation of this gene may imply a mechanism for estrogenic control of the Nalp3 inflammasome activity, namely the ability of caspase-1 to cleave pro-cytokines IL-1 β and IL-18.

The clinical implications of estrogenic modulation of the inflammasome genes can be applied to macrophagic myofasciitis and several chronic inflammatory diseases, collectively known as cryopyrin-associated periodic syndrome (CAPS). CAPS is attributed to mutations within the Nalp3 gene (Hoffman, Mueller et al. 2001; Verma, Eriksson et al. 2010; Hoffman and Brydges 2011). Agostini et al. (2004) showed that monocytes from individuals with Muckle-Wells syndrome, a type of CAPS, constitutively secrete IL-1 β , and these levels were only slightly lower than stimulated macrophages from normal donors. Murine bone marrow derived macrophages containing a Nalp3 mutation that is common among sick individuals produce constitutive IL-1β and the addition of pathogen associated molecular patterns induced further secretion without the aid of a Nalp3 activator, such as ATP (Meng, Zhang et al. 2009). Therapy for CAPS often addresses the suppression of IL-1 β , via antagonists, namely, Anakinra. Macrophage myofasciitis is a newly described chronic inflammatory disease which is attributed to the inability of the body to remove the aluminum that is injected with a vaccine and thus causing chronic stimulation of macrophages at the site; the presence of aluminum within macrophages at the injection site of vaccines have repeatedly been reported (Lacson, D'Cruz et al. 2002; Lach and Cupler 2008). If estrogen has the ability to regulate the inflammatory response, via Nalp3 inflammasomal modulation, then an increase in systemic estrogen has the potential to influence the degree and duration of chronic inflammation in patients with macrophage myofasciitis or CAPS.

Preliminary data from our lab show that in the presence of 17β-estradiol Nalp3 gene expression is up-regulated when the peritoneal macrophage PMJ2 is stimulated with ATP; however, this modulation is not present when stimulated with aluminum hydroxide (unpublished). If estrogen modulates the expression of Nalp3 inflammasome genes the diseases associated with it can be understood on a gender basis. Also, controlling hormonal levels in these patients may be viewed as a possible therapeutic approach.

There are many factors involved in the processing of both steroids and cytokines, making it difficult to obtain a clear understanding of estrogen's impact on IL-1 β . Currently, there are no studies that have observed estrogen's effects on aluminum adjuvancy. This study seeks to understand estrogen effects on Nalp3 associated gene expression. Determining the impact that estrogen may have on aluminum induced cytokine secretion and on Nalp3 inflammasome gene expression may help understand how estrogens impact aluminum adjuvancy. It is hypothesized that 17 β -estradiol will: (i) decrease aluminum induced IL-1 β secretion from murine bone marrow-derived macrophage, and (ii) up-regulate SPI-6 gene expression via estrogen receptor β . In support of this hypothesis, cells treated with E2 and diarylpropionitrile (DPN), an ER β agonist, are expected to show lower levels of IL-1 β in the supernatant, and this will correlate with an increase in gene expression of SPI-6.

1.1 Purpose of the Study

No study has addressed the influence of estradiol on aluminum adjuvancy. This study investigates how estradiol modulates the genes of the Nalp3 inflammasome, which is a key factor in aluminum adjuvancy. The genes of direct interest to the inflammasome are: Nalp3, ASC, and Caspase-1. Other genes that influence the function or outcome of the inflammasome were also investigated: namely, IL-1 β and SPI-9. This study also attempts to determine the estrogen receptor responsible for the modulation of Nalp3 inflammasome genes by using specified estrogen receptor agonists.

7

2 EXPERIMENTS

2.1 Animals

Female BALB/c mice (Charles River Laboratories, Inc., Wilmington, MA) were housed together in an individually ventilated cage with a 12hr day and night cycle (lights from 7am – 7pm). Cages and water were replaced once a week. Chow was isoflavone and soy-free (Stewart's Feed, #5K96). All animal procedures were conducted in agreement with the institutions policies and by the guidelines of the IACUC approved protocol.

2.2 Ovariectomy

At the age of 10.5 weeks, mice were anesthetized with 5% isoflurane gas in 100% medical grade oxygen. Mice were administered subcutaneously buprenorphine (0.1 mg/kg). While on a heating pad the mice's ovaries were removed by dorsal entry. The incisions were stapled and allowed to fall off naturally.

2.3 Isolation of Bone Marrow Derived Macrophages (BMDM)

At 6 months of age, mice were anesthetized with 5% isoflurane gas in 100% medical grade oxygen and euthanized by axillary cut-down. Femur and tibia bones were removed and tissue was removed. The bone epiphyses were removed in a biosafety cabinet and bone marrow was flushed out with RPMI1640 (Gibco) until the bone was white. The marrow plugs were mechanically disrupted with a 19-G needle and filtered through a 70 µm cell strainer. Each mouse's bone marrow was cultured in two non-culture treated petri dishes with 20 ml of complete media (PRMI1640 supplemented with 10% FBS, 5 mM glutamine, 1 mM HEPES, and 1% Pen-Strep (Gibco), plus 10 ng/ml M-CSF (Leinco). On day 3, 5 ml of media was removed and an

additional 10 ml of complete media was added; the complete media was replaced on day 6. On day 7 cells were identified by flow cytometry with staining for F4/80 (Invitrogen) and CD11(b) (BD Pharmingen).

2.4 Cell Culture Stimulation

On day 8 of culture, cells were counted and plated in 12-well plates at 5×10^5 cells per well in 2 ml. After 24 hrs the media was replaced with 1 ml of hormone-free media (complete media without phenol red and containing charcoal-stripped FBS) supplemented with either ethanol dissolved 17 β -estradiol (E2) at 1 or 10 nM, propypyrazoletriphenol (PPT) at 5 nM, or diarylpropionitril (DPN) at 10 nM, or vehicle (ethanol at 0.001%) for 12 hrs. PPT served as an ER α agonist (Stauffer, Coletta et al. 2000), whereas DPN served as an ER β agonist (Meyers, Sun et al. 2001). Cells were then primed with lipopolysaccharide (LPS) at 100 ng/ml for 1 hr with their respective hormone. This was replaced with media containing their respective hormone, LPS, and aluminum hydroxide (AI) at 125 µg/ml for 6 hrs. Positive controls were primed/stimulated with LPS without hormones or aluminum for 7 hrs and ATP at 5 mM for the last 20 min. All reagents were obtained from Sigma-Aldrich.

2.5 Supernatant and RNA collection

Supernatants were removed and centrifuged at 10,000 x g for 10 min at 4°C, then frozen at -80°C. Aluminum containing groups were washed 3X with PBS before RNA extraction. RNA was obtained using the RNeasy kit (Qiagen) according to manufactures protocol, and stored at -20°C.

2.6 ELISA

ELISA was conducted using a mouse IL-1β ELISA Set (BD OptEIA). Coating buffer was a 1 M phosphoric acid solution. Assay diluent was a 5% FBS in PBS solution. The stop buffer was a 1 M phosphoric acid solution. Samples were diluted 1/50. None stimulated groups (no aluminum or LPS) were conducted in either duplicates or triplicates. Aluminum containing samples were assayed in triplicates. Multiple assays were performed.

2.7 Real-Time PCR

Total RNA (2 μ l) was reverse transcribed with AMV reverse-transcriptase (Roche) in the presence of RNase inhibitor (Roche) using random primers (Roche). Diluted (1/20) cDNA underwent real-time PCR using Taqman[®] Gene Expression Assay (Applied Biosystems) in a 7500 Fast-Realtime machine (Applied Biosystems). Relative quantification was derived by $\Delta\Delta$ Ct, with β -actin as the endogenous control.

2.8 Flow Cytometry

Cells (3x10⁵) were stained for either anti-mouse F4/80 PE conjugate (Invitrogen) or antimouse CD11(b) FITC conjugate (BD Pharmingen). F4/80-PE antibody was compared to an isotype control (Biolegend), whereas the CD11(b)-FITC antibody was compared to an unstained cell sample.

2.9 Statistical Analysis

One-way ANOVA analysis was conducted with post-analysis using Tukey's multiple comparison test to determine significance. A p-value of <0.01 was considered statistically significant.

3 RESULTS

3.1 BMDM were Positive for F4/80 and CD11(b)

After 7 days of culture, bone marrow derived macrophages (BMDM) were stained for F4/80 expression to determine the purity of the derived macrophages. Over 98% of the cells expressed F4/80, deeming them murine macrophages (Figure 1.1). These cells were also assayed for a common monocyte/macrophage receptor, CD11(b), which was expressed at >90% of the cells (Figure 1.2).



Figure 1.1 BMDM were F4/80 Positive. Bone Marrow Derived Macrophages (BMDM) $(3x10^5$ cells) were stained with F4/80 antibody or an isotype control (PE conjugates) and analyzed by flow cytometry. Percentage is of 20,000 recorded events.



Figure 1.2 BMDM were CD11(b) Positive. Bone Marrow Derived Macrophages (BMDM) $(3x10^5$ cells) were stained with CD11(b) antibody (FITC conjugate) or unstained and analyzed by flow cytometry. Percentage is of 20,000 recorded events.

3.2 17β-estradiol Increased Nalp3 Expression

To determine how Nalp3 gene expression responds to particular estrogen receptor agonists, BMDM were stimulated with aluminum hydroxide (AI) and LPS and cDNA was assayed by real-time PCR (Figure 2.1-2). Estradiol (E2) up-regulated Nalp3 gene expression in a dose dependant manner; in unstimulated cells the gene expression increased 2 and 2.5-fold with E2 at 1 and 10 nM, respectively. This similar pattern was observed in the stimulated groups, however it was not statistically significant (p>0.05). The Nalp3 gene expression in the no hormone group increased ~56-fold after stimulation; whereas in the presence of 1 nM E2 the increase was ~62-fold, and at 10 nM E2 it was some ~68-fold. This increase – after treatment with 10 nM E2 – in stimulated macrophages was 20% more than the no hormone group. After stimulation and in the presence of diarylpropionitrile (DPN), an estrogen receptor (ER) beta agonist, Nalp3 gene expression increased to a similar degree as macrophages stimulated in the presence of E2 at 1 nM; propylpyrazoletriphenol (PPT), the ER α agonist, did not express this similar effect. Unstimulated macrophages treated with either PPT or DPN did not up-regulate Nalp3 expression.

The increase in endogenous expression of Nalp3 expression in response to E2 and not either of the ER agonists suggests that this modulation is attributed to the heterodimerization of ER α and β , both of which bind E2. The up-regulation in the presence of E2 may deem these cells more readily responsive to Nalp3 activating molecules.



Figure 2.1 17 β -estradiol Increased Endogenous Nalp3 Gene Expression. BMDM were cultured in the presence or absence of estrogen receptor agonists for 17 hrs. "No Hormones" group contains no estrogen receptor agonists, and is the reference point of expression. Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agoinist. Error bars represent the standard deviation of triplicate samples assayed in triplicates. ** (p<0.001) derived using Tukey's multiple comparison test.



Figure 2.2 17 β -estradiol Increased Nalp3 Gene Expression in Stimulated Macrophages. BMDM were cultured in the presence or absence of estrogen receptor agonists for 12 hrs prior to stimulation with LPS (100 ng/ml) for 1 hr and another 6 hrs with aluminum hydroxide (125 µg/ml). Quantification is normalized to a non-stimulated and no hormone containing group (not shown). Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agoinist. Error bars represent the standard deviation of triplicate samples assayed in triplicates.

3.3 Stimulation of BMDM Decreased the Gene Expression of ASC

For Nalp3 to active caspase-1, via proteolytic cleavage, the protein ASC is required to link the two proteins. Figure 3.1 shows how estrogen receptor agonists affect ASC gene expression before and after stimulation of BMDM with Al and LPS. Estradiol did not affect the endogenous expression of ASC. The estrogen receptor α and β agonists, PPT and DPN, respectively, down-regulated the expression of this gene in unstimulated cells; however, the reduction was minor. After stimulation, the macrophages expressed about half as much ASC. This decrease was not altered in the presence of any of the hormones tested.



Figure 3.1 Stimulation of BMDM with Aluminum and LPS Down-regulated ASC Gene Expression. BMDM were cultured in the presence or absence of estrogen receptor agonists for 12 hrs prior to stimulation with LPS (100 ng/ml) for 1 hr and another 6 hrs with aluminum hydroxide. Quantification is normalized to the unstimulated and no hormone containing group. Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agonist. Error bars represent the standard deviation of triplicate samples assayed in triplicates.

3.4 Estrogen Receptor Agonists Modulated Caspase-1

The expression of caspase-1 is vital to the adjuvancy effects of aluminum on macrophages, so it was of interests to determine whether estrogen modulates the expression of this gene (Figure 4.1). DPN, an ER β agonist, up-regulated caspase-1 in stimulated macrophages; DPN resulted in over a 3-fold increase, whereas no hormones yielded a 2.5-fold increase. The increase was modest when cells were treated with estradiol, for both 1 nM and 10 nM concentrations. PPT, an ER α agonist, increased the gene expression 3-fold, slightly higher than estradiol. This would suggest that the gene may be modulated in stimulated macrophages by ER α , and to a greater extent, ER β . Endogenous levels of the gene were not altered in the presence of any of the tested hormones.



Figure 4.1 Estrogen Receptor Beta Increased Caspase-1 Gene Expression. BMDM were cultured in the presence or absence of estrogen receptor agonists for 12 hrs prior to stimulation with LPS (100 ng/ml) for 1 hr and another 6 hrs with aluminum hydroxide. Quantification is normalized to a non-stimulated and no hormone containing group. Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agoinist. Error bars represent the standard deviation of triplicate samples assayed in duplicates (Unstimulated) or triplicates (Stimulated). *(p<0.01), ** (p>0.001), derived using Tukey's multiple comparison test as compared with stimulated No Hormone group.

3.5 Serpinb9 was Up-regulated in Stimulated Macrophages via ERβ

To determine a possible mechanism by which estrogen may inhibit IL-1 β secretion, the expression of the gene *Serpinb9*, which codes for a natural inhibitor of caspase-1, was determined (Figure 5.1). In the BMDM cell model, activation with LPS and aluminum increased the expression of *Serpinb9*; this suggests that *Serpinb9* is an inflammatory response gene. After activation, macrophages in the presence of DPN, an ER β agonist, increased *Serpinb9* expression 5-fold, as compared to the unstimulated and no hormone containing group; whereas a stimulated cells without hormones only increased gene expression by roughly 2-fold. This deems *Serpinb9* an ER β responsive gene, but up-regulation only occurs after cell stimulation/activation; DPN, and PPT, down-regulated the endogenous gene expression levels of *Serpinb9* to half the level of both no hormone and estradiol containing groups.



Figure 5.1 Estrogen Receptor Beta Increased *Serbinb9* Expression in Stimulated Macrophages. BMDM were cultured in the presence or absence of estrogen receptor agonists for 12 hrs prior to stimulation with LPS (100 ng/ml) for 1 hr and another 6 hrs with aluminum hydroxide. Quantification is normalized to a non-stimulated and no hormone containing group. Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agoinist. Error bars represent the standard deviation of triplicate samples assayed in triplicates. ** (p>0.001), derived using Tukey's multiple comparison test as compared with stimulated No Hormone group.

16

3.6 BMDM Constitutively Expressed IL-1 β mRNA, but at Much Lower Levels then when Stimulated

IL-1 β secretion is central to the Nalp3 inflammasome's function, so determining how estradiol might influence its gene expression can help in determining possible modulation of aluminum adjuvancy by estradiol. BMDM constitutively express IL-1 β mRNA, and PPT and DPN down-regulated these levels (Figure 6.1). However, these levels are minute when compared to stimulated cells. After stimulated with LPS and Al, IL-1 β gene expression in BMDM increased about 650-fold. Unlike the endogenous levels, in this model none of the estrogen receptor agonists influenced the levels of IL-1 β gene expression in activated cells (Figure 6.1).



6.1 BMDM Increased Production of IL-1 β mRNA after Stimulated. BMDM were cultured in the presence or absence of estrogen receptor agonists for 12 hrs prior to stimulation with LPS (100 ng/ml) for 1 hr and another 6 hrs with aluminum hydroxide. Quantification is normalized to a non-stimulated and no hormone containing group. Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agoinist. Error bars represent the standard deviation of triplicate samples assayed in triplicates. *(p<0.05), derived using Tukey's multiple comparison test as compared with non-stimulated No Hormone group.

3.7 17β-estradiol did not Modulate IL-1β Secretion in BMDM Stimulated with Aluminum and LPS

To determine if estrogen receptor agonists modulate the amount of IL-1 β secreted from BMDM in response to AI and LPS, ELISA assays were conducted on the supernatants from the cell cultures stimulated with these reagents. No significant modulation was observed in any of the stimulation groups (Figure 2.1). No secretion was detected in groups that were not stimulated with AI and LPS (data not shown).



Figure 7.1 Estrogen Receptor Agonists Did Not Modulate IL-1 β Secretion in BMDM Stimulated with Aluminum and LPS. All groups were cultured in hexlicates, except E2 (10 nM) which was conducted in triplicates. Cells were stimulated with LPS (100ng/ml) for 1 hr, followed by aluminum hydroxide (125µg/ml) for 6 hr, except the Positive Control, which was stimulated with LPS for 7 hr and ATP (5 mM) for the last 20 min. Hormone containing groups are: E2, estradiol; PPT, ER α agonist; DPN, ER β agoinist. Error bars represent standard error of the mean from six different samples, except E2 (10 nM) which is of three culture samples, assayed three different times in triplicates each time.

4 DISCUSSION

Estrogen is known to modulate several parameters of the immune system (reviewed in Bouman, Heineman, and Faas 2005). Classical estrogen receptors, ER α and ER β , are well established transcription factors that function by dimerization in either a homo- or a hetero- fashion (Kumar and Chambon 1988; Ogawa, Inoue et al. 1998). Here we investigated how 17 β -estradiol (E2) might modulate the gene expression of the Nalp3 inflammasome genes - Nalp3, ASC, and Caspase-1 – with and without lipopolysaccharide (LPS) and aluminum hydroxide (AI) activation of macrophages. In addition to E2, two estrogen receptor (ER) agonists were used: propylpyrazoletriphenol (PPT), an ER α agonist, and diarylpropionitrile (DPN), an ER β agonist. Bone marrow derived macrophages (BMDM), which express the characteristic macrophage markers F4/80 and CD11(b) (Fig. 1.1 & 1.2, respectively), are a common *in vitro* model in the study of the Nalp3 inflammasome. We also investigated the influence E2 and the ER agonists might have on IL-1 β gene expression and secretion in these cells.

If E2 has a role in modulating Nalp3 expression, then the implications may be extensive. Several chronic inflammatory diseases, collectively called CAPS, result from a constitutively active Nalp3 protein; expression of Nalp3 leads to uncontrolled secretion of IL-1 β , an endogenous pyrogenic cytokine. Estrogen has been implicated in rheumatoid arthritis (reviewed in Islander, Jochems et al. 2011), and Nalp3 induced IL-1 β has been implicated in gout (Joosten, Netea et al. 2010). To date, no reported study has determined whether E2 regulates the Nalp3 gene. Our results show that E2 up-regulates the endogenous levels of Nalp3 mRNA, and that this upregulation is likely a result of heterodimerization of ER α and β ; PPT and DPN, ER α and ER β agonists, respectively, did not show any up-regulation in this gene. Further, the subsequent increase in Nalp3 expression after increasing the E2 concentration to 10 nM suggests that this up-regulation is dose-dependent. This up-regulation by E2 was not apparent in the stimulated macrophages. Instead, stimulated cultures that were treated with E2 at 10 nM show a slight increase in Nalp3 gene expression, but the significance was not evident. The clear up-regulation of the Nalp3 gene in macrophages that have not been stimulated with LPS and Al may enable the cells to respond more strongly or quickly to a Nalp3 activator and hence would express the phenotypic manifestation of caspase-1 activation more readily. This would be a kinetically based response and whose effects have peaked within the six hours of stimulation that the current model used, and whose residual effects are seen by the slight variation in the stimulated cells. To verify this hypothesis, a kinetic study with multiply time points would have to be conducted. Even though E2 did not significantly influence Nalp3 gene expression in stimulated cells, the statistically significant endogenous up-regulation of Nalp3 by E2 may be implicated in the cryopyrin-associated periodic syndrome, wherein increased estrogen levels may lead to an immune flare-up.

ASC is an adaptor protein, linking caspase-1 to Nalp3, resulting in caspase-1 activation. The ASC gene was shown to decrease 2-fold in response to the LPS and Al stimuli. Indeed, Taxman et al. (2006) observed a sharp decrease in ASC mRNA expression in the human monocytic cell line THP-1, after 6 hrs of infection with *Porphyromonas gingivalis*, a gram-negative bacterium. The down-regulation of ASC in response to 6 hrs of stimulation may suggest a negative feedback to the inflammatory response, preventing the cell from inducing inflammation for too long of a period. The hormonal effects on the expression of ASC were not seen in either the unstimulated or stimulated cultures, suggesting that this gene is not modulated by estrogen.

The presence of caspase-1 is vital for an effective Nalp3 response. Caspase-1 is constitutively expressed and is up-regulated in response to LPS and Al (Figure 4.1). The endogenous level of caspase-1 gene expression is not affected by estrogen receptor agonists in cells that are not stimulated. However, DPN was found to significantly increase the levels of caspase-1 mRNA in stimulated macrophages. This increase suggests that ERß is an activator of this gene; interestingly though, neither of the E2 concentrations led to this increase. An increase in caspase-1 gene expression was observed in response to E2 in pigs' endometrium (Ashworth, Ross et al. 2010); however, these animals were not challenged with any type of PAMP and thus do not correlate with our data. Our data suggest that ERß up-regulates caspase-1 gene expression only in the presence of other stimuli, likely LPS, which is known to recruit a multitude of transcription factors resulting in an interaction. This suggests that $ER\beta$ requires the presence of other transcription factors to function as an activator of transcription. Indeed, Quaedackers et al. (2007) showed ERα to directly interact NF-κB proteins p50 and p65, which are up-regulated in LPS stimulated cells (Zandi, Rothwarf et al. 1997). Also, E2 was shown to increase NF-kB activity in splenic macrophages (Suzuki, Yu et al. 2008). Thus, it is possible to imagine how NF-kB transcription factors may interact with estrogen receptors. Because 17β-estradiol binds both ERα and β , the lack of caspase-1 gene expression increase after stimulation of macrophages in the presence of E2 may be due to antagonism by ERa, whereby ERa acts as a decoy for E2, preventing enough ER β ligation for the activation of caspase-1.

Several studies have suggested that the protein SPI-6, gene name *Serpinb9*, can inhibit caspase-1 activity (Annand, Dahlen et al. 1999; Kannan-Thulasiraman and Shapiro 2002). Fur-thermore, *Serpinb9* has been shown to be up-regulated in response to estrogens (Krieg, Krieg,

and Shapiro 2001; Krieg, Krieg et al. 2004). As mentioned above, caspase-1 is crucial to the secretion of IL-1 β and its inhibition would result in lower secretion levels of IL-1 β ; thus, if estrogen modulates IL-1 β secretion it would be of interest to determine if *Serpinb9* up-regulation would correlate. Our data show that endogenous levels of Serpinb9 were not affected by either of the E2 concentrations, but that the ER α and β agoinsts, PPT and DPN, respectively, decreased the expression levels roughly 2-fold (Figure 5.1); however this was not statistically significant. LPS is known to increase the expression of Serpinb9 (Kannan-Thulasiraman and Shapiro 2002; Andrew, Simkins et al. 2008), which would explain the up-regulation of the gene among all the stimulated groups. More interestingly though, was the large and significant up-regulation of Serpinb9 in response to DPN after stimulating the macrophages. This is contrary to Dr. David J. Shaprio's team at the University of Illinois, which consistently attributes ER α to the upregulation of Serpinb9 (Krieg, Krieg et al. 2004; Jiang, Ellison et al. 2007). Our data does suggest an increase similar to that of E2, at 1 nM, in response to the ER α agonist. Taking this into account, it is worth noting the lack in increase in E2 at 10 nM, which suggest an E2 threshold in acting as an activator of transcription. However, these data points contain lots of variation and are not statistically significant.

Several studies have attributed ER β to decreased immune responses in microglia (Baker, Brautigam and Watters 2004; Lewis, Johnson et al. 2008), the macrophages of the central nervous system. The 5-fold increase in *Serpinb9* in response to DPN, as compared to the roughly 2fold increase in other stimulated hormonal groups, would suggest that ER β has a potential to inhibit IL-1 β secretion via SPI-6. However, our data did not support such a hypothesis; the IL-1 β secretion levels remained similar. One possibility for a lack of reduction in IL-1 β secretion can be due to *Serpinb9* not being translated and thus no phenotypic modulation is seen. Indeed, macrophages where negative for PI-9, the human homolog, in lymphoid tissue samples; how-ever, these samples were derived from normal human tissues (Bladergroen, Strik et al. 2001).

The secretion of IL-1 β from macrophages stimulated with LPS and Al for six hours did not differ in the presence of any of the ER agonists tested (Figure 7.1). These data correlate well with the lack of modulation of IL-1 β gene expression (Figure 6.2), which is in agreement with human monocyte-derived macrophages (Corcoran, Meydani et al. 2010). The inability of the DPN up-regulated SPI-6, whose increased protein levels are assumed due to the increased mRNA levels, to reduce the amount of IL-1 β secretion may be offset by a corresponding upregulation of caspase-1. Another explanation is that *Serpinb9* might not have been upregulated before the six hour mark. Without kinetic gene expression data we cannot determine if the observed up-regulation of *Serpinb9* by DPN only just occurred at the six hour mark, and that the manifestation of caspase-1 inhibition – reduced IL-1 β secretion – can only be observed after this time point. Lewis et al. (2008) showed that DPN decreased IL-1 β secretion from LPS stimulated microglia, after 24 hours of stimulation. Our study chose six hours because LPS stimulated IL-1 mRNA peaks at this time point (Dinarello, Muegge, and Durum 2000).

Other factors that may have led to the lack of observable differences in IL-1 β secretion is the presence of M-CSF, a growth factor essential to BMDM, or the presence of LPS concentrations that are too high. Cells respond upon reaching a threshold level of stimulation; further, the response is not expressed until intracellular biochemical responses reach a particular threshold. This phenomenon explains the varying cellular responses to different concentrations of stimuli. M-CSF is known to influence IL-1 β secretion and activation of macrophages in general (Evans, Kamdar et al. 1995; Popova, Kzhyshkowska et al. 2011). Its presence was perhaps not essential, but is common in the macrophage milieu. The M-CSF concentrations used in this study (10 ng/ml) could have masked the effects that E2 might have had on the secretion of IL-1 β by exacerbating the macrophage's response to the LPS and Al stimuli. The LPS concentration, 100 ng/ml, may also have been too high. A response curve was not conducted on these cells, so similarly to the M-CSF, LPS might have masked the potential effects of estrogen on IL-1 β secretion.

5 CONCLUSIONS

Estrogen receptor agonists estradiol, PPT, and DPN modulate gene expression of particular Nalp3 inflammasome genes. Nalp3 gene expression was increased by estradiol at 1 nM and to a higher degree with 10 nM. Nalp3 modulation by estradiol may be attributed to heterodimerization of estrogen receptor (ER) α and β ; PPT and DPN, ER α and ER β agonist, respectively, did not show any increase in gene expression. Caspase-1 was up-regulated by all three agonists; caspase-1 gene expression is modulated by both ER α and ER β , but ER β seems to have a higher degree of influence. IL-1 β gene expression was not modulated by estradiol, PPT, or DPN in this model. SPI-6 gene expression in macrophages was up-regulated in response to stimulation with Al and LPS, and this up-regulation was greatly exacerbated in the presence of DPN. However, the up-regulation of SPI-6 gene expression did not result in decreased IL-1 β secretion. In conclusion, estrogen receptor agonists did not modulated IL-1 β secretion in murine bone marrow-derived macrophages stimulated with aluminum and LPS, while in the presence of M-CSF.

6 REFERENCES

- Agostini, L., M. Martinon, et al. (2004) "NALP3 forms an IL-1β-Processing inflammasome with increased activity in Muckle-Wells autoinflammatory Disorder." <u>Immunity</u> **20**: 319-325
- Andrew, K.A., H.M.A., Simkins, et al. (2008) "Dendritic Cells treated with lipopolysaccharide upregulate Serine Protease Inhibitor 6 and remain sensitive to killing by cytotoxic T Lymphocytes in vivo." Journal of Immunology 181: 8356-8362
- Annand, R. R., J. R. Dahlen, et al. (1999) "Caspase-1 (interleukin-1β-converting enzyme) is inhibited by the human serpin analogue proteinase inhibitor 9." <u>Biochemical Journal</u> **342**: 655-665
- Ashworth, M. D., J. W. Ross, et al. (2010). "Endometrial caspase 1 and interleukin-18 expression during the estrous cycle and peri-implantation period of porcine pregnancy and response to early exogenous estrogen administration." <u>Reproductive Biology and Endocrinology</u> **8**: 33
- Baker, A. E., V. M. Brautigam and J. J. Watters (2004). "Estrogen modulates microglial inflammatory mediator production iva interactions with estrogen receptor β." <u>Endocrinology</u> 145: 5021-5032
- Bladergroen, B.A., M.C.M. Strik, et al. (2001). "The granzyme B inhibitor, protease inhibitor 9, is mainly expressed by dendritic cells and at immune-privileged sites." <u>Journal of Immunology</u> 166: 3218-3225
- Bots M., E. de Bruin, et al. (2007). "Proteinase inhibitor-9 expression is induced by maturation in dendritic cells via p38 MAP kinase." <u>Human Immunology</u> **68**: 959-964
- Bouman A., M. J. Heineman and M. M. Fass (2005). "Sex hormones and the immune response in humans." <u>Human Reproduction Update</u> **11**: 411-423
- Card, J. W., M. A. Carey, et al. (2006). "Gender differences in murine airway responsiveness and lipopolysaccaride-induced inflammation." Journal of Immunology **177**: 621-630
- Carlinfanti, E. (1950). "Quantitative studies on diphtheria prophylactic and some considerations on the assessment of antigenicity." Journal of Immunology **64**: 373-379
- Church L. D., G. P. Cook, et al. (2008). "Primer: inflammasomes and interleukin 1β in inflammatory disorders." <u>Nature Clinical Practice Rheumatology</u> **4**: 34–42
- Cioffi, M., K, Esposito, et al. (2002). " Cytokine pattern in postmenopause." <u>Maturitas</u> **41**: 187-192

- Cook, I. F. (2008). "Sexual dimorphism of humoral immunity with human vaccines." <u>Vaccine</u> **26**: 3551-3555.
- Corcoran, M. P., M. Meydani, et al. (2010). "Sex hormone modulation of proinflammatory cytokine and C-reactive protein expression in macrophages from older men and postmenopausal women." Journal of Endocrinology **206**: 217-224.
- Dao, T., K. Ohashi, et al. (1996). "Interferon-γ-inducing factor, a novel cytokine, enhances Fas ligand-mediated cytotoxicity of murine T helper 1 cells." <u>Cell Immunology</u> **173**: 230-235
- Deshpande, R., H. Khalili, et al. (1997). "Estradiol down-regulates LPS-induced cytokine production and NFkB activation in murine macrophages." <u>American Journal of Reproductive Im-</u> <u>munology</u> **38**: 46-54
- Dinarello, C. A. (1998). "Interleukin-1β, interleukin-18, and the interleukin-1β converting enzyme." <u>Annals of the New York Academy of Sciences</u> **865**: 1-11
- Dinarello, C. A. (2006). "Interleukin 1 and interleukin 18 as mediators of inflammation and the aging process." <u>American Journal of Clinical Nurtition</u> **83**: 447S-455S
- Dinarello, C. A. (2009). "Immunological and inflammatory functions of the interleukin-1 family." <u>Annual Review of Immunology</u> **27**: 519-550
- Duncan, J. A., D. T. Bergstralh, et al. (2007). "Cryoprin/NALP3 binds ATP/dATP, is an ATPase, and requires ATP binding to mediate inflammatory signaling." <u>Proceedings of the National</u> <u>Academy of Sciences of the Unisted States of America</u> **104**: 6764-6769
- Eisenbarth, S. C., O. R. Colegio, et al. (2008). "Crucial role for the Nalp3 inflammasome in the immunostimulatory properties of aluminium adjuvants." <u>Nature</u> **453**: 1122-1126
- Evans, R. E., S. J. Kamdar, et al. (1995). "The potential role of the macrophage colonystimulating factor, CSF-1, in inflammatory responses: characterization of macrophage cytokine gene expression." Journal of Leukocyte Biology **55**: 99-107
- Franchi, L. and N. Gabriel (2008). "The NIrp3 inflammasome is critical for aluminium hydroxidemediated IL-1β secretion but dispensable for adjuvant activity." <u>European Journal of Immu-</u> <u>nology</u> **38**: 2085–2089.
- Ghisletti, S., C. Meda, et al. (2005). "17β-estradiol inhibits inflammatory gene expression by controlling NF-κB intracellular localization." <u>Molecular and Cellular Biology</u> **25**: 2957-2968

- Gionet, N., D. Jansson, et al. (2009). "NF-κB and estrogen receptor α interactions: differential function in estrogen receptor-negative and –positive hormone-independent breast cancer cells." Journal of Cellular Biochemistry **107**: 448-459
- Girón-González, J., F. J. Moral, et al. (2000). "Consistent production of a higher T_H1:T_H2 cytokine ration by stimulated T cells in men compared with women." <u>European Journal of Endocrino-</u> <u>logy</u> **143**: 31-36
- Gregorio, E. D., E. Tritto, et al. (2008). "Alum adjuvanticity: Unraveling a century old mystery." <u>European Journal of Immunology</u> **38**: 2068-2071
- Gu, Y., K. Kuida, et al. (1997). "Activation of interferon-γ inducing factor mediated by interleukin-1β converting enzyme." <u>Science</u> **275**: 206-209
- Gupta, R. K. (1998). "Aluminum compounds as vaccine adjuvants." <u>Advanced Drug Delivery Re-</u> views **32**: 155-172
- Harris, M. T., R. S. Feldberg, et al. (2000). "Expression of proinflammatory genes during estrogen-induced inflammation of the rat prostate." <u>The Prostate</u> **44**: 19-25
- Hoffman, H. M., J. L. Mueller, et al. (2001). "Mutation of a new gene encoding a putative pyrinlike protein causes familial cold autoinflammatory syndrome and Mucle-Wells syndrome." <u>Nature</u> 29: 301-305
- Hoffman, H. M. and S. D. Brydges. (2011). "The genetic and molecular basis of inflammasomemediated disease." Journal of Biological Chemistry **286**: 10889-10896
- Hu, S. K., Y. L. Mitcho, et al. (1988). "Effect of estradiol on interleukin 1 synthesis by macrophages." <u>International Journal of Immunopharmacology</u> **10**: 247-252
- Islander, U., C. Jochems, et al. (2011). "Estrogens in rheumatoid arthritis; the immune system and bone." <u>Molecular and Cellular Endoccrinology</u> **335**: 14-29
- Jiang, X., S. J. Ellison, et al. (2007). "Interplay between the levels of estrogen and estrogen receptor controls the level of the granzyme inhibitor, proteinase inhibitor 9 and susceptibility to immune surveillance by natural killer cells." <u>Oncogene</u> **26**: 4106-4114
- Kannan-Thulasiraman, P. and D. J. Shapiro (2002). "Modulators of inflammation use nuclear factor-κB and activator protein-1 sites to induce the caspase-1 and granzyme B inhibitor, proteinase inhibitor 9." Journal of Biological Chemistry **227**: 41230-41239
- Kool, M., V. Petrilli, et al. (2008). "Cutting Edge: Alum Adjuvant Stimulates Inflammatory Dendritic Cells through Activation of the NALP3 Inflammasome." <u>The Journal of Immunology</u> 181: 3755-3759.

- Kramer, P. R. and S. Wray (2002). "17-β-Estradiol regulates expression of genes that function in macrophage activation and cholesterol homeostasis." <u>Journal of Steroid Biochemistry &</u> <u>Molecular Biology</u> 81: 203-216
- Krieg, S. A., A. J. Krieg and D.J. Shapiro (2001). "A unique downstream estrogen responsive unit mediates estrogen induction of proteinase inhibitor-9, a cellular inhibitor of IL-1βconverting enzyme (caspase 1)." <u>Molecular Endocrinology</u> **15**: 1971-1982
- Krieg, A. J, S. A. Krieg, B. S. Ahn, and D. J. Shapiro (2004). "Interplay between estrogen response element sequence and ligands controls *in vivo* binding of estrogen receptor to regulated genes." <u>The Journal of Biological Chemistry</u> 279: 5025-5034
- Kumar, N. and P. Chambon (1988). "The estrogen receptor binds tightly to its responsive element as a ligand-induced homodimer." <u>Cell</u> **55**: 145-156
- Lach, B. and E. J. Cupler (2008). "Macrophagic myofasciitis in children is a localized reaction to vaccination." Journal of Child Neurology **23**: 614-619
- Lacson, A.G., C.A. D'Cruz, et al. (2002). "Aluminum phagocytosis in quadriceps muscle following vaccination in children: relationship to macrophagic myofasciitis." <u>Journal of Immunology</u> **177**: 621-630
- Lambert, C. K., E. M. Curran, et al. (2004). "Estrogen receptor-α deficiency promotes increased TNF-α secretion and bacterial killing by murine macrophages in response to microbial stimuli in vitro." Journal of Leukocyte Biology **75**:1166-1172
- Lewis, D.K., A.B. Johnson, et al. (2008). "Effects of estrogen receptor agonists on regulation of the inflammatory response in astrocytes from young adult and middle-aged female rats." Journal of Neuroimmunology **195**: 47-59
- Li, H., S. Nookala, et al. (2007). "Aluminum Hydroxide Adjuvants Activate Caspase-1 and Induce IL-1 β and IL-18 Release." <u>The Journal of Immunology</u> **178**: 5271-5276
- Li, H., S. B. Willingham, et al. (2008). "Cutting edge: inflammasome Activation by Alum and Alum's Adjuvant Effect Are Mediated by NLRP3." <u>The Journal of Immunology</u> **181**:17-21
- Mariathasan, S., K. Newton, et al. (2004). "Differential activation of the inflammasome by caspase-1 adaptor ASC and Ipaf." <u>Nature</u> **430**: 213-218
- Marriott, I., K. L. Bost, et al. (2006). "Sexual dimorphism in expression of receptors for bacterial Lipopolysaccharides in murine macrophages: A possible mechanism for gender-based differences in endotoxic shock susceptibility." Journal of Reproductive Immunology **71**: 12-27

- Masseoud, F. N. (2008). "Estrogen-Induced modulation of innate and adaptive immune function." Unpublished, Georgia State University 2008, Atlanta, Georgia, USA
- Mayor, A., F. Martinon, et al. (2007). "A crucial function of SGT1 and HSP90 in inflammasome activity links mammalian and plant innate immune responses." <u>Nature Immunology</u> 8: 497-503
- Meng, G., F. Zhang, et al. (2009). "A mutation in the NIrp3 gene causing inflammasome hyperactivation potentiates Th17 cell-dominant immune responses." <u>Immunity</u> **30**: 860-874
- Meyers, M. J., J. Sun, et al. (2001). "Estrogen receptor-β potency-selective ligands: structureactivity relationship studies of diarylpropionitriles and their acetylene and polar analogues." Journal of Medicinal Chemistry **44**: 4230-4251
- Morishita, M., M. Miyagi and Y. Iwamoto. (1999). "Effects of sex hormones on production of interleukin-1 by human peripheral monocytes." Journal of Periodontology **70**: 757-760
- Munding, C., M. Keller, et al. (2006). "The estrogen-responsive B box protein: a novel enhancer of interleukin-1β secretion." <u>Cell Death and Differentiation</u> **13**: 1938-1949
- Murakami, Y., M. Otsuki, et al. (2005). "Estrogen inhibits interleukin-18 mRNA expression in the mouse uterus." Journal of Reproductive Development **51**: 639-647
- Nguyen, D. C., F. Masseoud, et al. (2011). "17β-Estradiol restores antibody responses to an influenza vaccine in a postmenopausal mouse model." <u>Vaccine</u> **29**: 2515-2518
- Ogawa, S., S. Inoue, et al. (1998). "The complete primary structure of human estrogen receptor β (hER β) and its heterodimerization with ER α *in vivo* and *in vitro*." <u>Biochemical and Biophysical Research Communications</u> **243**: 122-126

Parham, P. (2009). <u>The Immune System. 3rd ed.</u> New York, Garland Science.

- Park, W. H. and M. C. Schroder. (1932). "Diphtheria toxin-antitoxin and toxoid: a comparison." <u>American Journal of Public Health</u> **22**: 7-16
- Pétrilli, V., C. Dostert, et al. (2007). "The inflammasome: a danger sensing complex triggering innate immunity." <u>Current Opinion in Immunology</u> **19**: 615-622
- Pioli, P. A., L. K. Weaver, et al. (2006). "Lipopolysaccharide-Induced IL1β production by human uterine macrophages up-regulates uterine epithelial cell expression of human β-Defensin2" <u>The Journal of Immunology</u> 176: 6647-6655
- Polan, M. L., J. Loukides, et al. (1989). "Gonadal steroids modulate human monocyte interleukin-1 (IL-1) activity." <u>Fertility and Sterility</u> **49**: 964-968

- Polan, M. L., J. Loukides, et al. (1989). "Progesterone and estradiol modulate interleukin-1β messanger ribonucleic acid levels in cultured human peripheral monocytes." <u>Journal of Clin-</u> <u>ical Endocrinology and Metabolism</u> **69**: 1200-1206
- Popova, A., J. Kzhyshkowska, et al. (2011). "Pro- and anti-inflammatory control of M-CSFmediated macrophage differentiation." Immunobiology **216**: 164-172
- Quaedackers, M. E., C. E. van den Brink, et al. (2007). "Direct interaction between estrogen receptor α and NF-κB in the nucleus of living cells." <u>Molecular and Cellular Endocrinology</u> **273**: 42-50
- Rhoads, C. P. (1930). "Immunization with aluminum hydroxide mixture of poliomyelitis virus." <u>Science</u> **72**: 608
- Rogers, A. and R. Eastell (1998) "Effects of estrogen therapy of postmenopausal women on cytokines measured in peripheral blood." Journal of Bone and Mineral Research **13**: 1577-1586
- Ruh, M. F., Y. Bi, et al. (1998). "Effect of estrogens on IL-1β promoter activity." Journal of Steroid Biochemical and Molecular Biology **66**: 203-210
- Stauffer, S. R., C. J. Coletta, et al. (2000). "Pyrazole ligands: structure-affinity/activity relationships and estrogen receptor-α-selective agonists." <u>Journal of Medicinal Chemistry</u> **43**: 4934-4947
- Stoll, S., H. Jonuleit, et al. (1998). "Production of functional IL-18 by different subtypes of murine and human dentritic cells (DC): DC-derived IL-18 enhances IL-12-dependent Th1 development." <u>European Journal of Immunology</u> 28: 3231-3239
- Stout-Delgado, H.W., Y. Getachew, B.C. Miller and D.L. Thiele (2007). "Intrahepatic lymphocyte expression of dipeptidyl peptidase I-processed granzyme B and perforin induces hepatocyte expression of serine proteinase inhibitor 6 (serpinb9/SPI-6)." Journal of Immunology 179: 6561-6567
- Suzuki, T., H. Yu, et al. (2008). "Estrogen-mediated activation of non-genomic pathway improves macrophages cytokine production following trauma-hemorrhage." <u>Journal of Cellu-</u> <u>lar Physiology</u> **214**: 662-672
- Taxman, D. J., J. Zhang, et al. (2006). "ASC mediates the induction of multiple cytokines by *Porphyromonas gingivalis* via caspase-1-dependent and –independent pathways." <u>The Journal of Immunology</u> **177**: 4252-4256

- Thornberry, N. A., H. G. Bull, et al.(1992). "A novel heterodimeric cysteine protease is required for interleukin-1β processing in monocytes." <u>Nature</u> **356**: 768-774
- Verma, D., P. Eriksson, et al. (2010). "Two adult siblings with atypical cryopryin-associated periodic syndrome due to a novel M299V mutation in *NLRP3*." <u>Arthritis and Rheumatism</u> 62: 2138-2143
- Volk, V. K. and W. E. Bunney (1939). "Diphtheria immunization with fluid toxoid and alum precipitated toxoid-preliminary report." <u>American Journal of Public Health</u> **29**: 197-204
- Volk, V. K. and W. E. Bunney (1940). "Diphtheria immunization with fluid toxoid and alum precipitated toxoid-progress report." <u>American Journal of Public Health</u> **30**: 42-43
- Yoshida, Y., Y. Nakamura, et al. (1996). "Changes in interleukin-1 production of peritoneal macrophages during estrous cycle in golden hamsters." <u>Endocrine Journal</u> **43**: 151-156
- Zandi, E., D. M. Rothwarf, et al. (1997). "The IκB Kinase Complex (IKK) contains two kinase subunits, IKKα and IKKβ, necessary for IκB phosphorylation and NF-κB activation." <u>Cell</u> **91**: 243-245