

Georgia State University

ScholarWorks @ Georgia State University

---

Psychology Faculty Publications

Department of Psychology

---

2012

## The role of mindfulness and psychological flexibility in somatization, depression, anxiety, and general psychological distress of a non-clinical college sample.

Akihiko Masuda

Georgia State University, amasuda@gsu.edu

Erin Tully

Georgia State University, etully2@gsu.edu

Follow this and additional works at: [https://scholarworks.gsu.edu/psych\\_facpub](https://scholarworks.gsu.edu/psych_facpub)



Part of the [Psychology Commons](#)

---

### Recommended Citation

Masuda, Akihiko and Tully, Erin, "The role of mindfulness and psychological flexibility in somatization, depression, anxiety, and general psychological distress of a non-clinical college sample." (2012).

*Psychology Faculty Publications*. 88.

[https://scholarworks.gsu.edu/psych\\_facpub/88](https://scholarworks.gsu.edu/psych_facpub/88)

This Article is brought to you for free and open access by the Department of Psychology at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Psychology Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

The Role of Mindfulness and Psychological Flexibility in Somatization, Depression, Anxiety,  
and General Psychological Distress of a Non-clinical College Sample

Akihiko Masuda, Ph.D.\* and Erin C. Tully, Ph.D.

Georgia State University

\*Corresponding author at: Department of Psychology, Georgia State University

Atlanta, GA 30303, USA.

Tel.: +1 (404) 413-6298

Fax: +1 (404) 413-6207.

Email Address: [amasuda@gsu.edu](mailto:amasuda@gsu.edu)

Short Title: Mindfulness and Psychological Flexibility

Word Count: 4,014

## Abstract

The current study investigated whether mindfulness and psychological flexibility uniquely and separately accounted for variability in psychological distress (somatization, depression, anxiety, and general psychological distress). An ethnically diverse, non-clinical sample of college undergraduates ( $N = 494$ , 76% female) completed a web-based survey that included the self-report measures of interest. Consistent with prior research, psychological flexibility and mindfulness were positively associated with each other, and tested separately, both variables were negatively associated with somatization, depression, anxiety, and general psychological distress. Results also revealed that psychological flexibility and mindfulness accounted for unique variance in all four measures of distress. These findings suggest that mindfulness and psychological flexibility are interrelated but not redundant constructs, and that both constructs are important for understanding the onset and maintenance of somatization, depression, anxiety, and general distress.

*Key words:* Psychological distress, somatization, depression, anxiety, mindfulness, psychological flexibility, experiential avoidance

## **Introduction**

Recently, cognitive behavioral therapies have been expanding to include mindfulness and psychological flexibility into their conceptual frameworks,<sup>1</sup> as growing evidence has shown their salutary role.<sup>2,3</sup> In theory, the two processes are often conceptualized as adaptive regulation and coping processes that reflect greater psychological health.<sup>3,4</sup> Literature also supports the effectiveness of acceptance- and mindfulness-based cognitive behavioral therapies that are designed to promote greater wellbeing through targeting these two processes.<sup>5-7</sup> Although psychological flexibility and mindfulness are often theorized to be similar but distinct processes, evidence supporting this conceptual position is still limited. As such, the present cross-sectional study aimed to understand the nature of the relationship among mindfulness, psychological flexibility, and psychological distress. In particular, this study quantified the redundant versus unique contributions of these two constructs to depression, anxiety, somatization, and general psychological distress.

### **Psychological Flexibility**

Psychological flexibility is roughly conceptualized as an overarching regulation process of (a) experiencing the present moment as it is without judgment and avoidance and (b) persisting or changing behavior when doing so serves valued-ends.<sup>6</sup> Psychological flexibility has been of great interest in recent years as accumulating evidence has supported its salutary effects. As such, a model has been developed to explain the relationship between psychological flexibility and psychological health. According to the psychological flexibility model,<sup>6</sup> greater psychological well-being is characterized by open and flexible contact with one's own internal and external environment and by commitment to value-consistent activities. Conversely, many forms of psychopathology are conceptualized in terms of diminished psychological flexibility,

which is marked by the excess of maladaptive affect/behavior regulations (e.g., thought suppression and avoidance) and by the deficits of contingency-sensitive and valued-directed behaviors. Accumulating evidence has shown that psychological flexibility is positively associated with psychological well-being<sup>3</sup> and inversely associated with a wide range of distress, including depression,<sup>8</sup> anxiety,<sup>9</sup> and general psychological distress.<sup>10-13</sup>

### **Mindfulness**

Mindfulness is another construct that has been widely incorporated into cognitive behavioral therapies in recent years.<sup>14</sup> Although the definition of mindfulness varies across investigations, it is often conceptualized as an adaptive regulation process of enhanced attention to, and nonjudgmental awareness of, present moment experiences.<sup>15</sup> Mindfulness, when defined in this way, is found to be positively associated with psychological well-being<sup>16, 17</sup> and inversely associated with a wide range of psychological outcomes, including depression,<sup>18</sup> anxiety,<sup>18</sup> rumination,<sup>19</sup> and general distress.<sup>19, 20</sup> Of particular importance to the present study, studies have consistently found positive associations between mindfulness and psychological flexibility.<sup>21,</sup>

22 23

Conceptually, these findings are interesting as psychological flexibility and mindfulness reflect functional and process-based understandings of psychopathology.<sup>1</sup> As discussed elsewhere,<sup>6, 21</sup> the two constructs reflect an overarching regulation process of *how* a person contacts and responds to one's internal and external environments in the present moment, not necessarily *what* the person experiences (e.g., hopelessness, fear, etc). A growing body of evidence suggests that various forms of psychopathology are best understood in terms of such underlying regulation processes in a given context.<sup>3, 4, 24</sup>

### **Relationship among Psychological Flexibility, Mindfulness, and Distress**

As stated above, research has consistently supported the negative association between psychological flexibility and major forms of psychological distress,<sup>6, 13</sup> the link between mindfulness and these forms of distress,<sup>2, 10, 19</sup> and the link between psychological flexibility and mindfulness.<sup>21, 22</sup> These findings raise questions about whether mindfulness and psychological flexibility uniquely and separately account for psychological distress or perhaps uniquely and separately account for variance in some forms of psychological distress but not others.

### **Current study**

Following from previous research,<sup>12, 13, 21, 22</sup> the present cross-sectional study first examined the associations among psychological flexibility, mindfulness, and psychological distress (i.e., somatization, depression, anxiety, and general psychological distress) with the expectation that psychological flexibility and mindfulness would be positively associated with one another and negatively associated with all forms of psychological distress. Then, the main study hypothesis was tested by examining the extent to which the two processes, psychological flexibility and mindfulness, accounted for unique variance in psychological distress. It was hypothesized that although mindfulness and psychological flexibility are related constructs and would account for some of the same variance in psychological distress, they would each also account for significant unique variances in distress.

## **Method**

### **Participants**

The current study was conducted at a large, public 4-year university in Georgia. Participants were recruited from undergraduate psychology courses through a web-based research participant pool. Six hundred eighty four participants ( $n_{Female} = 501$ ; 73% female) completed a survey containing several instruments, with a mean completion time for the survey

of approximately 32 minutes ( $SD = 15.75$ ). As employed in previous studies,<sup>11</sup> those who completed the survey in less than 15 minutes or more than 45 minutes were removed from the study because of the questionable validity of their responses. Five hundred fifty participants remained ( $n_{Female} = 413$ ; 75% female). Participants ranged in self-reported age from 16 to 50 years ( $M = 20.97$ ,  $SD = 4.96$ ). Additionally, 56 participants who were aged 26 years old or older were further excluded based on outlier analysis of age. The final participants consisted of 494 college undergraduates ( $n_{Female} = 373$ ; 76% female), ranging in age from 16 to 25 ( $M = 19.55$ ,  $SD = 1.64$ ). The ethnic composition of the sample was representative of the university with 40% ( $n = 195$ ;  $n_{Female} = 147$ ) identifying as “European American,” 28% ( $n = 137$ ;  $n_{Female} = 108$ ) identifying as “African American,” 18% ( $n = 87$ ;  $n_{Female} = 61$ ) identifying as “Asian American/Pacific Islander,” 6% ( $n = 28$ ;  $n_{Female} = 21$ ) identifying as “Hispanic American,” and 8% ( $n = 47$ ;  $n_{Female} = 36$ ) identifying as “bicultural,” “other,” or “Native American”.

### **Procedure and measures**

The current study was approved and monitored by the university Institutional Review Board. Participants who enrolled in the study were asked to complete an anonymous web-based survey. The purpose of the study and instructions for completing the survey were presented at the beginning of the survey. Participants anonymously provided demographic information and completed the measures. The following measures were used to assess psychological distress, psychological flexibility, and mindfulness.

**Psychological Distress.** The *General Health Questionnaire-12* (GHQ-12)<sup>25</sup> is a measure of global psychological distress. Participants are asked to rate frequency with which they experience common types of distress. Using a Likert-scale format, items are scored on a 4-point scale, ranging from 0 (*not at all*) to 3 (*much more than usual*), with a total score derived from the

sum of all responses (e.g., “Have you recently lost much sleep over worry?”). Total scores range from 0 to 36, with higher scores indicating greater distress. A recent study with a non-clinical college undergraduate sample has shown an adequate Chronbach’s alpha of .88.<sup>26</sup> In the present study, Chronbach’s alpha of this measure was .87.

The *Brief Symptom Inventory 18* (BSI-18)<sup>27</sup> is a measure of psychological distress designed to screen for depressive, anxious, and somatic symptoms. The BSI-18 contains 18 items and employs a 5-point Likert scale ranging from 0 (*not at all*) to 4 (*extremely*). The global severity index (GSI) score is derived from the sum of all item scores, ranging from 0 to 72 with greater scores suggesting greater psychological distress. Additionally, scores can be obtained for the somatization (six items; e.g., “faintness”), depression (six items; e.g., “no interest”), and anxiety (six items; e.g., “nervousness”) dimensions. The BSI has been shown to be a reliable and valid measure, with an adequate internal consistency ( $\alpha = .74, .84, .79,$  and  $.89$ , for somatization, depression, anxiety, and GSI, respectively).<sup>27</sup> In the present study Chronbach’s alpha of somatization, depression, anxiety, and GSI were .78, .85, .82, and .91, respectively.

**Psychological flexibility.** The *Acceptance and Action Questionnaire-16* (AAQ-16)<sup>10</sup> was used to measure psychological flexibility for this study. The AAQ is a 16-item questionnaire designed to assess willingness to accept undesirable thoughts and feelings (e.g., “It is OK to feel depressed or anxious”), while acting in a way that is consistent with one’s values and goals (e.g., “I am able to take action on a problem even if I am uncertain of the right thing to do”). The measure employs a 7-point Likert scale, ranging from 1 (*Never true*) to 7 (*Always true*). Total scores range from 16 to 112, with higher scores indicating greater psychological flexibility. Research has indicated that the AAQ has good psychometric properties.<sup>6</sup> In a previous study

conducted with a non-clinical sample,<sup>10</sup> alpha coefficients for this measure ranged from .72 to .79. Chronbach's alpha of this measure in the present study was .62.

**Mindfulness.** The Mindful Attention Awareness Scale (MAAS)<sup>15</sup> is a 15-item, self-report measure, which is designed to assess the frequency of mindlessness, the opposite of the construct of mindfulness, over time (e.g., "It seems I am running automatic without much awareness of what I'm doing"). Participants rate the extent to which they function mindlessly in daily life, using a six-point Likert scale ranging from 1 (*almost always*) to 6 (*almost never*). Total scores range from 15 to 90, with higher scores denoting greater mindfulness. The MAAS has good Cronbach's alpha, ranging from .82 to .87.<sup>15</sup> Chronbach's alpha of MAAS in the present study was .89.

### **Data analysis**

A series of multiple regressions were conducted to investigate the unique role of mindfulness and psychological flexibility on the general and specific forms of distress. Age, gender (i.e., coded as 1 = male, 2 = female), and ethnicity (e.g., coded as 0 = Non-European American, 1 = European American) were covaried on all regression analyses.

## **Results**

### **Associations among Psychological Flexibility, Mindfulness, and Distress**

Descriptive statistics and correlations among the study variables are shown in Table 1. Being a female was also associated with greater general distress, greater somatization, greater anxiety, and lower psychological flexibility. There was a positive association between psychological flexibility and mindfulness. Psychological flexibility (AAQ) and mindfulness (MAAS) were negatively associated with all forms of psychological distress (subscales and GSI of BSI-18 and GHQ).

### **Explaining Variance in Psychological Distress**

Age, gender, and ethnicity were included as covariates in all regression analyses (Table 2). Ethnicity predicted anxiety. Being an ethnic minority was associated with greater levels of anxiety. Mindfulness and psychological flexibility both separately accounted for unique variance in general psychological distress measured with the GHQ-12 and BSI-18 GSI. Mindfulness and psychological flexibility also uniquely and separately accounted for the variance in somatization, depression, and anxiety.

### **Discussion**

Employing an ethnically diverse, non-clinical sample of college students, the present study examined whether mindfulness and psychological flexibility separately accounted for unique variance in somatization, depression, anxiety, and general psychological distress. Consistent with previous findings,<sup>13, 19, 21, 23</sup> the study demonstrated that both mindfulness and psychological flexibility were inversely associated with somatization, depression, anxiety, and general psychological distress. The study also extended the extant literature by demonstrating the unique and distinct variance in each of these forms of psychological distress that is accounted for by psychological flexibility and mindfulness.

The current study has important theoretical implications. First, the elucidation of the significant and distinct roles of these mindfulness and psychological flexibility support process-based explanations for psychopathology.<sup>6, 24, 28</sup> Process-based accounts posit that an individual's responses to internal and external experiences are at least as crucial as the experiences themselves in the onset and maintenance of psychopathology. In particular, the present findings suggest that regulation processes, such as mindful awareness and psychological openness without avoidance, play crucial roles in maintenance of somatization, anxiety,

depression, and general distress. Second, their associations with a range of distress also support the transdiagnostic and unifying nature of mindfulness and psychological flexibility, suggesting their applicability to broader clinical contexts.

Clinically, the present study suggests an important role for mindfulness and psychological flexibility in the treatment of psychological distress. This clinical implication is consistent with recent cognitive and behavioral therapies that incorporate these two processes into their theories and practices.<sup>14, 29</sup> A growing body of evidence has demonstrated that these therapies promote positive clinical outcomes by improving mindfulness and psychological flexibility.<sup>1, 6, 7</sup> The present study concurs with this research suggesting that interventions should not only target psychological symptoms but should also target underlying processes, such as psychological flexibility and mindfulness, and that studies of interventions that target both mindfulness and psychological flexibility might be fruitful.

The current investigation has several notable limitations. Given the use of non-clinical sample, the present study should not be treated as a clinical investigation of psychopathology. The number of variables included in the study was intentionally limited in order to gain a preliminary understanding of the role of mindfulness and psychological flexibility in a range of distress. However, this empirical approach might have undermined the significance of the present findings as recent studies have shown the interaction effects of adaptive and maladaptive regulation strategies on distress. In particular, maladaptive regulation processes, such as rumination and thought suppressions, have been found to be more strongly associated with a range of psychological distress than adaptive regulation strategies,<sup>24, 30</sup> and that an inverse association between adaptive regulation and distress is established only at high levels of

maladaptive strategies.<sup>31</sup> Therefore, it is important to investigate the roles of mindfulness and psychological flexibility along with some of the major maladaptive strategies.

As mentioned elsewhere,<sup>11</sup> the scales used in the present study have not been fully validated across diverse ethnic groups. This concern is particularly to the case with the AAQ-16. Although the AAQ-16 is a most widely used measure of psychological flexibility,<sup>6</sup> it is still unclear whether the measure reflects the construct of psychological flexibility. Given its lower Chronbach's alpha found in the present study, it is important to investigate the construct validity of this measure across diverse populations further.

Similarly, it should be noted that, given the exclusive use of MAAS, the present conceptualization of mindfulness does not encompass other features that are often included in the definitions of mindfulness.<sup>21</sup> In other words, mindfulness in the present study reflects the present moment awareness, but it does not capture other features, such as the absence of impulsivity, non-judgment, and purposeful action.<sup>1, 21</sup> As the latter features of mindfulness overlap with the construct of psychological flexibility, results of the associations among psychological flexibility, distress, and mindfulness are very likely to change should other measures of mindfulness are used.

External validity of the present study is somewhat limited given that data were derived from college students attending an urban university in the southeastern United States. From a socio-cultural perspective, some demographic factors, such as gender role, ethnicity, regional context, and university culture, are likely to shape the variables of the present study in systematic ways. Although gender was covaried out in all analyses, our findings derived from a predominantly female undergraduate sample may not be applicable to more diverse samples,

including those that are less educated, more clinical, or older. Nevertheless, the sample was diverse in terms of ethnicity and social economic status.

Finally, perhaps the largest limitation was the reliance on a cross-sectional and correlational design with the use of self-report measures exclusively. The analytic strategy of the present study did not permit elucidating the direction of associations or making causal inferences about functional associations among the constructs of interest. Therefore, the interpretation of the present findings should be made with cautions. Conceptually, mindfulness and psychological flexibility are regulation processes referring to individual interactions with internal and external experiences in a given moment in a given context. For this reason, the exclusive reliance on self-reported measures is unlikely to capture the dynamic and ongoing nature of these two processes.

### **Conclusion**

This study addresses a novel question, employs a large, ethnically and economically diverse sample, and uses multiple measures of psychological distress. It extends the existing literature on regulation processes underlying a range of distress by suggesting that mindfulness and psychological flexibility separately and independent accounts for unique variance in general and specific forms of distress. The current study also suggests that it is beneficial to continue investigating the role of mindfulness in psychological flexibility and their associations with a range of distress, particularly with treatment studies and studies that can bear out causal relationships among these variables.

**Acknowledgement**

The present cross-sectional study was conducted at Georgia State University, Atlanta, GA. The study had not presented elsewhere. There was no assistance in writing the manuscript or other help that did not merit authorship.

**Author Contributions**

Masuda wrote the first draft of the manuscript and finalized the manuscript. Tully extensively revised and edited the manuscript. All authors contributed to and approved the final manuscript.

**Declaration of Conflicting Interests**

The authors declared no potential conflicts of interests with respect to the authorship and/or publication of this article.

**Funding**

The authors received no financial support for the research and/or authorship of this article.

**Ethical Approval**

The present study was reviewed, approved, and monitored by the institutional review board at the university, with which the authors of the paper is affiliated.

## References

1. Hayes SC, Villatte M, Levin M, et al. Open, aware, and active: Contextual approaches as an emerging trend in the behavioral and cognitive therapies. *Annu Rev Clin Psychol.* 2011;7:141-168.
2. Brown KW, Ryan RM, Creswell JD. Mindfulness: Theoretical foundations and evidence for its salutary effects. *Psychological Inquiry.* 2007;18(4):211-237.
3. Kashdan TB, Rottenberg J. Psychological flexibility as a fundamental aspect of health. *Clin. Psychol. Rev.* 2010;30(4):467-480.
4. Chambers R, Gullone E, Allen NB. Mindful emotion regulation: An integrative review. *Clinical Psychology Review.* 2009;29(6):560-572.
5. Hofmann SG, Sawyer AT, Witt AA, et al. The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *J. Consult. Clin. Psychol.* 2010;78(2):169-183.
6. Hayes SC, Luoma JB, Bond FW, et al. Acceptance and commitment therapy: Model, processes and outcomes. *Behav. Res. Ther.* 2006;44(1):1-25.
7. Shapiro SL, Oman D, Thoresen CE, et al. Cultivating mindfulness: Effects on well-being. *J Clin Psychol.* 2008;64(7):840-862.
8. Bond FW, Bunce D. Mediators of change in emotion-focused and problem-focused worksite stress management interventions. *J. Occup. Health Psychol.* 2000;5(1):156-163.
9. Kashdan TB, Barrios V, Forsyth JP, et al. Experiential avoidance as a generalized psychological vulnerability: Comparisons with coping and emotion regulation strategies. *Behav. Res. Ther.* 2006;44(9):1301-1320.

10. Bond FW, Bunce D. The role of acceptance and job control in mental health, job satisfaction, and work performance. *J. Appl. Psychol.* 2003;88(6):1057-1067.
11. Masuda A, Anderson PL, Wendell JW, et al. Psychological flexibility mediates the relation between self-concealment and negative psychological outcomes. *Personality and Individual Differences.* 2011;50:243-247.
12. Plumb JC, Orsillo SM, Luterek JA. A preliminary test of the role of experiential avoidance in post-event functioning. *J. Behav. Ther. Exp. Psychiatry.* 2004;35(3):245-257.
13. Tull MT, Gratz KL, Salters K, et al. The role of experiential avoidance in posttraumatic stress symptoms and symptoms of depression, anxiety, and somatization. *J. Nerv. Ment. Dis.* 2004;192(11):754-761.
14. Baer RA. *Mindfulness-based treatment approaches: Clinician's guide to evidence base and applications.* San Diego, CA: Elsevier Academic Press; 2006.
15. Brown KW, Ryan RM. The benefits of being present: Mindfulness and its role in psychological well-being. *J. Pers. Soc. Psychol.* 2003;84(4):822-848.
16. Howell AJ, Digdon NL, Buro K. Mindfulness predicts sleep-related self-regulation and well-being. *Personality and Individual Differences.* 2010;48(4):419-424.
17. Howell AJ, Digdon NL, Buro K, et al. Relations among mindfulness, well-being, and sleep. *Personality and Individual Differences.* 2008;45(8):773-777.
18. Roemer L, Lee JK, Salters-Pedneault K, et al. Mindfulness and emotion regulation difficulties in generalized anxiety disorder: Preliminary evidence for independent and overlapping contributions. *Behavior Therapy.* 2009;40(2):142-154.

19. Coffey KA, Hartman M. Mechanisms of action in the inverse relationship between mindfulness and psychological distress. *Complementary Health Practice Review*. 2008;13(2):79-91.
20. Masuda A, Wendell JW, Chou Y-Y, et al. Relationships among self-concealment, mindfulness and negative psychological outcomes in asian american and european american college students. *International Journal for the Advancement of Counselling*. 2010;32(3):165-177.
21. Baer RA, Smith GT, Hopkins J, et al. Using self-report assessment methods to explore facets of mindfulness. *Assessment*. 2006;13(1):27-45.
22. Masuda A, Anderson PL, Sheehan ST. Mindfulness and mental health among african american college students. *Complementary Health Practice Review*. 2009;14(3):115-127.
23. Mitmansgruber H, Beck TN, Höfer S, et al. When you don't like what you feel: Experiential avoidance, mindfulness and meta-emotion in emotion regulation. *Personality and Individual Differences*. 2009;46(4):448-453.
24. Aldao A, Nolen-Hoeksema S, Schweizer S. Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review*. 2010;30(2):217-237.
25. Goldberg D. *Manual of the general health questionnaire*. Windsor: National Foundation for Educational Research; 1978.
26. Masuda A, Wendell JW. Mindfulness mediates the relation between disordered eating-related cognitions and psychological distress. *Eating Behaviors*. 2010;11(4):293-296.
27. Derogatis LR. *Brief symptom inventory-18 (bsi-18) administration, scoring, and procedure manuals*. 3rd ed. Minneapolis: NCS Pearson; 2001.

28. Segal ZV, Teasdale JD, Williams JMG. Mindfulness-based cognitive therapy: Theoretical rationale and empirical status. In: Hayes SC, Follette VM, Linehan MM, eds. *Mindfulness and acceptance: Expanding the cognitive-behavioral tradition*. New York, NY US: Guilford Press; 2004:45-65.
29. Hayes SC, Follette VM, Linehan MM. *Mindfulness and acceptance: Expanding the cognitive-behavioral tradition*. New York, NY US: Guilford Press; 2004.
30. Aldao A, Nolen-Hoeksema S. Specificity of cognitive emotion regulation strategies: A transdiagnostic examination. *Behaviour Research and Therapy*. 2010;48(10):974-983.
31. Aldao A, Nolen-Hoeksema S. When are adaptive strategies most predictive of psychopathology? *J Abnorm Psychol*. 2011.

Table 1

*Means, Standard Deviations, Coefficient Alphas, and Zero-Order Relations between all Variables*

	1	2	3	4	5	6	7	8	9	10
1. Psychological Distress (GHQ)	--									
2. Psychological Distress (BSI-18 GSI)	.64**	--								
3. Somatization (BSI-18 Somatization)	.38**	.80**	--							
4. Depression (BSI-18 Depression)	.68**	.85**	.44**	--						
5. Anxiety (BSI-18 Anxiety)	.54**	.91**	.66**	.67**	--					
6. Psychological Flexibility (AAQ)	-.46**	-.42**	-.24**	-.42**	-.38**	--				
7. Mindfulness (MAAS)	-.43**	-.47**	-.37**	-.42**	-.41**	.39**	--			
8. Age	-.03	-.03	.05	-.00	-.03	.09	.03	--		
9. Gender	.14**	.12**	.12**	.06	.13**	-.14**	-.07	-.06	--	
10. Ethnicity	-.03	.04	.00	.02	.07	.09*	-.01	.07	.00	--
<i>M</i>	12.16	12.85	3.65	4.91	4.29	71.21	57.58			
<i>SD</i>	6.16	10.87	3.81	4.73	4.19	8.93	12.21			
<i>α</i>	.87	.91	.78	.85	.82	.62	.89			

Note:  $N = 494$ , \* $p < .05$ , \*\* $p < .01$ , GHQ = General Health Questionnaire; BSI-18 = Brief Symptom Inventory-18 item; GSI = Global Severity Index; AAQ = Acceptance and Action Questionnaire; MAAS = Mindfulness Attention Awareness Scale

Table 2.

*Regression Analyses to Investigate the Unique Role of Mindfulness and Psychological Flexibility on Various Forms of Distress*

Variable	$\beta$	<i>B</i>	<i>SE B</i>	<i>t</i>	<i>p</i>
<b>General Psychological Distress (GHQ)</b>					
Age	.01	.04	.15	.26	.794
Gender	.07	1.03	.55	1.86	.063
Ethnicity	-.00	-.01	.49	-.03	.980
Mindfulness (MAAS)	-.29	-.15	.02	-7.00	.000
Psychological Flexibility (AAQ)	-.34	-.23	.03	-7.98	.000
$R^2 = .29$					
<b>General Psychological Distress (BSI-18 GSI)</b>					
Age	.06	.42	.25	1.65	.099
Gender	.06	1.42	.97	1.46	.144
Ethnicity	.06	1.26	.85	1.48	.138
Mindfulness (MAAS)	-.36	-.32	.04	-8.68	.000
Psychological Flexibility (AAQ)	-.28	-.34	.05	-6.68	.000
$R^2 = .29$					
<b>Somatization (BSI-18 Somatization)</b>					
Age	.08	.18	.10	1.84	.066
Gender	.08	.72	.37	1.92	.055
Ethnicity	.00	.03	.33	.08	.934
Mindfulness (MAAS)	-.33	-.11	.01	-7.20	.000
Psychological Flexibility (AAQ)	-.11	-.05	.02	-2.26	.019
$R^2 = .16$					
<b>Depression (BSI-18 Depression)</b>					
Age	.03	.09	.11	.76	.447
Gender	-.00	-.03	.43	-.07	.947
Ethnicity	.04	.45	.38	1.18	.240
Mindfulness (MAAS)	-.30	-.12	.02	-7.03	.000
Psychological Flexibility (AAQ)	-.32	-.17	.02	-7.36	.000
$R^2 = .26$					
<b>Anxiety (BSI-18 Anxiety)</b>					
Age	.06	.15	.10	1.52	.129
Gender	.08	.73	.39	1.89	.060
Ethnicity	.09	.79	.34	2.32	.021
Mindfulness (MAAS)	-.30	-.10	.02	-6.94	.000
Psychological Flexibility (AAQ)	-.27	-.13	.02	-6.20	.000
$R^2 = .24$					

Note.  $N = 494$ , GHQ = General Health Questionnaire; BSI = Brief Symptom Inventory; GSI = Global Severity Index, AAQ = Acceptance and Action Questionnaire; MAAS = Mindfulness Attention Awareness Scale.