

Georgia State University

## ScholarWorks @ Georgia State University

---

Kinesiology Faculty Publications

Department of Kinesiology and Health

---

1991

### Creatine Kinase, Creatine Kinase-MM, and the Isoforms of Creatine Kinase-MM Following a Competitive Swimming Workout

Mark H. Bean

Hugh M. Neisler

Walter R. Thompson

*Georgia State University, wrthompson@gsu.edu*

Margot Hall

Tony Young

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.gsu.edu/kin\\_health\\_facpub](https://scholarworks.gsu.edu/kin_health_facpub)



Part of the [Kinesiology Commons](#)

---

#### Recommended Citation

Bean, M.H., H.M. Neisler, W.R. Thompson, M. Hall, T. Young, and J. Pittington: Creatine kinase, creatine kinase-MM, and the isoforms of creatine kinase-MM following a competitive swimming workout, *JOURNAL OF SWIMMING RESEARCH*, 7(3):1991, 21-25. <http://www.swimmingcoach.org/pdf/pub/jsr1990c.pdf>.

This Article is brought to you for free and open access by the Department of Kinesiology and Health at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Kinesiology Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

---

**Authors**

Mark H. Bean, Hugh M. Neisler, Walter R. Thompson, Margot Hall, Tony Young, and John Pittington

THE JOURNAL OF  
**SWIMMING RESEARCH**

ISSN: 0747-5993  
Vol. 7, No. 3, Fall 1991

CONTENTS

Editor's Preview .....	4
ORIGINAL INVESTIGATIONS	
Effectiveness of Biokinetic Training on Swimming Performance in Collegiate Swimmers <i>A.J. Roberts, B. Termin, M.F. Reilly, D.R. Pendergast</i> .....	5
Evaluation of Anaerobic Power and Capacity in Competitive Swimmers <i>D.M. Rohrs, J.M. Stager</i> .....	12
The Effectiveness of Cycle Ergometer Training in Maintaining Aerobic Fitness During Detraining from Competitive Swimming <i>A. Bohrer-Claude, R.L. Sharp</i> .....	17
Creatine Kinase and the Isoforms of Creatine Kinase—MM Following a Competitive Swimming Workout <i>M.H. Bean, H.M. Neister, W.R. Thompson, M. Hall, T. Young, J. Pittington</i> .....	21
Effect of Stroke Rate and Body Mass on $VO_2$ in Crawl Swimming <i>P.P. Klentrou, R.R. Monpetit</i> .....	26
Author Guidelines .....	32



# THE JOURNAL OF SWIMMING RESEARCH



A publication of the ASCA in cooperation with the USS Sports Medicine Committee

## PUBLISHER AND EDITOR

Rick Sharp  
American Swimming Coaches Association  
304 SE 20th St.  
Fort Lauderdale, FL 33316

## BOARD OF ASSOCIATE EDITORS

Jan Prins, Ph.D.  
University of Hawaii  
Honolulu, HI 96822

Joel M. Stager, Ph.D.  
Indiana University  
Bloomington, IN 47405

John P. Troup, Ph.D.  
United States Swimming  
Colorado Springs, CO 80909-5760

Ernest Maglischo, Ph.D.  
California State College, Bakersfield  
Bakersfield, CA 93309

Jack H. Wilmore, Ph.D.  
University of Texas, Austin  
Austin, TX 78712

David L. Costill, Ph.D.  
Ball State University  
Muncie, IN 47306

William Morgan, Ed.D.  
University of Wisconsin  
Madison, WI 53706

## REVIEW BOARD

Cheryl W. Maglischo, Ph.D.  
California State University  
Chico, CA

Ted Becker, Ph.D., R.P.T., A.T.C.  
Evert Pacific Industrial Rehabilitation  
Evert, WA 98201

Robert E. Schleihauf, Ph.D.  
82 Brambach Road  
Scarsdale, NY 10583

James Councilman, Ph.D.  
Indiana University  
Bloomington, IN 47405

## ASSOCIATION OFFICERS

Doug Ingram, President  
Southern Illinois University  
Carbondale, Ill.

Paul Blair  
Little Rock Racquet Club  
Little Rock, AR

Phill Hansel  
University of Houston  
Houston, TX

Jack Nelson  
Fort Lauderdale Swim Team  
Fort Lauderdale, FL

Nort Thornton, Vice President  
University of California  
Berkeley, CA

George Block  
Alamo Area Aquatics  
San Antonio, TX

Ira Klein  
Las Vegas Gold  
Las Vegas, NV

Nancy Jean Schlueter  
Tempe, AZ

Chuck Warner, Vice President  
Sarasota Y Sharks  
Sarasota, FL

Peter Daland  
University of Southern Cal.  
Los Angeles, CA

Kathy McKee  
Dynamo Swim Team  
Tucker, GA

Mark Schubert  
University of Texas  
Austin, TX

Dick Hannula  
Tacoma Swim Club  
Tacoma, WA

Jim Montrella  
The Ohio State University  
Columbus, OH

Richard Shoulberg  
Germantown Academy  
Ft. Washington, PA

THE JOURNAL OF SWIMMING RESEARCH is an official publication of the American Swimming Coaches Association in cooperation with United States Swimming's Sports Medicine Committee. The American Swimming Coaches Association (ASCA) is an educational and professional service organization for swimming coaches. Its national office is located at 304 SE 20th Street, Fort Lauderdale FL 33316. United States Swimming (USS) is the National Governing Body for competitive swimming in the United States. It is a Group A member of the United States Olympic Committee. Its national headquarters is located at 1750 East Boulder Street, Colorado Springs, CO 80909.

COPYRIGHT 1989 American Swimming Coaches Association. It is the policy of the American Swimming Coaches Association to reserve the rights to all articles published in this journal. Coaches and swimming researchers are hereby given permission to make single or multiple copies of single articles for personal scholarly use without explicit written permission, providing that no such copies are sold or used in such a way as to substitute for a subscription to this journal. The American Swimming Coaches Association considers it the obligation of any user to acknowledge the authors and THE JOURNAL OF SWIMMING

RESEARCH for all use of this material as a reprint or reference and that United States Swimming be acknowledged when use is made of research funded by them.

THE JOURNAL OF SWIMMING RESEARCH entitles the subscriber to 4 issues as they are published (not necessarily quarterly) by the American Swimming Coaches Association from 304 SE 20th Street, Fort Lauderdale FL 33316. Telephone (305) 462-6267. Subscription rates are \$55.00 (foreign), \$45.00 (Canada & Mexico), and \$35.00 (USA). The journal is printed by the Western Newspaper Publishing Co., Inc. of Indianapolis, IN.

Acceptance of advertising material does not imply endorsement by the American Swimming Coaches Association or United States Swimming. Author's opinions expressed in the articles are not necessarily those of the staff and/or Boards of the American Swimming Coaches Association or United States Swimming.

POSTMASTER: Send address changes to THE JOURNAL OF SWIMMING RESEARCH, 304 SE 20th Street, Fort Lauderdale FL 33316.

# Creatine Kinase and the Isoforms of Creatine Kinase-MM Following a Competitive Swimming Workout

MARK H. BEAN, PH.D.  
HUGH M. NEISLER, PH.D.  
WALTER R. THOMPSON, PH.D.  
MARGOT HALL, PH.D.  
TONY YOUNG, M.S.  
JOHN PITTINGTON M.S.

*Laboratory of Applied Physiology  
School of Health, Physical Education and Recreation  
Department of Medical Technology  
University of Southern Mississippi  
Hattiesburg, Mississippi 39406*

*Northeast Louisiana University  
Monroe, Louisiana 71209*

## Abstract

*Serum levels of total creatine kinase (CK), CK-MM and the isoforms of CK-MM were measured in 14 male competitive collegiate swimmers. The purpose of this study was to observe changes in CK and the isoforms of CK-MM following a competitive swim training session. Venous blood samples were taken before and immediately following a 5550 yard training session. The main series in the workout consisted of a 27 minute, minimal rest interval set designed to obtain a moderately heavy physical effort. Total CK, composed predominantly of CK-MM, increased significantly ( $p < 0.05$ ) following the swim training session. CK-MM3 (skeletal muscle form) as well as the CK-MM3/CK-MM1 ratio increased significantly ( $p < 0.05$ ) from pre- to post-exercise. CK-MM2 (intermediate form) increased significantly ( $p < 0.05$ ) from pre- to post-exercise with CK-MM1 (serum form) decreasing significantly ( $p < 0.05$ ) from before to after the swim training session. These results indicate that significant changes will occur in serum CK and the isoforms of CK-MM following a competitive swimming workout.*

## Introduction

Creatine kinase (CK) catalyzes the phosphorylation of creatine by adenosine triphosphate (ATP). This rapid reaction creates creatine phosphate (CP) which is an immediate source of energy during exercise. Creatine kinase is found abundantly in skeletal muscle tissue and has demonstrated significant increases following different forms of exercise (9,14,19). Two factors seem to determine the amount of increase in serum enzymes following exercise. First, activities involving eccentric contractions as in downhill running have demonstrated significantly greater increases in serum CK when compared to equal amounts of level running (16). Second, there appears to be a high correlation between increases in serum enzymes and the degree of intramuscular

pressure during a particular exercise (12). These increases have been associated with possible changes in skeletal muscle cell permeability (12).

Studies involving single bouts of swimming have demonstrated conflicting results when serum CK has been observed. Critz and Cunningham (7) and Haralambie and Senser (8) observed increases in serum CK immediately following a 12-minute maximal swimming effort and a 5,200 yard continuous swim, respectively. Conflicting results were reported by Kirwan et al. (10) and Symanski et al. (17) who demonstrated no significant changes in serum CK following swimming activity.

Three sub-types (isoforms) of the CK-MM isoenzyme have been identified and utilized as earlier and more sensitive indicators of acute myocardial infarction (AMI)

(22), as an accurate noninvasive indicator of myocardial reperfusion following AMI (4), and as an effective means of more precisely predicting the time of necrotic onset following the diagnosis of AMI (13). The isoforms which are recognized in various ways in the literature will be identified in the present study as: CK-MM3 (muscle tissue form), CK-MM2 (intermediate form), and CK-MM1 (serum form). CK-MM3, the most cathodic form during electrophoresis, is released directly from muscle tissue into blood where it is converted to CK-MM2 and then to CK-MM1 (1). The CK-MM3 isoform is composed of two unmodified M subunit monomers designated as tissue ( $M_T$ ) and serum ( $M_S$ ) subunits by Wu (20). Serum carboxypeptidase-N (1) post-translationally cleaves a C terminal lysine group from  $M_T$  to form  $M_S$ . Different combinations of  $M_T$  and  $M_S$  create the CK-MM subtypes observed in serum. These alterations allow the CK-MM isoforms to be quantitated by different clinical laboratory methods including high voltage electrophoresis, chromatofocusing, and liquid chromatography (22).

Previous exercise investigations have indicated that the CK-MM isoforms will demonstrate significant changes following exercise activities (2,3,6). To date the isoforms of CK-MM have not been observed following competitive swimming. The purpose of this study was to observe indicators of change in skeletal muscle cells following competitive swim training by measuring total CK and the isoforms of CK-MM. This study also examined the discrepancies between previous investigations involving total CK following swimming.

## Methods

**Subjects/training.** Fourteen male collegiate swimmers between the ages of 17 and 22 years ( $x$  age =  $19.50 \pm 1.40$  yrs.) from a Division I AA university participated in this study. All subjects signed written forms of consent after being advised of the risks and benefits of participating in the study. This procedure is in accordance with the policy statement of the American College of Sports Medicine and the Human Subjects Protection Review Committee. Swim training was designed to include a gradual increase in volume (yardage swum per day) from September through December and included a progressive weight-training program. Data were collected following the first thirteen days of training which averaged 3500 yards swum per day. Blood samples were taken 36 hours following a previous weight-training session.

**Protocol.** The subjects in this study performed a regularly scheduled swim training session of approximately 90 minutes in duration. All subjects arrived at the pool at approximately 5:45 a.m. The training session began at 6:00 a.m. and ended at approximately 7:30 a.m. The subjects began and ended the training session

as a group. The training session consisted of two warm-up swims, two main series, and a cool-down swim totaling 5550 yards. The main series were designed to obtain a moderately heavy physical effort.

**Blood collection.** Venous blood samples were taken before and immediately following the completion of the training session. Samples were drawn by two Registered Medical Technologists using routine venapuncture techniques within ten minutes ( $\pm 3$  minutes) following the swim training session. Seven milliliters of whole blood was obtained from each subject. Blood samples were obtained by a single, aseptic, nontraumatic venapuncture of an antecubital vein using a sterile 18 gauge Vacutainer brand needle. Blood was collected, allowed to clot for 15 minutes, and centrifuged at 3500 rpm to obtain a clear serum sample. The serum sample was then frozen at  $-190^\circ$  F and transported to the institution laboratories conducting this research (approximately four hours away) where they were thawed at  $37^\circ$  C and assayed immediately. Freezing and thawing serum samples have been demonstrated to have no effect on the isoform pattern (20).

**Assay procedures.** Single measures were taken for total CK and the Isoforms of CK-MM. Total CK was determined according to the method of Szasz et al. (18) using a Gilford™ CK assay kit (Gilford Diagnostics, Cleveland, Ohio 44135). Serum was incubated with NADH and observed at 60 second intervals for changes in absorbency at 340 nm. Total CK, in U/L were calculated from the changes in absorbency readings. The CK isoenzymes were determined electrophoretically using the Corning™ ACI System (Corning Laboratories, Palo Alto, CA 94360). Serum samples were applied to agarose gel plates using the formulation of Rosalki (15) and then electrophoresed at 90 volts for 20 minutes. The agarose plates were then incubated with fluorometric substrate for 20 minutes and dried. The dried plates were scanned with a recording fluorometric densitometer and quantified. The CK-MM isoforms were determined by high-voltage electrophoresis in accordance with procedures established by Helena Laboratories (Helena Laboratories, Beaumont, TX). Serum samples were applied to an acetate gel then electrophoresed at 800 volts for 12 minutes. The acetate plate was blotted and layered on a substrate plate (sandwich technique) and then incubated for six minutes at  $45^\circ$  C. Following incubation the plates were separated and dried at  $56^\circ$  C for five minutes. The isoforms were quantitated with a computer integrated scanning densitometer. Typical within-run %CV's for the CK-MM isoforms using high voltage electrophoresis range from 3% to 5% with a sensitivity of roughly 5 U/L per isoform band (21).

**Units of activity.** Activity (U/L) for the CK isoenzyme was derived by multiplying the percent for each isoenzyme fraction (CK-MM, CK-MB, & CK-BB), as

determined electrophoretically, times the total CK activity units. Activity (U/L) for the CK-MM isoforms was derived by multiplying the percent of each isoform (CK-MM3, CK-MM2, and CK-MM1) by the total CK-MM activity. This was also determined electrophoretically.

*Statistical analysis.* One-way analysis of variance (ANOVA) for repeated measures was used to determine any significant changes in pre- to post-data. Changes were considered significant at  $p < 0.05$ .

## Results

All samples contained exclusively the skeletal muscle isoenzyme CK-MM as determined by electrophoresis. Data for total CK and the isoforms of CK-MM are presented in Table 1. Total CK increased significantly

Table 1  
 Total CK, CK-MM, CK-MM isoforms, Hematocrit, Hemoglobin before and immediately following the swim training session.

	Pre training session	Post training session	Prob. F
Total CK (U/L)	219.9 ± 81.6	238.8 ± 87.8	0.0081
CK-MM3 (U/L)	23.1 ± 12.8	39.1 ± 20.1	0.0219
CK-MM2 (U/L)	61.1 ± 30.2	77.3 ± 31.0	0.0005
CK-MM1 (U/L)	135.1 ± 48.6	123.8 ± 47.1	0.0355
MM3/MM1 ratio	0.18 ± .08	.30 ± .14	0.0170

Values are mean ± SE (N=14). CK = creatine kinase; CK-MM = creatine kinase-MM isoenzyme, CK-MM<sub>3</sub> = muscle isoform; CK-MM<sub>2</sub> = intermediate isoform; CK-MM<sub>1</sub> = serum isoform. All changes were statistically significant.

from before to immediately following the swim training session. A significant difference was observed in total CK despite a wide range in both the pre and post samples. CK-MM3, CK-MM2, and the CK-MM3/CK-MM1 ratio also increased following the training session. There was a significant decrease in CK-MM1 following the training session.

## Discussion/Conclusions

Skeletal muscle structural alterations following exercise have been of interest to physiologists as well as athletic coaches for many years. Changes in the concentration of serum enzymes have become a common method of determining alterations of cellular membranes which possibly indicates damage to the muscle cell (12). These alterations of cell structure allow for the release of enzymes into general circulation increasing the concentration of those enzymes in sera which can be quantiated by common clinical laboratory methods (21). The conventional method of estimating damage to skeletal muscle cells has been to observe changes in serum levels of total CK (12).

Research to date indicates that serum CK concentra-

tions will increase following most exercise activities (12). Conflicting results have been reported following swimming. Kirwan et al. (10) saw no significant changes in serum CK immediately following a sub-maximal (95%  $\dot{V}O_{2,max}$ ) front crawl swim while Symanski et al. (17) reported no significant change in serum CK following a one hour continuous submaximal (70%  $\dot{V}O_{2,max}$ ) tethered swim. Haralambie and Sensor (8) reported marked increases in serum CK following a 5,200 yard (90 minute) continuous swim in agreement with Critz and Cunningham (7) who also reported a 16% increase in serum CK following a 12 minute maximal effort swim. The primary difference in the studies reporting conflicting results seems to be the training state of the swimmers observed. In those studies showing increases in total serum CK the subjects were either novice swimmers as in Critz and Cunningham (7) or classified as "generally well trained" as in Haralambie and Sensor (8). A closer look at the swimming speeds of the subjects in the latter study (0.99 m/sec for 5,500 yards) would indicate a less well trained group. In both investigations where no significant change was observed, the subjects were classified as "highly trained" (10,17).

The swimmers in the present study were all collegiate competitive swimmers. However, data were collected after only thirteen days of training. Our results agree with those studies showing an increase in serum CK following a competitive swimming workout. The present investigation also indicates that the training state of swimmers will greatly influence the amount of CK released into general circulation following swimming. Studies in which CK was observed in competitive swimmers during a complete season of training also demonstrated a decrease in resting and immediate post exercise samples as the season progressed (5,11). These studies also indicate that a diminished serum CK will accompany an increase in conditioning.

The present study was also designed to assess changes in serum levels of the isoforms of CK-MM following a competitive swim training session. Previous exercise studies observing the isoforms of CK-MM have demonstrated significant increases in both CK-MM3 and the CK-MM3/CK-MM1 ratio following repeated eccentric contractions of the forearm flexors (2) as well as after repeated isometric contractions of the knee extensor muscles (6). Changes in the CK-MM isoforms in these studies were observed two hours following the cessation of exercise with changes in total CK not occurring until six hours following the completion of the exercise protocol. The results of the present study demonstrated significant increases in CK-MM3, CK-MM2, and the CK-MM3/CK-MM1 ratio within ten minutes following the exercise intervention. A significant decrease in CK-MM1 following the swim training session was also observed.

The present study indicates that there will be a release of CK into serum during an intense swim training session in lesser trained swimmers. The increase in total CK, in this group, argues in favor of an increased release of CK from muscle cells, which is predominately CK-MM. It is, however, interesting to note that there was a concomitant decrease in the absolute amount in CK-MM1. Although it is tempting to hypothesize that there was increased clearance and/or inactivation of the CK-MM1 isoform due to increased metabolism during swimming, the research methodology did not reveal the mechanism. Additional research is needed to address these findings.

The results of the present study also demonstrated that there will be a significant change in serum levels of the isoforms of CK-MM. Also in this study both total CK and the isoforms of CK-MM demonstrated significant changes upon sampling. It is impossible to know whether the change in the isoforms could be detected before the change in total CK as has been demonstrated with other exercise studies (2,6). Serum carboxypeptidase-N, which is responsible for the conversion of the CK-MM isoforms, was not measured in the present study. The authors would like to point out that it is difficult to determine whether the changes in the percentages of the CK-MM isoforms were due in part to changes in serum carboxypeptidase-N. The increase in total CK indicates a release of CK-MM from muscle tissue and therefore indicates that the changes in the isoforms of CK-MM were due to this release.

In conclusion, total CK and the isoforms of CK-MM will demonstrate significant changes following competitive swim training. These changes will be more likely to occur in lesser trained swimmers. Although it is tempting to suggest that these analytes may be useful to the coach as a marker of conditioning, in most cases this would not be practical. More research that could directly quantitate skeletal muscle damage in relation to serum levels of total CK and the isoforms of CK-MM could bring more value to measuring these analytes.

#### Acknowledgments

The authors would like to thank Helena Laboratories, Beaumont, Texas for their assistance with the assays of the CK-MM isoforms. The authors would also like to thank Dr. Alan H.B. Wu, University of Texas Health Science Center in Houston for his suggestions concerning the study.

Address for correspondence: Mark H. Bean, Ph.D., North Mississippi Medical Center, Wellness Center, 1030 South Madison Street, Tupelo, MS 38801.

#### References

1. Abendschein, D.R., H. Serota, T.H. Plummer, K. Amirarian, A. W. Strauss, B.E. Sobel, and A.S. Jaffe. Conversion of MM creatine kinase in human plasma by

- carboxypeptidase N. *J. Lab. Clin. Med.* 110:798-806, 1987.
2. Apple, F.S., Y. Hellston, and P. M. Clarkson. Early detection of skeletal muscle injury by assay of creatine kinase MM isoforms in serum after acute exercise. *Clin. Chem.* 34:1102-1104, 1988.
3. Apple, F.S., M.A. Rogers, and J.L. Ivy. Creatine kinase isoenzyme MM variants in skeletal muscle and plasma from marathon runners. *Clin. Chem.* 32:41-44, 1986.
4. Apple, F.S., S.W. Sharkey, M. Werdick, K.J. Elsperger, and R.T. Tilbury. Analyses of creatine kinase isoenzymes and isoforms in serum to detect reperfusion after acute myocardial infarction. *Clin. Chem.* 33:507-511.
5. Burke, E.R., H.L. Falsetti, R.D. Feld, G.S. Patton, and C.C. Kennedy. Creatine kinase levels in competitive swimmers during a season of training. *Scand. J. Sports Sci.* 4:1-4, 1982.
6. Clarkson, P.M., F.S. Apple, W.C. Byrnes, K.M. McCormick, and P. Triffletti. Creatine kinase isoforms following isometric exercise. *Muscle & Nerve*, 10:41-44, 1987.
7. Critz, J.B. and D.A. Cunningham. Plasma enzyme levels in man after different physical activities. *J. Sports Med.* 12:143-149, 1972.
8. Haralambie, G., and L. Senser. Metabolic changes in man during long-distance swimming. *Eur. J. of Appl Phys.* 43:115-125, 1980.
9. Kanter, M.M., G.R. Lesmes, L.A. Kaminsky, J. La HamSaeger, and N. Nequin. Serum creatine kinase and lactate dehydrogenase changes following an eighty kilometer race. *Eur. J. Appl. Physiol.* 57:60-63, 1988.
10. Kirwan, J.P., D.L. Costill, M.G. Flynn, J.B. Mitchell, W.J. Fink, P.D. Neuffer, and J.A. Houmard. Physiological responses to successive days of intense training in competitive swimmers. *Med. Sci. Sports Exerc.* 20:255-259, 1987.
11. Millard, M.C. Zauner, R. Cade, R. Reese. Serum CPK levels in male and female world class swimmers during a season of training. *J. Swim. Res.* 1:12-16, 1985.
12. Noakes, T.D. Effects of exercise on serum enzyme activities in humans. *Sports Med.* 4:245-267, 1987.
13. Panteghini, M., C. Cuccia, A. Malchiodi, M. Calarco, and N. Pagnoni. Isoforms of creatine kinase MM and MB in acute myocardial infarction: a clinical evaluation. *Clinica Chimica Acta.* 155:1-10, 1986.
14. Rogers, M.A., G.A. Stull, and F.S. Apple. Creatine kinase isoenzyme activities in men and women following a marathon race. *Med. Sci. Sports Exerc.* 17:670-682, 1985.
15. Rosalki, S.B. An improved procedure for serum creatine phosphocreatine determination. *J. Lab. and Clin. Med.* 69:696-705, 1967.
16. Schwane, J.A., J.S. Williams, and J.H. Sloan. Effects of training on delayed muscle soreness and serum creatine kinase activity after running. *Med. Sci. Sports Exerc.* 19:548-589, 1987.
17. Symanski, J.D., R.G. McMurray, L.M. Silverman, B.W. Smith, and A.J. Siegel. Serum creatine and CK-MB isoenzyme responses to acute and prolonged swimming in trained athletes. *Clin. Chem. Acta.* 129:181-187, 1983.
18. Szasz, G., W. Gruber, and E. Berndt. Creatine kinase in serum: I. Determination of optimum reaction conditions. *Clin. Chem.* 22:650-656, 1976.
19. Tiidus, P.M., and C.D. Ianuzzo. Effects of intensity and duration of muscular exercise on delayed soreness and serum enzyme activities. *Med. Sci. Sports Exerc.*



- 15:461-465, 1983.  
20. Wevers, R.A., H.P. Olthuis, J.C. Van Neil, et al. A study on the dimeric structure of creatine kinase. *Clin Chem Acta* 75:377-385, 1977.  
21. Wu, A.H. Creatine kinase isoforms in ischemic heart

- disease. *Clin. Chem.* 35:7-13, 1989.  
22. Wu, A.H., T.G. Gornet, V.H. Wu, R.E. Brockie, and A. Nishikawa. Early diagnosis of acute myocardial infarction by rapid analysis of creatine kinase isoenzyme-3 (CK-MM) sub-types. *Clin. Chem.* 33:358-362, 1987.

# ASCA American Swimming Coaches Association Book List for Swim Coaches



**1. Swimmer and Team**  
by Don Gambriel and Alfred Bay  
Practical, how-to information on the swim team. 117 pages.  
Member Price- \$11.75 Non-members- \$12.75



**2. Swimming Faster**  
by Ernie Maglisco  
The most current comprehensive coaching text. Emphasis on physiology and training, includes bio-mechanics and many other topics. 472 pages.  
Member Price- \$34.75 Non-members- \$35.95




**3. Complete Book of Swimming**  
by Dr. James Counsilman  
Another excellent text covering all aspects of swimming by the most respected Coach/Researcher in Aquatic history. 178 pages.  
Member Price- \$8.75 Non-members- \$9.75



**4. The Aquazoids**  
by Brent Rutemiller  
A complete stretching program which has been successful in increasing muscular range of motion of athletes. Developed by the author over the past 10 years. 39 pages.  
Member Price- \$7.25 Non-members- \$8.25



**5. Coaching Swimming Effectively**  
by Jean Larabee  
An excellent text for the new coach. Outlines for teaching strokes, organizing practices, etc. 112 pages.  
Member Price- \$15.75 Non-members- \$16.75



**6. The Incomplete Book of Training Games and Gimmicks**  
by Bob Steele  
Incorporates fun into the training program by the expert of the topic. How to make workouts more enjoyable.  
Member Price- \$8.25 Non-members- \$9.25



**7. 1990 Championship Swimming Results**  
Edited by Joann Brislin and Doug McDonald  
Championship results from High School State Meets, College Conferences, National Collegiate Championships and Junior/Senior/YMCA Nationals. 474 pages.  
Member Price- \$22.75 Non-members- \$25.75



**8. Climb to the Top**  
by Dennis Pursley  
The USA National Team Director lays out a detailed description of a club swimming program.  
Member Price- \$14.25 Non-members- \$15.25



**9. Tide Teamwork**  
by Don Gambriel and Jonty Skinner  
A model training approach, with categorization of training work that is being done. Coach Skinner adds computer coaching skills to Olympic Coach Gambriel's experience. 192 pages.  
Member Price- \$19.75 Non-members- \$20.75



**10. Parent, Coach & Athlete**  
by John Leonard  
A handbook for age group swimming parents. 1st edition used by coaches around the world as an introduction to the sport for new parents. 2nd edition available only (new, 1990).  
Member Price- \$9.75 Non-members- \$10.75  
Team Discount for 24 or more- \$6.00



**11. Stretching**  
by Bob Anderson  
Revised edition of exercises for swimming, running and other endurance sports. Well illustrated. 192 pages.  
Member Price- \$9.75 Non-members- \$10.75



**12. Mastering Masters Swimming**  
by George Bole  
Excellent resource for Masters Swimmers. Filled with information you'll need to know.  
Member Price- \$14.75 Non-members- \$15.75

Book Number	# of Copies	\$ per Book	Total	Ship the books listed on the left to:	
				NAME _____	
				STREET _____	
				CITY _____ STATE _____ ZIP _____	
				COUNTRY _____	
				No P.O. Boxes please!	
				Payment By: <input type="checkbox"/> Check <input type="checkbox"/> MasterCard <input type="checkbox"/> Visa	
				Acct.No. _____ Exp.Date _____	
				Signature _____	
Total of Above				Clip, mail, and make checks payable to:	
Postage & Handling				<b>ASCA</b>	
Total Enclosed				304 SE 20th Street, Fort Lauderdale, FL 33316	

**POSTAGE & HANDLING**

U.S.A. Add \$2.00 for first item and \$0.75 for each additional item.

Non-U.S.A. Foreign Surface Mail. Add \$3.00 for first item and \$1.25 for each additional item.

Please copy this page for additional ordering space.