

Georgia State University

ScholarWorks @ Georgia State University

---

Neuroscience Institute Faculty Publications

Neuroscience Institute

---

2011

## Serotonergic Lesions of the Periaqueductal Gray, a Primary Source of Serotonin to the Nucleus Paragigantocellularis, Facilitate Sexual Behavior in Male Rats

Joseph J. Normandin

Georgia State University, jnormandin1@gsu.edu

Anne Z. Murphy PhD

Georgia State University, amurphy@gsu.edu

Follow this and additional works at: [https://scholarworks.gsu.edu/neurosci\\_facpub](https://scholarworks.gsu.edu/neurosci_facpub)



Part of the [Neuroscience and Neurobiology Commons](#)

---

### Recommended Citation

Normandin, J. J. & Murphy, A. Z. (2011). Serotonergic Lesions of the Periaqueductal Gray, a Primary Source of Serotonin to the Nucleus Paragigantocellularis, Facilitate Sexual Behavior in Male Rats. *Pharmacology Biochemistry and Behavior*, 98(3), 369–375. doi: 10.1016/j.pbb.2011.01.024

This Article is brought to you for free and open access by the Neuroscience Institute at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Neuroscience Institute Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

1  
2  
3  
4 **Serotonergic lesions of the periaqueductal gray, a primary source of serotonin to**  
5 **the nucleus paragigantocellularis, facilitate sexual behavior in male rats**  
6  
7  
8  
9

10 Joseph J. Normandin<sup>a,b,1</sup>, \*Anne Z. Murphy<sup>a,b</sup>

11  
12 <sup>a</sup>Department of Biology, Center for Behavioral Neuroscience,  
13 Georgia State University, Atlanta, Georgia 30302-4010

14  
15 <sup>b</sup>Neuroscience Institute, Georgia State University,  
16 Atlanta, Georgia 30302-5010  
17  
18  
19  
20

21 January 21, 2011  
22  
23

24 Running Head: Serotonergic vPAG lesions and sexual behavior

25 Present address: <sup>1</sup>Netherlands Institute for Neuroscience  
26 Neuroendocrinology Group  
27 Meibergdreef 47  
28 1105 BA, Amsterdam  
29 The Netherlands  
30 j.normandin@nin.knaw.nl  
31  
32  
33  
34  
35  
36  
37  
38

39 Correspondence to: \*Anne Z. Murphy, Ph.D.  
40 Neuroscience Institute  
41 Georgia State University  
42 PO Box 5030  
43 Atlanta, GA 30302-5030  
44 amurphy@gsu.edu  
45 Phone: 404.413.5332  
46 Fax: 404.413.5301  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

57 Section Editor: Gery Schulteis, Ph.D.  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7 **Abstract**  
8

9 While selective serotonin reuptake inhibitors (SSRIs) are widely used to treat  
10 anxiety and depression, they also produce profound disruptions of sexual function  
11 including delayed orgasm/ejaculation. The nucleus paragigantocellularis (nPGi), a  
12 primary source of inhibition of ejaculation in male rats, contains receptors for serotonin  
13 (5-HT). The ventrolateral periaqueductal gray (vlPAG) provides serotonin to this region,  
14 thus providing an anatomical and neurochemical basis for serotonergic regulation of the  
15 nPGi. We hypothesize that 5-HT acting at the nPGi could underlie the SSRI-induced  
16 inhibition of ejaculation in rodents. To this end, we produced 5-HT lesions of the source  
17 of 5-HT to the nPGi (the vlPAG) and examined sexual behavior. Removing the source  
18 of 5-HT to the nPGi facilitated genital reflexes, but not other aspects of sexual behavior,  
19 consistent with our hypothesis. Namely, 5-HT lesions produced a significant increase in  
20 the mean number of ejaculations and a significant decrease in ejaculation latency as  
21 compared to sham lesioned animals, while latency to mating and the post-ejaculatory  
22 interval did not differ. These data suggest that the serotonergic vlPAG-nPGi pathway is  
23 an important regulatory mechanism for the inhibition of ejaculation in rats, and supports  
24 the hypothesis that this circuit contributes to SSRI-induced inhibition of ejaculation.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 **Keywords:** genital reflexes, ejaculation, copulation, periaqueductal gray, SSRI  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7 **1 Introduction**  
8

9 Selective serotonin reuptake inhibitors (SSRIs) are the most commonly  
10 prescribed class of drugs for individuals with the clinical diagnosis of depression (Arroll  
11 et al., 2009; Koenig and Thase, 2009). SSRIs increase synaptic availability of serotonin  
12 (5-HT) by blocking reuptake at axon terminals (Hiemke and Hartter, 2000; Richelson,  
13 1994). A well-documented side effect of SSRI treatment is sexual dysfunction  
14 (Kennedy and Rizvi, 2009; Schweitzer et al., 2009), with symptoms of decreased libido,  
15 premature ejaculation, delayed ejaculation, and anorgasmia most commonly reported.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 The interaction between 5-HT and the neural circuitry underlying mammalian  
27 sexual behavior is not completely understood, although experiments in the male rat  
28 have suggested that serotonin can either inhibit or facilitate sexual behavior, depending  
29 on the brain site in question and/or the type of receptor targeted (Bitran and Hull, 1987;  
30 de Jong et al., 2006). For example, systemic administration of 5-HT or 5-  
31 hydroxytryptophan (a precursor to 5-HT synthesis) to male rats increases latency to  
32 ejaculation (Ahlenius and Larsson, 1991; Gonzales et al., 1982), whereas the 5-HT  
33 synthesis inhibitor p-chlorophenylalanine decreases ejaculation latency (McIntosh and  
34 Barfield, 1984). In addition, general increases in 5-HT levels through chronic systemic  
35 administration of an SSRI produce delayed ejaculation in male rats (Ahlenius et al.,  
36 1980; de Jong et al., 2005; Mos et al., 1999; Vega Matuszcyk et al., 1998) much like in  
37 humans (Hsu and Shen, 1995). In these studies, 5-HT was administered or  
38 manipulated systematically; therefore the specific brain region(s) mediating the effects  
39 of 5-HT on sexual behavior are not known.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 The nucleus paragigantocellularis (nPGi) of the rostroventrolateral medulla in  
5 rats, and the homologous structure in humans, the nucleus paragigantocellularis  
6 lateralis (Zec and Kinney, 2001), is the hypothesized site of descending inhibition of  
7 genital reflexes in both rats (Marson and McKenna, 1990) and humans (Johnson,  
8 2006). The nPGi receives projections from upstream sites related to sexual behavior,  
9 including the paraventricular nucleus of the hypothalamus, medial preoptic area, and  
10 periaqueductal gray (Murphy and Hoffman, 2001; Murphy et al., 1999; Normandin and  
11 Murphy, 2008), and in turn, projects to the spinal motoneurons (pudendal motoneurons)  
12 innervating the bulbospongiosus and ischiocavernosus muscles (Hermann et al., 2003;  
13 Marson and Carson 3rd, 1999; Marson and McKenna, 1996; Tang et al., 1999), which  
14 are critical for ejaculation in male rats (Miura et al., 2001; Pescatori et al., 1993) and  
15 humans (Hsu et al., 2004; Shafik et al., 2009).

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33 The rat nPGi and the human homologue contain 5-HT immunoreactive cells  
34 (Azmitia and Gannon, 1986; Marson and McKenna, 1992) and application of 5-HT to  
35 the spinal targets of the nPGi in male rats blocks the urethro-genital reflex (Marson and  
36 McKenna, 1992). Similarly, Clément, et al. (2007) found that in male rats, systemic  
37 administration of the SSRI dapoxetine reduced fictive ejaculatory reflexes, which was  
38 reversed by lesions of the nPGi. In addition, lesions of the nPGi in male rats block the  
39 inhibitory effects of systemic fluoxetine (an SSRI) on sexual behavior (Yells et al.,  
40 1994). These studies make it clear that 5-HT acts at the level of the spinal cord to  
41 inhibit genital reflexes, and that 5-HT from the nPGi is a necessary antecedent for  
42 normal inhibition of genital reflexes. It is unclear, however, whether 5-HT may also be  
43 acting at the level of the nPGi to produce the observed effects.

1  
2  
3  
4 The nPGi, contains receptors for 5-HT including the 5-HT<sub>1A</sub> (Thor et al., 1992,  
5 1990), 5-HT<sub>1C</sub> (Hoffman and Mezey, 1989), 5-HT<sub>2A</sub> (Fay and Kubin, 2000; Fonseca et  
6 al., 2001), 5-HT<sub>2C</sub> (Fonseca et al., 2001), and 5-HT<sub>3</sub> (Fonseca et al., 2001) subtypes,  
7 although only the 5-HT<sub>1A</sub> and 5-HT<sub>2C</sub> subtypes are found in abundance (Fonseca et al.,  
8 2001; Thor et al., 1990). The primary source of 5-HT to the nPGi is the ventrolateral  
9 periaqueductal gray (vlPAG). vlPAG neurons project directly to the nPGi (Murphy and  
10 Hoffman, 2001; Normandin and Murphy, 2008), and these vlPAG-nPGi cells co-localize  
11 with 5-HT (Bago et al., 2002) and sexual behavior-induced Fos (Normandin and  
12 Murphy, 2008), suggesting a role for the serotonergic vlPAG-nPGi pathway in male  
13 sexual behavior.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 The present studies were designed to test the hypothesis that the effects of  
29 SSRIs on male sexual behavior are mediated, at least in part, via the nPGi. Our  
30 working hypothesis is that removal of the source of 5-HT to the nPGi would alter the  
31 expression of genital reflexes but not other aspects of sexual behavior.  
32  
33  
34  
35  
36  
37  
38  
39  
40

## 41 **2 Methods**

### 42 **2.1 Subjects**

43 Thirty male Sprague-Dawley Rats (*Rattus Norvegicus*; Charles River; 275-375g)  
44 were same-sex double-housed in a temperature-controlled vivarium in reverse light  
45 (lights on 7:00pm, off 7:00am) with ad libitum access to food and water. All experiments  
46 were approved by the Georgia State Institutional Animal Care and Use Committee, with  
47 pain and suffering minimized in accordance with the Committee's policies.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2.2 Ovariectomy and gonadal-steroid replacement in stimulus females

To induce sexual receptivity, ovariectomized female rats were injected with  $\beta$ -estradiol-3-benzoate (10ug/0.1ml sesame oil s.c.; Sigma Aldrich) 48 hours before testing and progesterone (500ug/0.1ml sesame oil s.c.; Fluka) 4 hours before testing as previously described (Barfield and Lisk, 1970; McEwen et al., 1987; Normandin and Murphy, 2010; Quadagno et al., 1972).

## 2.3 vIPAG 5-HT lesions

All surgeries were performed under aseptic conditions. Male rats (n=30) were anesthetized by inhalation of isoflurane (2-5%; Henry Schein) and placed in a stereotaxic frame. The skull was leveled such that bregma and lambda were at the same dorsoventral plane, and a craniotomy was performed dorsal to the PAG. vIPAG serotonergic-specific lesions were produced by bilateral injection of 5,7-dihydroxytryptamine creatine sulfate (5,7-DHT; 3mg/ml in 0.9% saline 0.1% ascorbic acid; Fluka; n=22 males) using a 1 $\mu$ l Hamilton syringe. Sham animals received equivolume injections of vehicle (0.9% saline 0.1% ascorbic acid; n=8 males). Thirty minutes prior to 5,7-DHT administration, all animals received an injection of desipramine hydrochloride (25mg/kg in 0.9% Saline i.p.; Sigma) to block uptake of the 5-HT neurotoxin by noradrenergic transporters on noradrenergic cells (Bjorklund et al., 1975). The coordinates for the vIPAG were (in mm): AP-8.5 bregma, ML +/-0.75, DV-5.75. 5,7-DHT or vehicle (300nl/side) was slowly injected over 2 minutes and the syringe was left in place for 10 min. before being removed. A small amount of bone wax was placed in the burr-hole and the animals' skin was wound-clipped. Animals received buprenorphine (0.1mg/kg s.c.; Henry Schein) for pain relief and baytril (5mg/kg i.m.;

1  
2  
3  
4 Henry Schein) as a prophylactic antibiotic. Animals recovered in clean heated cages  
5  
6 before being returned to the housing facility.  
7  
8

#### 9 **2.4 Sexual behavior tests**

10  
11 Sexual behavior tests were conducted in acrylic aquariums (61cm x 30.5cm x  
12  
13 30.5cm). Animals were acclimated to the arena prior to the initiation of the experiment  
14  
15 for 10 min. on two consecutive days. Three and six days later, males engaged in 1-  
16  
17 hour mating bouts with stimulus females to gain sexual experience. A third mating bout  
18  
19 on day nine served as the baseline (pre-lesion) measure. Experimental animals then  
20  
21 underwent lesion (or sham lesion) surgery as described above. Following a seven-day  
22  
23 recovery, a final 1-hour mating bout served as the post-lesion measure. All sexual  
24  
25 behavior bouts were recorded and the number of mounts (attempted intromissions  
26  
27 without the stereotyped pelvic thrust associated with an intromission), intromissions,  
28  
29 and ejaculations, as well as the latency to begin mating (from placement of female into  
30  
31 mating chamber to first mount or intromission), ejaculation latency, and post-ejaculatory  
32  
33 interval was scored. Data were tabulated for each ejaculatory series and then  
34  
35 compared across series.  
36  
37  
38  
39  
40  
41  
42

#### 43 **2.5 Perfusion / fixation / tissue preparation**

44  
45 Following a euthanizing dose of SleepAway (0.5ml i.p.; Henry Schein), all  
46  
47 animals were transcardially perfused with 250 ml of 0.9% sodium chloride/2% sodium  
48  
49 nitrite, followed by 300 ml of 4% paraformaldehyde 2.5% acrolein (Polysciences) in 0.1  
50  
51 M phosphate buffer then 150ml of the sodium chloride/sodium nitrite solution. Following  
52  
53 perfusion/fixation, brains were removed and stored at 4°C in 30% sucrose solution until  
54  
55 sectioned. Brains were cut into 25µm coronal sections in a 1:4 series through the PAG  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 with a Leica 2000R freezing microtome and stored free-floating in cryoprotectant-  
5  
6 antifreeze solution (Watson et al., 1986) at -20°C.  
7  
8

## 9           **2.6    Immunohistochemistry**

10           Midbrain tissue sectioned at 25µm in a 1:4 series was processed for NeuN or 5-  
11  
12 HTP for lesion verification as previously described (Loyd and Murphy, 2006; Murphy  
13  
14 and Hoffman, 2001). Briefly, sections were removed from the cryoprotectant solution,  
15  
16 rinsed extensively in potassium phosphate buffered saline (KPBS; pH 7.4), and then  
17  
18 reacted for 20 minutes in 1% sodium borohydride to remove excess aldehydes.  
19  
20 Sections were then incubated in primary antibody solution directed against either NeuN  
21  
22 (Millipore, MAB377; monoclonal, raised in mouse; 1:70,000) or 5-HTP (Immunostar,  
23  
24 24446; polyclonal, raised in rabbit; 1:1,000) in KPBS containing 0.1% Triton-X for 1 hour  
25  
26 at room temperature followed by 48 hours at 4°C. After primary antibody incubation,  
27  
28 tissue was rinsed in KPBS, incubated for 1 hour in biotinylated goat-anti mouse (mouse  
29  
30 anti-NeuN primary antibody) or goat anti-rabbit (rabbit anti-5-HTP primary antibody) IgG  
31  
32 (Jackson Immunoresearch) at a concentration of 1:600, rinsed in KPBS, followed by a  
33  
34 1-hour incubation in avidin-biotin peroxidase complex (Vector Labs, ABC Elite Kit PK-  
35  
36 6100) at a concentration of 1:10. After rinsing in KPBS and sodium acetate (0.175 M;  
37  
38 pH 6.5), NeuN and 5-HTP were visualized as a black reaction product using nickel  
39  
40 sulfate intensified 3,3'-diaminobenzidine solution containing 0.08% hydrogen peroxide  
41  
42 in sodium acetate buffer. The reaction product was terminated after approximately 15  
43  
44 minutes by rinsing in sodium acetate buffer. Sections were mounted out of saline onto  
45  
46 gelatin-subbed slides, air dried overnight, dehydrated in a series of graded alcohols,  
47  
48 cleared in HistoClear, and cover-slipped using Permount.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2.7 Lesion analysis

5,7-DHT lesion and vehicle sham lesion sites were verified by light-microscopic analysis of vIPAG-containing brain sections immunohistochemically stained for NeuN or 5-HTP. The position of the bottom of the injection track was noted in the NeuN stained sections, and the absence or presence of 5-HTP was noted in 5-HTP stained sections. Only those injections centered in the vIPAG and with an absence of 5-HTP immunoreactivity in the vIPAG were considered for analysis.

## 2.8 Statistical analysis

Planned comparisons of mean sexual behavior within the lesion or sham groups were analyzed with two-tailed paired t-tests. Between group (lesion vs. sham) comparisons of means of sexual behavior were conducted using independent sample t-tests (two-tailed). All statistical comparisons were made with the alpha value set at 0.05.

# 3 Results

## 3.1 Lesion Verification

An example of 5-HTP immunoreactivity in a lesioned animal and sham-lesioned animal is provided in Figure 1. Serotonergic lesions were restricted to the vIPAG in the lesion (n=9) group, with slight spread to the dorsal raphe nucleus, and encompassing an approximate 1mm<sup>3</sup> volume. Syringe paths in our sham-lesion group (n=7) ended just dorsal to, or within the vIPAG. The two animals with misplaced lesions (dorsal to the vIPAG) did not exhibit any obvious sexual behavior changes. In addition, no obvious changes in non-sexual behaviors were noted in any group.

### 3.2 Sexual Behavior

Serotonergic lesions of the vIPAG did not affect behaviors not directly related to genital reflex function. For example, there was no change from baseline in the mean latency to initiate copulation, or in the mean post-ejaculatory interval within either group (Figure 2). By contrast, serotonergic lesions of the vIPAG facilitated male sexual behavior in many measures related to genital reflex functions. While there was no change from baseline in the mean **total** of mounts or intromissions within either group (Figure 3A & B), there was an increase from baseline in the mean number of ejaculations within the lesion group ( $p=0.005$ ) but not the sham group ( $p=1.000$ ). In addition, the mean number of ejaculations in the sexual behavior test was significantly greater in the lesion group than in the sham group ( $p=0.039$ , Figure 3C). There was a decrease from baseline in the mean number of intromissions required for ejaculation within both groups (lesion:  $p=0.004$ ; sham:  $p=0.039$ ; Figure 3D). There was a trend approaching statistical significance for a decrease from baseline in the mean ejaculation latency within the lesion group ( $p=0.054$ ; Figure 3E), and this trend is supported by the significantly shorter mean ejaculation latency in the sexual behavior test in the lesion group versus the sham group ( $p=0.026$ ; Figure 3E).

## 4 Discussion

Overall, serotonergic lesions of the vIPAG in male rats produced a facilitation of sexual behavior. In particular, 5-HT lesions of the vIPAG significantly increased the number of ejaculations during the mating bout, and there was a trend for a decrease in the mean latency to ejaculation. 5-HT vIPAG lesioned animals also had significantly

1  
2  
3  
4 shorter ejaculation latencies as compared to sham. There were no differences in the  
5  
6 mean latency to mate, or in the post-ejaculatory interval for either group indicating that  
7  
8 the effect of serotonergic vIPAG lesions is limited to those behaviors directly associated  
9  
10 with genital reflexes.  
11  
12

13  
14 The facilitation of sexual behavior we observed was specific to those measures  
15  
16 associated with genital reflex function. Given the anatomical connections between  
17  
18 vIPAG 5-HT cells and the nPGi (Bago et al., 2002; Normandin and Murphy, 2008;  
19  
20 Underwood et al., 1999), as well as the concordance of effects in both vIPAG 5-HT  
21  
22 lesions and nPGi lesions (Liu and Sachs, 1999; Yells et al., 1992), it is likely that the  
23  
24 effects observed here are dependent on vIPAG 5-HT neurotransmission to the nPGi. In  
25  
26 the context of a normally behaving animal, one would expect that 5-HT from the vIPAG  
27  
28 serves to enhance nPGi activity, thereby increasing descending inhibition of genital  
29  
30 reflexes. This serotonergic signal from the vIPAG would be particularly active in  
31  
32 behavioral contexts where sexual behavior would not be appropriate (e.g. a “fight-or-  
33  
34 flight” response) and likely “de-activated” when mating conditions were optimal (e.g.  
35  
36 availability of a sexually receptive conspecific in the absence of a predator). This  
37  
38 serotonergic vIPAG-nPGi pathway may effectively act as a gating mechanism for  
39  
40 descending inhibition of genital reflexes.  
41  
42  
43  
44  
45  
46  
47

48 Regulation of nPGi function (and thereby genital reflexes) through vIPAG is an  
49  
50 interesting prospect, as midbrain periaqueductal gray (PAG) cells are known to be  
51  
52 involved in the coordination of cardiovascular responses to anxiety and stress (Johnson  
53  
54 et al., 2004; Moraes et al., 2008; Murphy et al., 1995), nociception (Haghparast and  
55  
56 Ahmad-Molaei, 2009; Loyd et al., 2007), as well as social behaviors (Lonstein and  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Stern, 1998; Pavesi et al., 2007), and could therefore signal the current behavioral  
5  
6 context to the nPGi. With respect to sexual behavior, we have previously found that the  
7  
8 medial preoptic area (MPOA) of the hypothalamus, a region critical in the expression of  
9  
10 male sexual behavior (Arendash and Gorski, 1983; Malsbury, 1971), including genital  
11  
12 reflexes (Giuliano et al., 1996; Marson and McKenna, 1994), provides input to the nPGi  
13  
14 both directly (Normandin and Murphy, 2008) and through a PAG relay (Murphy and  
15  
16 Hoffman, 2001). This MPOA-PAG pathway may provide necessary input to the nPGi  
17  
18 regarding sexual “tone.” Indeed electrolytic lesions of the PAG, including the vPAG,  
19  
20 blocks normal MPOA-elicited bulbospongiosus contractions (Marson, 2004), suggesting  
21  
22 that the MPOA-vPAG-nPGi circuit is required for the elicitation of genital reflexes.  
23  
24  
25  
26  
27

28  
29 The medially adjacent dorsal raphe nucleus (DR) also contains serotonergic  
30  
31 neurons that may have been affected by our lesions. However, results from previous  
32  
33 work suggest that our observed findings are primarily the result of vPAG lesions, and  
34  
35 not due to the spread of the toxin into the DR. For example, in male rats, lesions of the  
36  
37 DR, or application of 5-HT or 5-HT agonists to the DR have no effect on male  
38  
39 ejaculatory behavior (Albinsson et al., 1996; Fernandez-Guasti et al., 1992; Hillegaart,  
40  
41 1991), although one study has reported that 5-HT DR manipulations shortened both the  
42  
43 latency to ejaculate and the post-ejaculatory interval (McIntosh and Barfield, 1984).  
44  
45 Interestingly, lesions of the DR also have been found to facilitate lordosis in male rats  
46  
47 treated with estradiol (Kakeyama and Yamanouchi, 1992), leading to speculation that  
48  
49 the DR is not involved in male typical sexual behavior per se, but rather inhibits female-  
50  
51 typical behavior in male rats.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 5 Conclusions

Serotonergic lesions of the vIPAG in male rats facilitated genital reflex function in a manner similar to nPGi lesions. This result lends evidence to the functional serotonergic connectivity between the vIPAG and nPGi. This work has implications for the occurrence of SSRI-induced delayed ejaculation in human males, and in particular, suggests that these drugs may be producing an effect by increasing serotonergic neurotransmission within the vIPAG-nPGi pathway. Limiting serotonergic drugs for mood disorders in humans to receptors and loci that influence mood regulation systems, but not sexual behavior systems is an emerging area of research (Baldwin et al., 2006; Breuer et al., 2008; Kennedy and Rizvi, 2010). While it has been known that SSRIs can inhibit genital reflexes in male rats at the level of the spinal cord (Marson and McKenna, 1992), our work implies that this could also occur through the nPGi via vIPAG 5-HT input. Thus, in developing treatments for depression and anxiety, clinicians must consider both the spinal and supraspinal targets of serotonergic drugs in this genital reflex circuitry. In addition, SSRIs are often prescribed to men experiencing premature ejaculation (Hatzimouratidis et al., 2010; Hellstrom, 2009). A global increase in 5-HT is neither warranted nor desired in these patients, as other side effects may be incurred (Haddad and Dursun, 2008; Hellstrom, 2009) as a result of treatment. Targeting the serotonergic vIPAG pathway to the nPGi with appropriately specific drugs may be one way to circumvent global treatment of these patients with SSRIs. However, it remains to be elucidated which 5-HT receptor subtypes on nPGi cells are responsible for modulating nPGi activity.

1  
2  
3  
4  
5  
6  
7 **References**  
8

9 Ahlenius, S, Larsson, K. Opposite effects of 5-methoxy-N,N-di-methyl-tryptamine and 5-  
10 hydroxytryptophan on male rat sexual behavior. *Pharmacol Biochem Behav*  
11 1991;38:201-205.  
12

13 Ahlenius, S, Larsson, K, Svensson, L. Further evidence for an inhibitory role of central  
14 5-HT in male rat sexual behavior. *Psychopharmacology (Berl)* 1980;68:217-220.  
15

16 Albinsson, A, Andersson, G, Andersson, K, Vega-Matuszczyk, J, Larsson, K. The  
17 effects of lesions in the mesencephalic raphe systems on male rat sexual behavior and  
18 locomotor activity. *Behav Brain Res* 1996;80:57-63.  
19

20 Arendash, GW, Gorski, RA. Effects of discrete lesions of the sexually dimorphic nucleus  
21 of the preoptic area or other medial preoptic regions on the sexual behavior of male  
22 rats. *Brain Res Bull* 1983;10:147-154.  
23

24 Arroll, B, Elley, CR, Fishman, T, Goodyear-Smith, FA, Kenealy, T, Blashki, G, Kerse, N,  
25 Macgillivray, S. Antidepressants versus placebo for depression in primary care.  
26 *Cochrane Database Syst Rev* 2009:CD007954.  
27

28 Azmitia, EC, Gannon, PJ. The primate serotonergic system: a review of human and  
29 animal studies and a report on *Macaca fascicularis*. *Adv Neurol* 1986;43:407-468.  
30

31 Bago, M, Marson, L, Dean, C. Serotonergic projections to the rostroventrolateral  
32 medulla from midbrain and raphe nuclei. *Brain Res* 2002;945:249-258.  
33

34 Baldwin, D, Bridgman, K, Buis, C. Resolution of sexual dysfunction during double-blind  
35 treatment of major depression with reboxetine or paroxetine. *J Psychopharmacol*  
36 2006;20:91-96.  
37

38 Barfield, MA, Lisk, RD. Advancement of behavioral estrus by subcutaneous injection of  
39 progesterone in the 4-day cyclic rat. *Endocrinology* 1970;87:1096-1098.  
40

41 Bitran, D, Hull, EM. Pharmacological analysis of male rat sexual behavior. *Neurosci*  
42 *Biobehav Rev* 1987;11:365-389.  
43

44 Bjorklund, A, Baumgarten, HG, Rensch, A. 5,7-Dihydroxytryptamine: improvement of its  
45 selectivity for serotonin neurons in the CNS by pretreatment with desipramine. *J*  
46 *Neurochem* 1975;24:833-835.  
47

48 Breuer, ME, Chan, JS, Oosting, RS, Groenink, L, Korte, SM, Campbell, U, Schreiber, R,  
49 Hania, T, Snoeren, EM, Waldinger, M, Olivier, B. The triple monoaminergic reuptake  
50 inhibitor DOV 216,303 has antidepressant effects in the rat olfactory bulbectomy model  
51 and lacks sexual side effects. *Eur Neuropsychopharmacol* 2008;18:908-916.  
52

1  
2  
3  
4 Clement, P, Bernabe, J, Gengo, P, Denys, P, Laurin, M, Alexandre, L, Giuliano, F.  
5 Supraspinal site of action for the inhibition of ejaculatory reflex by dapoxetine. *Eur Urol*  
6 2007;51:825-832.

7  
8  
9 de Jong, TR, Pattij, T, Veening, JG, Dederen, PJ, Waldinger, MD, Cools, AR, Olivier, B.  
10 Citalopram combined with WAY 100635 inhibits ejaculation and ejaculation-related Fos  
11 immunoreactivity. *Eur J Pharmacol* 2005;509:49-59.

12  
13 de Jong, TR, Veening, JG, Waldinger, MD, Cools, AR, Olivier, B. Serotonin and the  
14 neurobiology of the ejaculatory threshold. *Neurosci Biobehav Rev* 2006;30:893-907.

15  
16 Fay, R, Kubin, L. Pontomedullary distribution of 5-HT<sub>2A</sub> receptor-like protein in the rat. *J*  
17 *Comp Neurol* 2000;418:323-345.

18  
19  
20 Fernandez-Guasti, A, Escalante, AL, Ahlenius, S, Hillegaart, V, Larsson, K. Stimulation  
21 of 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors in brain regions and its effects on male rat sexual  
22 behaviour. *Eur J Pharmacol* 1992;210:121-129.

23  
24 Fonseca, MI, Ni, YG, Dunning, DD, Miledi, R. Distribution of serotonin 2A, 2C and 3  
25 receptor mRNA in spinal cord and medulla oblongata. *Brain Res Mol Brain Res*  
26 2001;89:11-19.

27  
28  
29 Giuliano, F, Rampin, O, Brown, K, Courtois, F, Benoit, G, Jardin, A. Stimulation of the  
30 medial preoptic area of the hypothalamus in the rat elicits increases in intracavernous  
31 pressure. *Neurosci Lett* 1996;209:1-4.

32  
33  
34 Gonzales, G, Mendoza, L, Ruiz, J, Torrejon, J. A demonstration that 5-  
35 hydroxytryptamine administered peripherally can affect sexual behavior in male rats.  
36 *Life Sci* 1982;31:2775-2781.

37  
38 Haddad, PM, Dursun, SM. Neurological complications of psychiatric drugs: clinical  
39 features and management. *Hum Psychopharmacol* 2008;23 Suppl 1:15-26.

40  
41 Haghparast, A, Ahmad-Molaei, L. Effects of electrolytic lesion of dorsolateral  
42 periaqueductal gray on analgesic response of morphine microinjected into the nucleus  
43 cuneiformis in rat. *Neurosci Lett* 2009;451:165-169.

44  
45  
46 Hatzimouratidis, K, Amar, E, Eardley, I, Giuliano, F, Hatzichristou, D, Montorsi, F, Vardi,  
47 Y, Wespes, E. Guidelines on Male Sexual Dysfunction: Erectile Dysfunction and  
48 Premature Ejaculation. *Eur Urol* 2010.

49  
50  
51 Hellstrom, WJ. Emerging treatments for premature ejaculation: focus on dapoxetine.  
52 *Neuropsychiatr Dis Treat* 2009;5:37-46.

53  
54 Hermann, GE, Holmes, GM, Rogers, RC, Beattie, MS, Bresnahan, JC. Descending  
55 spinal projections from the rostral gigantocellular reticular nuclei complex. *J Comp*  
56 *Neurol* 2003;455:210-221.

57  
58  
59 Hiemke, C, Hartter, S. Pharmacokinetics of selective serotonin reuptake inhibitors.  
60 *Pharmacol Ther* 2000;85:11-28.



1  
2  
3  
4 Hillegaart, V. Effects of local application of 5-HT and 8-OH-DPAT into the dorsal and  
5 median raphe nuclei on core temperature in the rat. *Psychopharmacology (Berl)*  
6 1991;103:291-296.  
7  
8 Hoffman, BJ, Mezey, E. Distribution of serotonin 5-HT<sub>1C</sub> receptor mRNA in adult rat  
9 brain. *FEBS Lett* 1989;247:453-462.  
10  
11 Hsu, GL, Hsieh, CH, Wen, HS, Hsu, WL, Wu, CH, Fong, TH, Chen, SC, Tseng, GF.  
12 Anatomy of the human penis: the relationship of the architecture between skeletal and  
13 smooth muscles. *J Androl* 2004;25:426-431.  
14  
15 Hsu, JH, Shen, WW. Male sexual side effects associated with antidepressants: a  
16 descriptive clinical study of 32 patients. *Int J Psychiatry Med* 1995;25:191-201.  
17  
18 Johnson, PL, Lightman, SL, Lowry, CA. A functional subset of serotonergic neurons in  
19 the rat ventrolateral periaqueductal gray implicated in the inhibition of  
20 sympathoexcitation and panic. *Ann N Y Acad Sci* 2004;1018:58-64.  
21  
22 Johnson, RD. Descending pathways modulating the spinal circuitry for ejaculation:  
23 effects of chronic spinal cord injury. *Prog Brain Res* 2006;152:415-426.  
24  
25 Takeyama, M, Yamanouchi, K. Lordosis in male rats: the facilitatory effect of  
26 mesencephalic dorsal raphe nucleus lesion. *Physiol Behav* 1992;51:181-184.  
27  
28 Kennedy, SH, Rizvi, S. Sexual dysfunction, depression, and the impact of  
29 antidepressants. *J Clin Psychopharmacol* 2009;29:157-164.  
30  
31 Kennedy, SH, Rizvi, SJ. Agomelatine in the Treatment of Major Depressive Disorder:  
32 Potential for Clinical Effectiveness. *CNS Drugs* 2010.  
33  
34 Koenig, AM, Thase, ME. First-line pharmacotherapies for depression - what is the best  
35 choice? *Pol Arch Med Wewn* 2009;119:478-486.  
36  
37 Liu, YC, Sachs, BD. Erectile function in male rats after lesions in the lateral  
38 paragigantocellular nucleus. *Neurosci Lett* 1999;262:203-206.  
39  
40 Lonstein, JS, Stern, JM. Site and behavioral specificity of periaqueductal gray lesions  
41 on postpartum sexual, maternal, and aggressive behaviors in rats. *Brain Res*  
42 1998;804:21-35.  
43  
44 Loyd, DR, Morgan, MM, Murphy, AZ. Morphine preferentially activates the  
45 periaqueductal gray-rostral ventromedial medullary pathway in the male rat: a potential  
46 mechanism for sex differences in antinociception. *Neuroscience* 2007;147:456-468.  
47  
48 Loyd, DR, Murphy, AZ. Sex differences in the anatomical and functional organization of  
49 the periaqueductal gray-rostral ventromedial medullary pathway in the rat: a potential  
50 circuit mediating the sexually dimorphic actions of morphine. *J Comp Neurol*  
51 2006;496:723-738.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Malsbury, CW. Facilitation of male rat copulatory behavior by electrical stimulation of  
5 the medial preoptic area. *Physiol Behav* 1971;7:797-805.  
6  
7 Marson, L. Lesions of the periaqueductal gray block the medial preoptic area-induced  
8 activation of the urethro-genital reflex in male rats. *Neurosci Lett* 2004;367:278-282.  
9  
10 Marson, L, Carson 3rd, CC. Central Nervous System Innervation of the Penis, Prostate,  
11 and Perineal Muscles: A Transneuronal Tracing Study. *Mol Urol* 1999;3:43-50.  
12  
13 Marson, L, McKenna, KE. CNS cell groups involved in the control of the  
14 ischiocavernosus and bulbospongiosus muscles: a transneuronal tracing study using  
15 pseudorabies virus. *J Comp Neurol* 1996;374:161-179.  
16  
17 Marson, L, McKenna, KE. The identification of a brainstem site controlling spinal sexual  
18 reflexes in male rats. *Brain Res* 1990;515:303-308.  
19  
20 Marson, L, McKenna, KE. A role for 5-hydroxytryptamine in descending inhibition of  
21 spinal sexual reflexes. *Exp Brain Res* 1992;88:313-320.  
22  
23 Marson, L, McKenna, KE. Stimulation of the hypothalamus initiates the urethro-genital  
24 reflex in male rats. *Brain Res* 1994;638:103-108.  
25  
26 McEwen, BS, Jones, KJ, Pfaff, DW. Hormonal control of sexual behavior in the female  
27 rat: molecular, cellular and neurochemical studies. *Biol Reprod* 1987;36:37-45.  
28  
29 McIntosh, TK, Barfield, RJ. Brain monoaminergic control of male reproductive behavior.  
30 I. Serotonin and the post-ejaculatory refractory period. *Behav Brain Res* 1984;12:255-  
31 265.  
32  
33 Miura, T, Kondo, Y, Akimoto, M, Sakuma, Y. Electromyography of male rat perineal  
34 musculature during copulatory behavior. *Urol Int* 2001;67:240-245.  
35  
36 Moraes, CL, Bertoglio, LJ, Carobrez, AP. Interplay between glutamate and serotonin  
37 within the dorsal periaqueductal gray modulates anxiety-related behavior of rats  
38 exposed to the elevated plus-maze. *Behav Brain Res* 2008;194:181-186.  
39  
40 Mos, J, Mollet, I, Tolboom, JT, Waldinger, MD, Olivier, B. A comparison of the effects of  
41 different serotonin reuptake blockers on sexual behaviour of the male rat. *Eur*  
42 *Neuropsychopharmacol* 1999;9:123-135.  
43  
44 Murphy, AZ, Ennis, M, Rizvi, TA, Behbehani, MM, Shipley, MT. Fos expression induced  
45 by changes in arterial pressure is localized in distinct, longitudinally organized columns  
46 of neurons in the rat midbrain periaqueductal gray. *J Comp Neurol* 1995;360:286-300.  
47  
48 Murphy, AZ, Hoffman, GE. Distribution of gonadal steroid receptor-containing neurons  
49 in the preoptic-periaqueductal gray-brainstem pathway: a potential circuit for the  
50 initiation of male sexual behavior. *J Comp Neurol* 2001;438:191-212.  
51  
52 Murphy, AZ, Rizvi, TA, Ennis, M, Shipley, MT. The organization of preoptic-medullary  
53 circuits in the male rat: evidence for interconnectivity of neural structures involved in  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 reproductive behavior, antinociception and cardiovascular regulation. *Neuroscience*  
5 1999;91:1103-1116.  
6

7  
8 Normandin, JJ, Murphy, AZ. Excitotoxic lesions of the nucleus paragigantocellularis  
9 facilitate male sexual behavior but attenuate female sexual behavior in rats.  
10 *Neuroscience* 2010;In press.  
11

12 Normandin, JJ, Murphy, AZ. Nucleus paragigantocellularis afferents in male and female  
13 rats: organization, gonadal steroid receptor expression, and activation during sexual  
14 behavior. *J Comp Neurol* 2008;508:771-794.  
15

16 Pavesi, E, Enck, MC, De Toledo, CA, Terenzi, MG. Disruption of maternal behaviour by  
17 acute conspecific interaction induces selective activation of the lateral periaqueductal  
18 grey. *Eur J Neurosci* 2007;26:2055-2065.  
19  
20

21 Paxinos, G, Watson, C. *The rat brain in stereotaxic coordinates*. Amsterdam ; Boston:  
22 Elsevier Academic Press, 2005.  
23

24 Pescatori, ES, Calabro, A, Artibani, W, Pagano, F, Triban, C, Italiano, G. Electrical  
25 stimulation of the dorsal nerve of the penis evokes reflex tonic erections of the penile  
26 body and reflex ejaculatory responses in the spinal rat. *J Urol* 1993;149:627-632.  
27  
28

29 Quadagno, DM, McCullough, J, Langan, R. The effect of varying amounts of exogenous  
30 estradiol benzoate on estrous behavior in the rat. *Horm Behav* 1972;3:175-179.  
31

32 Richelson, E. The pharmacology of antidepressants at the synapse: focus on newer  
33 compounds. *J Clin Psychiatry* 1994;55 Suppl A:34-39; discussion 40-31, 98-100.  
34

35 Schweitzer, I, Maguire, K, Ng, C. Sexual side-effects of contemporary antidepressants:  
36 review. *Aust N Z J Psychiatry* 2009;43:795-808.  
37  
38

39 Shafik, A, Shafik, AA, El Sibai, O, Shafik, IA. Electromyographic study of ejaculatory  
40 mechanism. *Int J Androl* 2009;32:212-217.  
41

42 Tang, Y, Rampin, O, Giuliano, F, Ugolini, G. Spinal and brain circuits to motoneurons of  
43 the bulbospongiosus muscle: retrograde transneuronal tracing with rabies virus. *J Comp*  
44 *Neurol* 1999;414:167-192.  
45

46 Thor, KB, Blitz-Siebert, A, Helke, CJ. Autoradiographic localization of 5HT1 binding  
47 sites in the medulla oblongata of the rat. *Synapse* 1992;10:185-205.  
48  
49

50 Thor, KB, Blitz-Siebert, A, Helke, CJ. Discrete localization of high-density 5-HT1A  
51 binding sites in the midline raphe and parapyramidal region of the ventral medulla  
52 oblongata of the rat. *Neurosci Lett* 1990;108:249-254.  
53

54 Underwood, MD, Arango, V, Bakalian, MJ, Ruggiero, DA, Mann, JJ. Dorsal raphe  
55 nucleus serotonergic neurons innervate the rostral ventrolateral medulla in rat. *Brain*  
56 *Res* 1999;824:45-55.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Vega Matuszcyk, J, Larsson, K, Eriksson, E. The selective serotonin reuptake inhibitor  
5 fluoxetine reduces sexual motivation in male rats. *Pharmacol Biochem Behav*  
6 1998;60:527-532.  
7

8  
9 Watson, RE, Jr., Wiegand, SJ, Clough, RW, Hoffman, GE. Use of cryoprotectant to  
10 maintain long-term peptide immunoreactivity and tissue morphology. *Peptides*  
11 1986;7:155-159.  
12

13 Yells, DP, Hendricks, SE, Prendergast, MA. Lesions of the nucleus  
14 paragigantocellularis: effects on mating behavior in male rats. *Brain Res* 1992;596:73-  
15 79.  
16

17  
18 Yells, DP, Prendergast, MA, Hendricks, SE, Nakamura, M. Fluoxetine-induced inhibition  
19 of male rat copulatory behavior: modification by lesions of the nucleus  
20 paragigantocellularis. *Pharmacol Biochem Behav* 1994;49:121-127.  
21

22 Zec, N, Kinney, HC. Anatomic relationships of the human nucleus paragigantocellularis  
23 lateralis: a Dil labeling study. *Auton Neurosci* 2001;89:110-124.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7 **Figure 1 - Photomicrograph of sham lesion and lesion sites**

8  
9 A photomicrograph of 5-hydroxytryptophan (5-HTP) immunolabeled cells of the  
10 ventrolateral periaqueductal gray (vIPAG) and dorsal raphe (DR) is shown on the left,  
11 with a diagram (modified from Paxinos and Watson, 2005) at the approximate level of  
12 the tissue section shown on the right. Intra-vIPAG injection of vehicle (photomicrograph  
13 left) preserved 5-HTP immunoreactivity in the vIPAG. Intra-vIPAG injection of 5,7-  
14 dihydroxytryptamine (photomicrograph right) markedly reduced immunoreactivity for 5-  
15 HTP in the vIPAG. Lesion sites are indicating by the gray circles in the diagram.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26 Aq=cerebral aqueduct.  
27  
28  
29  
30

31 **Figure 2 - Measures of sexual behavior unrelated to genital reflex function**

32  
33 Serotonergic lesions of the vIPAG did not alter the latency to mating (A) or post-  
34 ejaculatory interval (B) from baseline to sexual behavior test within either the sham or  
35 lesion groups. Error bars = standard error of the mean (SEM).  
36  
37  
38  
39  
40  
41  
42

43 **Figure 3 - Measures of sexual behaviors related to genital reflex function**

44  
45 Serotonergic lesions of the vIPAG did not alter the mean number of mounts (A) or  
46 intromissions (B) from baseline to sexual behavior test within either group. However,  
47 such lesions significantly increased the mean number of ejaculations (C) within the  
48 lesion group, which was significantly greater than in sham group. The mean number of  
49 intromission per ejaculation (D) significantly decreased within both the sham and lesion  
50 group. The mean ejaculation latency (min.; E) decreased within the lesion group,  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

approaching significance, which was significantly less than the sham group. \* =  $p < 0.05$ ,  
error bars = SEM.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## **Acknowledgements**

The authors would like to acknowledge the expert technical contributions of Hila Eichenbaum and Vincent Laufer to this work. This work was supported by National Institutes of Health, grant number DA16272 (to A.Z.M) and National Science Foundation grant number IBN9876754 (to the Center for Behavioral Neuroscience).

**Figure 1**  
[Click here to download high resolution image](#)






Figure 2  
[Click here to download high resolution image](#)




Figure 3  
[Click here to download high resolution image](#)

