

Georgia State University

ScholarWorks @ Georgia State University

---

Nutrition Theses

Department of Nutrition

---

Fall 12-15-2010

## The Effectiveness of a Short Food Frequency Questionnaire in Determining the Adequacy of Vitamin D Intake in Children

Caitlin Russell  
*Georgia State University*

Follow this and additional works at: [https://scholarworks.gsu.edu/nutrition\\_theses](https://scholarworks.gsu.edu/nutrition_theses)



Part of the [Nutrition Commons](#)

---

### Recommended Citation


Russell, Caitlin, "The Effectiveness of a Short Food Frequency Questionnaire in Determining the Adequacy of Vitamin D Intake in Children." Thesis, Georgia State University, 2010.


doi: <https://doi.org/10.57709/1717734>

This Thesis is brought to you for free and open access by the Department of Nutrition at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Nutrition Theses by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

ACCEPTANCE

This thesis, THE EFFECTIVENESS OF A SHORT FOOD FREQUENCY QUESTIONNAIRE IN DETERMINING THE ADEQUACY OF VITAMIN D INTAKE IN CHILDREN, by Caitlin S. Russell was prepared under the direction of the Master's Thesis Advisory Committee. It is accepted by the committee members in partial fulfillment of the requirements for the degree Master of Science in the College of Health and Human Sciences, Georgia State University. The Master's Thesis Advisory Committee, as representatives of the faculty, certify that this thesis has met all standards of excellence and scholarship as determined by the faculty.

  
\_\_\_\_\_  
Anita M. Nucci, PhD, RD, LD  
Committee Chair

  
\_\_\_\_\_  
Vijay Ganji, PhD, RD  
Committee Member

  
\_\_\_\_\_  
Barbara Hopkins, MMSc, RD, LD  
Committee Member

December 13, 2010  
Date

## AUTHOR'S STATEMENT

In presenting this thesis as a partial fulfillment of the requirements for the advanced degree from Georgia State University, I agree that the library of Georgia State University shall make it available for inspection and circulation in accordance with its regulations governing materials of this type. I agree that permission to quote, to copy from, or to publish this thesis may be granted by the professor under whose direction it was written, by the College of Health and Human Sciences director of graduate studies and research, or by me. Such quoting, copying, or publishing must be solely for scholarly purposes and will not involve potential financial gain. It is understood that any copying from or publication of this thesis which involves potential financial gain will not be allowed without my written permission.

*Caitlin S. Russell*

Signature of Author

## NOTICE TO BORROWERS

All theses deposited in the Georgia State University library must be used in accordance with the stipulations prescribed by the author in the preceding statement. The author of this thesis is:

Caitlin S. Russell  
3978 Land O' Lakes Drive  
Atlanta, GA 30342

The director of this thesis is:

Anita M. Nucci, PhD, RD, LD  
Assistant Professor and Graduate Program Director  
Division of Nutrition, School of Health Professions  
College of Health and Human Sciences  
Georgia State University  
Atlanta, Georgia 30302

## VITA

Caitlin S. Russell

ADDRESS: 3978 Land O' Lakes Drive  
Atlanta, GA 30342

EDUCATION: M.S. 2010 Georgia State University  
Health Sciences  
B.A. 1994 Sweet Briar College  
French

### PROFESSIONAL EXPERIENCE:

- Director, Tradeshows 2001-2004  
AMC, Inc., Atlanta, GA
- Web & Graphic Designer 2000-2001  
ExpoExchange, LLC, Decatur, GA
- Executive Director, International Development 1997-2000  
AMC, Inc., Atlanta, GA
- Assistant English Teacher 1994-1996  
Japan Exchange & Teaching Program, Kimitsu, Japan

### PROFESSIONAL SOCIETIES AND ORGANIZATIONS:

- Treasurer, Greater Atlanta Dietetic Association 2010-present
- American Dietetic Association 2006-present
- Georgia Dietetic Association 2006-present
- Greater Atlanta Dietetic Association 2006-present

### AWARDS AND PUBLICATIONS:

- American Dietetic Association's Outstanding Dietetics Student Award 2010
- Georgia State University's Outstanding Didactic Program in  
Dietetics Award – Graduate Student 2010
- Georgia Dietetic Association's Outstanding Student – Didactic Program 2010
- Dean's List – Georgia Perimeter College Dunwoody Center 2004
- Dean's List – Georgia Perimeter College Clarkston Campus 2004
- The Georgia State University Signal, Vol. 74, Issue 27; 2007  
*Fast, cheap and healthy recipes*

## ABSTRACT

### THE EFFECTIVENESS OF A SHORT FOOD FREQUENCY QUESTIONNAIRE IN DETERMINING THE ADEQUACY OF VITAMIN D INTAKE IN CHILDREN

by  
Caitlin S. Russell

**Background:** Studies have consistently found a high prevalence of vitamin D deficiency in adolescents. Few validated dietary intake assessment tools for vitamin D exist for adolescents.

**Objective:** The aim of this study was to determine if a short food frequency questionnaire (SFFQ) can be used to effectively assess vitamin D intake in adolescents compared to a previously validated long food frequency questionnaire (LFFQ).

**Participants/setting:** 140 healthy 6-12 year old (male  $n=81$ ) Caucasian and African American ( $n=94$ ) children from Pittsburgh, Pennsylvania completed a SFFQ and LFFQ at two time points 6 months apart.

**Main outcome measures:** Reliability and validity of a SFFQ by comparison with a previously validated LFFQ for children and adolescents.

**Statistical analysis:** Reliability, validity, sensitivity, specificity, positive, and negative predictive values were assessed using Pearson correlation coefficients.

**Results:** Mean vitamin D intake from the SFFQ (range, 434 to 485 IU) was higher than the LFFQ (range, 320 to 378 IU). Overall association between the SFFQ and the LFFQ for vitamin D intake was modest ( $r=0.36$ ,  $P<0.001$ ). When stratified by race, the overall degree of association was weak for African Americans ( $r=0.26$ ,  $P=0.001$ ) and moderate for Caucasians ( $r=0.57$ ,  $P<0.001$ ). Overall reliability testing results were modest and

significant for the LFFQ ( $r=0.28$ ,  $P=0.002$ ) and SFFQ ( $r=0.33$ ,  $P<0.001$ ). Association between mean vitamin D intake from LFFQs and SFFQs was used to determine validity. The association for validity was found to be modest ( $r=0.51$ ,  $P<0.001$ ). Sensitivity, specificity, positive predictive value, and negative predictive value for the SFFQ were 90%, 64%, 0.78, and 0.58, respectively.

**Conclusion:** The SFFQ was found to be modestly valid and reliable in an early adolescent population. Associations between African Americans were not as strong as Caucasians which may be due to errors in reporting dietary consumption related to higher body weight.

THE EFFECTIVENESS OF A SHORT FOOD FREQUENCY QUESTIONNAIRE IN  
DETERMINING THE ADEQUACY OF VITAMIN D INTAKE IN CHILDREN

by  
Caitlin S. Russell

A Thesis

Presented in Partial Fulfillment of Requirements for the Degree of

Master of Science in Health Sciences

The College of Health and Human Sciences

Division of Nutrition

Georgia State University

Atlanta, Georgia  
2010

## ACKNOWLEDGMENTS

I owe my deepest gratitude to Dr. Anita Nucci for her patience, constant support and advisement throughout this process. I am honored to have received her invaluable insight into research publishing. I would also like to thank Dr. Vijay Ganji and Barbara Hopkins for their expert assistance in writing this thesis. Also, thank you to Barbara for her indispensable guidance throughout my graduate studies. I am grateful for Dr. Kumaravel Rajakumar and his staff for beginning the initial research and for their support while writing this manuscript. Finally, I am indebted to my husband Scott Russell and my daughters Eva and Julia for their encouragement and unwavering support during these past years. Without them and the rest of my loving family and friends, I literally could not have accomplished this.

## TABLE OF CONTENTS

List of Tables .....	iv
Abbreviations .....	v
Chapter	
I. INTRODUCTION .....	1
II. LITERATURE REVIEW .....	4
Vitamin D Requirements in Humans .....	4
Vitamin D Status in Children and Adolescents .....	5
Dietary Assessment Methods in Children.....	7
24-Hour Dietary Recall.....	9
Food Record.....	11
Food Frequency Questionnaire .....	13
Table 1. Characteristics of studies that focused on vitamin D intake among adolescent populations .....	16
Vitamin D Intake in Children and Adolescents .....	17
References.....	19
III. MANUSCRIPT IN STYLE OF JOURNAL.....	27
Author's Page.....	28
Abstract .....	30
Introduction.....	32
Methods.....	35
Results.....	39
Discussion .....	41
Conclusion .....	50
Tables .....	51
References.....	57

## LIST OF TABLES

Table	Page
1. Mean age and median weight, height and body mass index measures by visit and race .....	51
2. Reported intake of milk, dairy products, and vitamin D-fortified orange juice per day from the short food frequency questionnaire.....	52
3. Vitamin D intake for the long (LFFQ) and short food frequency questionnaire (SFFQ) by visit and race .....	53
4. Association between the long (LFFQ) and short food frequency questionnaires (SFFQ) for vitamin D intake by visit and race.....	54
5. Reliability test for vitamin D intake between two long food frequency questionnaires (LFFQs) administered 6 months apart and two short food frequency questionnaires (SFFQs) administered 6 months apart.....	55
6. Probability statistics results for the short food frequency questionnaire (SFFQ).....	56

## ABBREVIATIONS

1,25(OH) <sub>2</sub> D	1,25-dihydroxyvitamin D
25(OH)D	25-hydroxyvitamin D
AAP	American Academy of Pediatrics
AI	Adequate Intake
BMI	Body Mass Index
cm	centimeter
CSFII	Continuing Survey of Food Intakes by Individuals
DRI	Dietary Reference Intake
EAR	Estimated Average Requirement
FFQ	Food Frequency Questionnaire
GA	Georgia
IOM	Institute of Medicine
IU	International Units
kg	kilogram
L	liter
LFFQ	Long Food Frequency Questionnaire
µg	microgram
mL	milliliter
ng	nanogram
NHANES	National Health and Nutrition Examination Survey
nmoL	nanomole

NPV	Negative Predictive Value
oz	ounce
PA	Pennsylvania
PPV	Positive Predictive Value
RDA	Recommended Dietary Allowance
SBNM	School-Based Nutrition Monitoring
SFFQ	Short Food Frequency Questionnaire
UPMC	University of Pittsburgh Medical Center
U.S.	United States of America
USDA	United States Department of Agriculture
UVB	Ultra-violet B

## CHAPTER I

### THE EFFECTIVENESS OF A SHORT FOOD FREQUENCY QUESTIONNAIRE IN DETERMINING THE ADEQUACY OF VITAMIN D INTAKE IN CHILDREN

#### Introduction

Several studies have reported the excessive prevalence of vitamin D insufficiency among children (1-9). The primary role of vitamin D in humans is to maintain serum calcium and phosphorus concentrations at levels that support proper cell function and bone mineralization (10,11). In children, deficiency of vitamin D is associated with rickets, when growing bones fail to mineralize. When a subject is deficient in vitamin D, calcium absorption is only 10 – 15% of what is normally absorbed, resulting in decreased bone mineralization and secondary hyperparathyroidism (12). Maintaining adequate vitamin D and calcium status during childhood may reduce the risk of developing chronic diseases such as osteoporosis, multiple sclerosis, hypertension, osteomalacia, and certain cancers, as well as reduce the risk of bone fracture, falls, depression, certain autoimmune diseases, muscle weakness, and cardiovascular disease later in life (11,13-15). Immune function and inflammation reduction are also regulated by vitamin D (16). Several factors determine vitamin D status and these include dietary intake of vitamin D, skin pigmentation, season, and latitude of residence (14,16). Because vitamin D is found in very few foods, fortified food and beverages provide the U.S. population with most of its dietary vitamin D. Fish liver oil, fatty fish, and egg yolks are natural sources of vitamin D (15,16). Researchers have measured the prevalence of vitamin D deficiency in healthy adolescents using serum 25-hydroxyvitamin D (25(OH)D) as a marker and dietary

assessment methods (2-4). These studies have consistently found a high prevalence of vitamin D deficiency or insufficiency in adolescents. Subjects with the lowest serum 25(OH)D tended to be African American, obese, and those who did not consume vitamin D-rich foods (2).

There is a range of dietary assessment methods that help determine individual and population nutritional intakes and habits. These methods play an essential role in establishing when a segment of the population is at risk for nutritional problems. There are limited data available on validated dietary assessment tools for adolescents (17-20). Previously validated tools include a food frequency questionnaire (FFQ) that provided reasonable estimates on vitamin D intake in children when compared to 3-day diaries (21), a calcium FFQ that was moderately associated with estimates from 24-hour dietary recalls in middle-school-aged children (22), a FFQ that showed good test-retest reliability for assessing food intake in schoolchildren when compared with observed food intake (23), and a FFQ that suitably ranked food and nutrient intake of adolescents when compared with food records (24). Fumagalli and colleagues (25) measured dietary intake in 5-10 year olds comparing 3-day dietary records with a FFQ that had previously been validated for use in adults. They found that the FFQ appears to overestimate usual energy and nutrient intakes in children and that portion sizes need to be adjusted before adoption of this instrument in children. When designing a valid assessment tool for this age group, consideration should be given to the populations' age, reading level, ability to understand questions, ability to understand conceptual thinking, and whether an adult caretaker will assist in completing the assessment (26).

At the Primary Care Center of the Children's Hospital of Pittsburgh of the University of Pittsburgh Medical Center (UPMC), Rajakumar and colleagues conducted a Vitamin D and Sunlight study between June 2006 and March 2008. This study was designed to determine the seasonal variation of vitamin D insufficiency in healthy 6-12 year old pre- and early adolescent African American and Caucasian children residing in Pittsburgh, PA. Included in this initial study was an evaluation of nutrition intake using both a long food frequency questionnaire (LFFQ) and a short food frequency questionnaire (SFFQ). For the current study, a secondary analysis was performed on the data generated from Rajakumar's initial investigation. By way of validity and reproducibility tests, the secondary analysis aimed to determine if the SFFQ could effectively assess vitamin D intake in adolescents compared to the LFFQ. It is important to use validity and reproducibility components because any interpretation of a FFQ should include the degree to which a questionnaire can truly measure dietary intake (17). We hypothesized that there will be no difference between adolescents' vitamin D intake calculated from a SFFQ when compared to a validated LFFQ.

## CHAPTER II

### Literature Review

#### **Vitamin D Requirements in Humans**

The primary role of vitamin D in humans is to maintain cellular calcium concentrations within the homeostatic 8-10 mg/dL range. Vitamin D, or its biologically active form 1,25-dihydroxyvitamin D (1,25(OH)<sub>2</sub>D), regulates calcium and phosphorus metabolism in the intestine and bone. Without adequate vitamin D, the small intestines only absorb about 10 – 15% of dietary calcium (15). Low serum calcium stimulates parathyroid hormone (PTH) which, along with 1,25(OH)<sub>2</sub>D, increases decalcification of bone leading to osteoporosis in adults and rickets in children (27). Having a sufficient concentration of serum vitamin D has been shown to reduce diastolic and systolic blood pressures in hypertensive patients and possibly decrease the risk of developing type 1 diabetes, cardiovascular disease, osteoporosis, and certain cancers such as prostate, breast, colon, and esophagus (10,15).

Vitamin D can be obtained through the diet and from supplements. Vitamin D is synthesized in the epidermis of the skin after its exposure to ultraviolet-B (UVB) radiation from sunlight, turning 7-dehydrocholesterol (provitamin D<sub>3</sub>) into previtamin D<sub>3</sub>, which later becomes thermodynamically stable vitamin D<sub>3</sub>. This stable form of vitamin D is called cholecalciferol and later becomes 25(OH)D (calcidiol) (15). However, when skin is protected from UVB radiation by melanin pigment, sunscreen, and clothing, cutaneous vitamin D production is greatly diminished (14). Living at a high latitude reduces the amount of vitamin D that can be synthesized in the skin since fewer solar

UVB photons reach the earth than for those living at a lower latitude. Natural sources of vitamin D include cod liver oil and oily fish such as salmon and sardines. In the U.S., milk, infant formulas, some orange juices, cereals, and breads are fortified with vitamin D. Obtaining accurate estimates of average dietary intakes of vitamin D can be difficult because of the variability of vitamin D content of fortified foods (11). Vitamin D supplements are available without a prescription and many supplements generally provide 400 IU (10 µg) of vitamin D.

Whether vitamin D is made in the skin or ingested in the diet, it enters circulation and is transported to the liver by vitamin D-binding protein. In the liver, vitamin D undergoes hydroxylation and becomes 25(OH)D, the major circulating form of vitamin D. To become biologically active, it is then hydroxylated to 1,25(OH)<sub>2</sub>D in the kidneys and other tissues.

According to the Institute of Medicine, the Dietary Reference Intake (DRI) for vitamin D is 600 IU/day (15 µg) for 1-70 year olds (28). The Estimated Average Requirement (EAR) for vitamin D is 400 IU/day (10 µg) for 1-71+ year olds. The American Academy of Pediatrics (AAP) recommends 400 IU/day of vitamin D intake for all infants, children, and adolescents (13). Vitamin D status in humans is measured using 25(OH)D (11,15).

### **Vitamin D Status in Children and Adolescents**

Vitamin D deficiency is defined by most experts as 25(OH)D  $\leq$  20 ng/mL (5,6,14,29-30). Numerous studies report vitamin D deficiency and insufficiency among children and adolescents (1-9). Using data from the National Health and Nutrition Examination Survey (NHANES) 2001-2004, Kumar and colleagues (2) analyzed data on

children and adolescents and found that 25(OH)D deficiency was common among 1-21 year old persons. In this nationally representative segment of the U.S. population, 61% or 50.8 million of the children had 25(OH)D concentrations between 15 and 29 ng/mL. Children with lower 25(OH)D concentration were girls, non-Hispanic blacks, obese and those who spent more time in front of a television, playing video games, or using a computer. Vitamin D deficiency was also associated with hypertension, low HDL concentrations, and elevated PTH concentrations. Suboptimal vitamin D status was also common among healthy, low-income, minority children in Atlanta, GA (9). Cole and colleagues reported that 22% of the children in their study ( $n=290$ ) had vitamin D deficiency and 74% had insufficiency. The study authors predicted that vitamin D intake would not influence vitamin D status in this study which was true. At a Boston primary care center, Gordon and colleagues (5) studied 380 pediatric patients and found that 12% were vitamin D deficient and 40% were below an accepted optimal threshold. Among the participants who had vitamin D deficiency, 32.5% showed evidence of demineralization after wrist and knee radiographic assessments. In light of growing data that support vitamin D's immunomodulatory effects, study authors stated that their findings support recommendations that vitamin D supplementation should be made available for all young children. The Atlanta and Boston studies contradict the assumption that children living at lower latitudes exhibit better vitamin D status than children living at higher latitudes.

Using  $\leq 15$  ng/mL as deficient, Gordon and colleagues (3) found that 24.1% ( $n=307$ ) of 11-18 year old children were vitamin D deficient, 42% were vitamin D insufficient ( $\leq 20$  ng/mL), and 4.6% were severely vitamin D deficient ( $\leq 8$  ng/mL). They found that African American adolescents were more likely to be vitamin D deficient than

other ethnic groups, which corresponds with lower vitamin D production among individuals with darker skin pigmentation. They also suggested additional research on health outcomes among children and adolescents after vitamin D supplementation. In order to estimate the proportion of vitamin D insufficiency among the pediatric African American population, Rajakumar and colleagues (4) studied the response among 42 preadolescent African American children who were given 400 IU of vitamin D daily for one month. They found that 49% of the subjects were vitamin D insufficient (10-20 ng/mL) at baseline. After one month of supplementation, the mean 25(OH)D concentration significantly increased in the vitamin D insufficient group, however, 18% persisted to be vitamin D insufficient. Study authors explained that the short duration of the study and minimal dosage of vitamin D supplementation might explain the persistent insufficiency at the end of the trial. They concluded that preadolescent African Americans residing in the Northeast were at risk of low vitamin D concentrations. In a southeastern U.S. prospective study, Willis and colleagues (8) found that vitamin D status declined with age for African American and Caucasian girls aged 4-8 years old. In this population, 18% of the participants had 25(OH)D concentration below 20 ng/mL. They found an inverse relationship between circulating 25(OH)D and muscle mass, indicating that an increase in fat-free soft tissue may have an increased utilization of vitamin D. This raises the question of whether more vitamin D is required during periods of growth among adolescents.

### **Dietary Assessment Methods in Children**

As children grow from birth through adolescence, they undergo enormous social, emotional, physical, and cognitive development. There are many factors that influence

this growth, including environmental factors, genetics, and nutritional status (26). To understand the dietary intake habits of adolescents, it is important to monitor their nutritional behavior using methods that quantify what is consumed. An important prerequisite for monitoring the nutritional status in children is the accurate assessment of nutrient intake using validated tools that have been shown to accurately portray consumption patterns. There are various methods used to assess children's dietary intake, ranging from food records, 24-hour dietary recalls, FFQs, and combinations of all three (18-25,31-44). Due to the challenging nature of dietary assessment tools, some of which require concerted effort to complete on behalf of the respondent, these methods are prone to reporting error, and dietary intakes may be erroneously interpreted (45). Subjects must be able to retrieve dietary information from memory and accurately estimate portion sizes. When a child is too young to complete an assessment on his own, parents and caregivers must be able to reliably estimate what the child has consumed. Parents and caregivers may be able to report intake from the home setting but may not accurately assess what is consumed at school and outside the home. It is essential to develop valid assessment tools for adolescents, which reduce response burden and achieve precise estimation of actual nutrient intake (18,20,37).

Dietary assessment methods can assess macro- and micronutrient components of the diet. Variability is lowest among nutrients that are consumed on a regular basis such as protein, fat, and carbohydrate. Micronutrient intake, such as vitamin D, is the most variable and requires many days of records to accurately capture true intake, though it is not known exactly how many days are needed (45).

### *24-Hour Dietary Recall*

A 24-hour dietary recall is a snapshot of what a person has recently eaten. It is usually a consultation, either in person or on the telephone, between a participant and trained personnel who interview the subject about everything he has eaten and drank over the past 24-hour period (18). The participant does not have to be motivated to complete any sort of dietary record when participating in a 24-hour dietary recall. A nutrient analysis program is needed in order to assess intake after a skilled nutritionist records the information. Limitations of this method include subjects not recalling all consumed foods, reporting information they feel the clinician wants to hear, and needing to collect multiple recalls in order to determine usual intake (17). As diet is known to vary from day to day, a 24-hour recall does not give an accurate portrayal of long-term eating habits for an individual, but it does provide estimates of population means for nutrients (18). Validity of 24-hour recalls depends on how accurately an individual can recall his intake in terms of both food eaten and portion sizes. Validity is also dependent on how accurately the trained personnel codes the intake recorded during the interview and how comprehensive the nutrient database is that is used to analyze the recorded food (17). Faggiano and colleagues (46) completed a validation trial in which subjects' ( $n=103$ ) actual food consumption was compared with their recollection of consumption on the following day. All food chosen for consumption was recorded and weighed before and after the meal. For the recall, researchers used photographs of each type of dish served, including seven different portion sizes for each dish. The following day, subjects were interviewed on what they consumed during the prior meal. Subjects overestimated portion sizes by  $>20\%$  for six foods and underestimated portion sizes by  $<20\%$  for four foods. Those who ate more than average tended to underreport and those who ate less

than average tended to overreport intake. This phenomenon is generally referred to as *flat slope syndrome*. In a 2009 validation study, Baxter and colleagues (47) investigated the effects of the elapsed time between a meal and the interview (retention interval) on the accuracy of 4<sup>th</sup> graders reporting dietary intake during 24-hour recalls. The children were observed eating two school meals and later interviewed to obtain 24-hour recall. Two target periods were used: prior 24 hours (immediately preceding the interview) and previous day (midnight to midnight of the day before the interview). Interviews were conducted in the morning, afternoon, and evening. Researchers found that accuracy was better for prior 24 hours than previous day recalls and the best accuracy was obtained in the afternoon and evening. After analyzing data in 2010 from the same 2009 retention-interval validation study, Baxter and colleagues (48) concluded that shortening the retention interval of dietary intake and report increases accuracy for reporting macronutrients among this age-group.

Reliability, which is also referred to as reproducibility or precision, expresses the degree to which an investigator is able to obtain the same results when a method is repeated under similar conditions (17). If a certain method is used to assess dietary intake and it does not give consistent results after repeated use under the same conditions, the method cannot be considered reliable. Using a USDA-funded School-Based Nutrition Monitoring (SBNM) secondary-level student questionnaire, Hoelscher and colleagues (49) analyzed reproducibility results for 8<sup>th</sup> grade middle school students' food choice behavior. Using test-retest study design for reproducibility, researchers administered the survey to students at two timepoints. They found that most questions on the SBNM questionnaire had acceptable reproducibility. For questions on food choices from the

previous day, agreement was 70% or greater ( $r=0.66$ ). As this questionnaire only represented one day and was not typical intake at the individual level, authors emphasized that it is appropriate to use their study findings to characterize groups, as in cross-sectional studies, but it is not appropriate for tracking individuals' food intake over time. Also using a test-retest study design for reproducibility of questionnaire items, Penkilo and colleagues (50) measured reproducibility coefficients of an elementary school-level SBNM questionnaire among 4<sup>th</sup> graders. Food and meal choice questions had relatively good reproducibility, however mean percent agreement was lower for the elementary school-level SBNM questionnaire versus the secondary school-level SBNM questionnaire mentioned previously (78.4% vs. 86.0% respectively). The authors concluded that the questionnaire could be a valuable tool for epidemiological studies when it is necessary to assess dietary intake in a large number of children in a relatively short time period.

### *Food Record*

Similar to the 24-hour dietary recall, the food record is based on actual foods and amounts consumed by an individual on one or more specific days (17). As with 24-hour recalls, the food record is used primarily as a means to assess the validity of a FFQ when the questionnaire is used as the primary collection instrument (17,21,24-26,32-34,40,42-44,51). There is considerable flexibility afforded for dietary data analysis when using a food record since this method can accommodate any level of diverse food description detail, which is also true with the 24-hour recall method. Also known as a food diary, the food record consists of a detailed listing of all food and beverages consumed by an individual for one or more days. To reduce relying on memory, intake should be recorded

by the subject at the time of consumption and the subject should be trained on the proper way to keep an accurate log (21,37). The food record is then reviewed by trained personnel in order to ensure sufficient detail of food descriptions, preparations, and amounts (17,32). An advantage of using a food record is that, if subjects are compliant with instructions, the foods and amounts eaten are recorded immediately and relying on memory is not necessary, unlike with 24-hour recalls. Keeping a food record requires a high level of motivation for the subject, which can lead to poor response rates if subjects are not inspired to keep accurate records. Taylor and colleagues (32) designed a study to determine the validity of a food frequency questionnaire compared to a 4-day food record for assessing calcium and vitamin D intake in adolescent girls with and without anorexia nervosa. Daily calcium and vitamin D intake did not differ between the food record and questionnaire. They found that there was better compliance with the questionnaire than with the food record, which reflects the general sentiment that food records are burdensome to complete. Multiple food record days are needed in order to correctly assess usual individual intake. The number of days necessary for an accurate estimation of intake depends on the daily variability of the nutrient of interest. Basiotis and colleagues (52) used food records from 29 participants who completed a 365 consecutive-day food record to determine the number of days of intake needed to estimate average nutrient intakes for individuals and groups of individuals. Food energy and 18 nutrients, except vitamin D, were analyzed. The number of days of food intake records needed to predict the usual nutrient intake of an individual varied substantially among individuals for the same nutrient and within individuals for different nutrients. Food energy required 31 days to predict usual intake (the least) and vitamin A required 433 days (the most).

For estimating mean nutrient intake for the group, 3 days were needed for food energy and 41 days were needed for vitamin A. Authors concluded that for even larger groups, fewer days of food records would be necessary. The Basiotis study, as well as a paper by Nelson and colleagues (53) that presents a comprehensive picture of within- and between-subject variation in nutrient intake, suggest that the minimum number of days of intake required for characterizing usual intake for energy and macronutrients ranges from 3-10 days. For food components, such as vitamins and minerals, with large day-to-day variability, the number of days required may range from 20 to >50 days (17). Participant motivation and logistics make this situation impractical for most researchers. In order to assess intake of energy and macronutrients, 4-5 days of food record intake are customarily selected as a reasonable compromise. For nutrients with high within-person variability that are found in few food sources such as vitamins and minerals, FFQs specifically designed to assess these micronutrients are usually selected as the method of choice.

#### *Food Frequency Questionnaire*

The 24-hour dietary recall and the food record differ in their methods of data collection from the FFQ. Whereas the 24-hour recall and the food record report specific foods and amounts consumed on a particular day, the FFQ is based on an individual's perception of typical intake over a period of time, usually from 1 month to a year (17,18). When designing a FFQ, decisions about whether to measure intake of specific nutrients or whether there is a need for a more comprehensive dietary assessment are critical. There are many factors to consider when designing a questionnaire, which include whether the primary objective is to rank individuals or to get a measure of absolute intake

of a nutrient. Also, the questionnaire must include food items that are consumed reasonably often by a majority of individuals. Developing the food list for a questionnaire takes careful consideration and can be approached by various methods: a review of published food composition tables using foods that contain large amounts of the nutrient of interest; or start with a long list of foods and reduce the list by pilot testing the questionnaire. Stepwise regression is a more sophisticated approach and has been used, among other approaches, to develop questionnaires that measured a fairly comprehensive list of nutrients that needed to fit in a relatively small amount of space (54,55). FFQs can be “short” or “long” depending on the number of questions included. In large epidemiological studies, it may not be feasible to collect multiple days of dietary intake via the 24-hour dietary recall or the food record. Therefore, the FFQ is generally the most appropriate method for measuring intake in this situation. A number of studies have used FFQs to investigate relationships between dietary intake and disease outcomes or to just assess energy, macronutrient, and micronutrient intakes in a pediatric population (19-25,32,33,36-44). In order to determine validity of their FFQ, most of these studies compared their questionnaire with either food records or dietary recalls. Only three studies (21,32,33) were found that focused on vitamin D intake among adolescent populations, and the studies are summarized in Table 1. Using 3-day diaries for reference, Marshall and colleagues (21) investigated the relative validities of the Iowa Fluoride Study questionnaire and the Block Kids’ Food Questionnaire in assessing vitamin D intakes. Children completed the Iowa Fluoride Study nutrient questionnaire ( $n=223$ ), the Block Kids’ Food Questionnaire ( $n=129$ ) and 3-day diaries during similar time periods. Nutrient correlations between intakes estimated from 3-day diaries and questionnaires

were similar for vitamin D intake. The authors concluded that a dietary assessment method that targets specific foods or nutrients could be as effective at estimating intakes as a more comprehensive assessment tool. They also stated that it is important to consider major dietary sources of the relevant nutrient and dietary habits of the population of interest when designing a targeted nutrient tool. Using a 3-day food record as a reference, Marshall and colleagues (33) designed a study to evaluate the validity of a beverage frequency questionnaire in assessing calcium and vitamin D intakes. They found that their questionnaire could provide a relative estimate of vitamin D intake. As consumers are decreasing their milk intake and increasing their soft drink and energy drink intake, they are consuming fewer beverages that contribute vital nutrients to the total diet such as calcium and vitamin D.

Table 1. Characteristics of studies that focused on vitamin D intake among adolescent populations

Author/year publication	Participants/age group	Dietary method(s)/ reference method	Findings	Limitations
Taylor <i>et al.</i> (32) (2009)	57 girls with anorexia nervosa 50 healthy girls 12-18 years	FFQ 4d food record	Greater compliance with FFQ (99%) than food record (71%). Daily vitamin D intake from food record and FFQ did not differ. Strong correlations observed for daily vitamin D intake derived from FFQ vs food record ( $r=0.78$ , $P<0.0001$ ).	Vitamin D concentrations were not measured as biomarker of intake. Socioeconomic status and parental education, which can affect dietary reporting, were not collected.
Marshall <i>et al.</i> (21) (2008)	223 children completed Iowa Fluoride Study questionnaire and 3 d food record at 9 years; 129 children completed the Block Kids' Food Questionnaire and 3 d food record at 8.3 years	Iowa Fluoride Study targeted nutrient questionnaire Block Kids' Food Questionnaire 3 d food record	Correlations with food records for vitamin D intakes reported on Iowa Fluoride Study questionnaires were similar to correlations with vitamin D reported on Block Kids' Food Questionnaire. Percentage of exact agreement was lower for vitamin D for intakes reported on Iowa Fluoride Study questionnaires compared to Block Kids' Food Questionnaires relative to food records.	Targeted nutrient questionnaires were not administered before standard 3 d food record, leading to potential bias. Parents reported data and may not know what/how much children are eating. Subjects reporting data may not remember specifics of intake. Participants primarily Caucasian from narrow geographic region – not a representative sample of the general population. Results not generalizable to wider population.
Marshall <i>et al.</i> (33) (2003)	240 children 6 months-5 years	Quantitative beverage questionnaire 3 d food record	Correlations between mean daily nutrient intakes estimated from questionnaires and food record were 0.60-0.80 for vitamin D.	Targeted beverage questionnaire was not administered before standard 3 d food record, leading to potential bias. Parents reported data and may not know what/how much children consumed. Participants primarily Caucasian from narrow geographic region – not a representative sample of the general population. Results should not be generalized to wider population.

## **Vitamin D Intake in Children and Adolescents**

The EAR for vitamin D is 400 IU/day (10 µg) for children age 1-13 years and the RDA is 600 IU/day (15 µg) (28). According to food consumption data collected in NHANES 1988-1994 (NHANES III), the 1994-1996 Continuing Survey of Food Intakes by Individuals (CSFII 1994-1996), and the 1998 Supplemental Children's Survey, the majority of Americans are not consuming the Adequate Intake (AI) of vitamin D (56). Although dairy was the largest food source of calcium and vitamin D, some avoided dairy products due to taste or lactose intolerance. For these persons, fortified juices can provide an alternative source of vitamin D. Using NHANES 2005-2006 data, Bailey and colleagues (57) estimated vitamin D intakes from food and dietary supplements. Participants completed two 24-hour recalls and a questionnaire. Among 9-13 year olds, 24% of males and 32% of females used a daily supplement containing vitamin D. Mean vitamin D intake among this age group was 8.4 µg or 336 IU/day for males and 8.0 µg or 320 IU/day for females. The prevalence of meeting the AI for vitamin D through diet was 53% and 47% among 9-13 year old males and females, respectively. When dietary supplement use was included, 66% of 9-13 year old males met the AI for vitamin D and 53% of 9-13 year old females met the AI for vitamin D. Decreased dairy consumption relates to inadequate intake of key nutrients, including vitamin D (58). Using data from CSFII 1994-1996, 1998 and NHANES 1999-2000 reports, Fulgoni and associates (59) showed that African American males across all age groups had significantly lower dairy intake than did their non-African American counterparts. Likewise, African American females age 2-18 years consumed significantly less dairy when compared with non-African Americans. In a segment of the population that is already at risk of low serum

vitamin D concentrations due to darker skin pigmentation, these results indicate a need for increased education about the role vitamin D plays in our health and the risks of not consuming enough vitamin D-fortified foods and beverages or a vitamin D supplement.

FFQs are typically validated by comparing results with food records as the reference method (18,21,24-26,32-34,36-40,43,44). Statistical tests are used to determine whether the two instruments reported similar values for nutrients. These tests assess reliability and validity, and statistical analysis includes paired sample *t* tests, Pearson's correlation coefficients, Spearman correlations, Fisher's exact test, Wilcoxon rank sum test, weighted  $\kappa$  statistics, and percentages of exact agreement.

## REFERENCES

1. Weng FL, Shults J, Leonard MB, Stallings VA, Zemel BS. Risk factors for low serum 25-hydroxyvitamin D concentrations in otherwise healthy children and adolescents. *Am J Clin Nutr.* 2007;86:150-158.
2. Kumar J, Muntner P, Kaskel FJ, Hailpern SM, Melamed ML. Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001-2004. *Pediatrics.* 2009;124:e362-370.
3. Gordon CM, DePeter KC, Feldman HA, Grace E, Emans SJ. Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med.* 2004;158:531-537.
4. Rajakumar K, Fernstrom JD, Janosky JE, Greenspan SL. Vitamin D insufficiency in preadolescent African-American children. *Clin Pediatr.* 2005;44:683-692.
5. Gordon CM, Feldman HA, Sinclair L, Williams AL, Kleinman PK, Perez-Rossello J, Cox JE. Prevalence of vitamin D deficiency among healthy infants and toddlers. *Arch Pediatr Adolesc Med.* 2008;162:505-512.
6. Dong Y, Pollock N, Stallmann-Jorgensen IS, Gutin B, Lan L, Chen TC, Keeton D, Petty K, Holick MF, Zhu H. Low 25-hydroxyvitamin D levels in adolescents: race, season, adiposity, physical activity, and fitness. *Pediatrics.* 2010;125:1104-1111.

7. Looker AC, Dawson-Hughes B, Calvo MS, Gunter EW, Sahyoun NR. Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III. *BONE*. 2002;30:771-777.
8. Willis CM, Laing EM, Hall DB, Hausman DB, Lewis RD. A prospective analysis of plasma 25-hydroxyvitamin D concentrations in white and black prepubertal females in the southeastern United States. *Am J Clin Nutr*. 2007;85:124-130.
9. Cole CR, Grant FK, Tangpricha V, Swaby-Ellis ED, Smith JL, Jacques A, Chen H, Schleicher RL, Ziegler TR. 25-hydroxyvitamin D status of healthy, low-income, minority children in Atlanta, Georgia. *Pediatrics*. 2010;125:633-639.
10. Shils ME, Shike M, Ross C, Caballero B, Cousins R. *Modern Nutrition in Health and Disease*. 10th ed. Baltimore, MD: Lippincot Williams & Wilkins; 2005.
11. Linus Pauling Institute Micronutrient Research for Optimum Health: Vitamin D. <http://lpi.oregonstate.edu/infocenter/vitamins/vitaminD/>. Updated November 7, 2008. Accessed November 22, 2009.
12. Misra M, Pacaud D, Petryk A, Collett-Solberg PF, Kappy M. Vitamin D deficiency in children and its management: review of current knowledge and recommendations. *Pediatrics*. 2008;122:398-417.
13. Wagner CL, Greer FR. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics*. 2008;122:1142-1152.
14. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357:266-281.
15. Holick MF. Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis. *Am J Clin Nutr*. 2004;79:362-371.

16. National Institutes of Health: Dietary Supplement Fact Sheet: Vitamin D.  
<http://dietary-supplements.info.nih.gov/factsheets/vitamind.asp#h3>. Updated November 13, 2009. Accessed November 22, 2009.
17. Willet W. *Nutritional Epidemiology*. 2nd ed. New York, NY: Oxford University Press; 1998.
18. Rockett HR, Berkey CS, Colditz GA. Evaluation of dietary assessment instruments in adolescents. *Curr Opin Clin Nutr Metab Care*. 2003;6:557-562.
19. Rockett HR, Wolf AM, Colditz GA. Development and reproducibility of a food frequency questionnaire to assess diets of older children and adolescents. *J Am Diet Assoc*. 1995;95:336-340.
20. Rockett HR, Berkey CS, Colditz GA. Comparison of a short food frequency questionnaire with the Youth/Adolescent Questionnaire in the Growing Up Today Study. *Int J Pediatr Obes*. 2007;2:31-39.
21. Marshall TA, Eichenberger Gilmore JM, Broffitt B, Stumbo PJ, Levy SM. Relative validity of the Iowa Fluoride Study targeted nutrient semi-quantitative questionnaire and the block kids' food questionnaire for estimating beverage, calcium, and vitamin D intakes by children. *J Am Diet Assoc*. 2008;108:465-472.
22. Harnack LJ, Lytle LA, Story M, Galuska DA, Schmitz K, Jacobs DR, Gao S. Reliability and validity of a brief questionnaire to assess calcium intake of middle-school-aged children. *J Am Diet Assoc*. 2006;106:1790-1795.
23. de Assis MAA, Kupek E, Guimaraes D, Calvo MCM, de Andrade DF, Bellisle F. Test-retest reliability and external validity of the previous day food questionnaire for 7-10-year-old school children. *Appetite*. 2008;51:187-193.

24. Araujo MC, Yokoo EM, Pereira RA. Validation and calibration of a semiquantitative food frequency questionnaire designed for adolescents. *J Am Diet Assoc.* 2010;110:1170-1177.
25. Fumagalli F, Pontes Monteiro J, Sartorelli DS, Vieira MN, de Lourdes Pires Bianchi M. Validation of a food frequency questionnaire for assessing dietary nutrients in Brazilian children 5 to 10 years of age. *Nutrition.* 2008;24:427-432.
26. Ortiz-Andrellucchi A, Henriquez-Sanchez P, Sanchez-Villegas A, Pena-Quintana L, Mendez M, Serra-Majem L. Dietary assessment methods for micronutrient intake in infants, children and adolescents: a systematic review. *Br J Nutr.* 2009;102 Suppl 1:S87-117.
27. Holick MF. Vitamin D and bone health. *J Nutr.* 1996;126:1159S-1164S.
28. Institute of Medicine of the National Academies. Dietary Reference Intakes for Calcium and Vitamin D. Washington, DC. National Academy of Sciences, November 2010.
29. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr.* 2006;84:18-28.
30. Gilbert-Diamond D, Baylin A, Mora-Plazas M, Marin C, Arsenault JE, Hughes MD, Willett WC, Villamor E. Vitamin D deficiency and anthropometric indicators of adiposity in school-age children: a prospective study. *Am J Clin Nutr.* 2010 epub ahead of print 1-6.
31. Mascarenhas MR, Zemel B, Stallings VA. Nutritional assessment in pediatrics. *Nutrition.* 1998;14:105-115.

- 32.** Taylor C, Lamparello B, Kruczek K, Anderson EJ, Hubbard J, Misra M. Validation of a food frequency questionnaire for determining calcium and vitamin D intake by adolescent girls with anorexia nervosa. *J Am Diet Assoc.* 2009;109:479-485, 485 e471-473.
- 33.** Marshall TA, Eichenberger Gilmore JM, Broffitt B, Levy SM, Stumbo PJ. Relative validation of a beverage frequency questionnaire in children ages 6 months through 5 years using 3-day food and beverage diaries. *J Am Diet Assoc.* 2003;103:714-720; discussion 720.
- 34.** Wu H, Gozdzik A, Barta JL, Wagner D, Cole DE, Vieth R, Parra EJ, Whiting SJ. The development and evaluation of a food frequency questionnaire used in assessing vitamin D intake in a sample of healthy young Canadian adults of diverse ancestry. *Nutr Res.* 2009;29:255-261.
- 35.** Bandini LG, Schoeller DA, Cyr HN, Dietz WH. Validity of reported energy intake in obese and nonobese adolescents. *Am J Clin Nutr.* 1990;52:421-425.
- 36.** Bellu R, Ortisi MT, Riva E, Banderali G, Cucco I, Giovannini M. Validity assessment of a food frequency questionnaire for school-age children in northern Italy. *Nutr Res.* 1995;15:1121-1128.
- 37.** Rockett HR, Breitenbach M, Frazier AL, Witschi J, Wolf AM, Field AE, Colditz GA. Validation of a youth/adolescent food frequency questionnaire. *Prev Med.* 1997;26:808-816.
- 38.** Cullen KW, Watson K, Zakeri I. Relative reliability and validity of the Block Kids Questionnaire among youth aged 10 to 17 years. *J Am Diet Assoc.* 2008;108:862-866.

39. Slater B, Philippi ST, Fisberg RM, Latorre MR. Validation of a semi-quantitative adolescent food frequency questionnaire applied at a public school in Sao Paulo, Brazil. *Eur J Clin Nutr.* 2003;57:629-635.
40. Watson JF, Collins CE, Sibbritt DW, Dibley MJ, Garg ML. Reproducibility and comparative validity of a food frequency questionnaire for Australian children and adolescents. *Int J Behav Nutr Phys Act.* 2009;6:62.
41. Buzzard IM, Stanton CA, Figueiredo M, Fries EA, Nicholson R, Hogan CJ, Danish SJ. Development and reproducibility of a brief food frequency questionnaire for assessing the fat, fiber, and fruit and vegetable intakes of rural adolescents. *J Am Diet Assoc.* 2001;101:1438-1446.
42. Zemel BS, Carey LB, Paulhamus DR, Stallings VA, Ittenbach RF. Quantifying Calcium Intake in School Age Children: Development and Validation of the Calcium Counts!© Food Frequency Questionnaire. *Am J Hum Biol.* 2010;22:180-186.
43. Klohe DM, Clarke KK, George GC, Milani TJ, Hanss-Nuss H, Freeland-Graves J. Relative validity and reliability of a food frequency questionnaire for a triethnic population of 1-year-old to 3-year-old children from low-income families. *J Am Diet Assoc.* 2005;105:727-734.
44. Hacker-Thompson A, Robertson TP, Sellmeyer DE. Validation of two food frequency questionnaires for dietary calcium assessment. *J Am Diet Assoc.* 2009;109:1237-1240.
45. Livingstone MB, Robson PJ, Wallace JM. Issues in dietary intake assessment of children and adolescents. *Br J Nutr.* 2004;92 Suppl 2:S213-222.

46. Faggiano F, Vineis P, Cravanzola D, Pisani P, Xompero G, Riboli E, Kaaks R. Validation of a method for the estimation of food portion size. *Epidemiology*. 1992;3:379-382.
47. Baxter SD, Hardin JW, Guinn CH, Royer JA, Mackelprang AJ, Smith AF. Fourth-grade children's dietary recall accuracy is influenced by retention interval (target period and interview time). *J Am Diet Assoc*. 2009;109:846-856.
48. Baxter SD, Guinn CH, Royer JA, Hardin JW, Smith AF. Shortening the retention interval of 24-hour dietary recalls increases fourth-grade children's accuracy for reporting energy and macronutrient intake at school meals. *J Am Diet Assoc*. 2010;110:1178-1188.
49. Hoelscher DM, Day RS, Kelder SH, Ward JL. Reproducibility and validity of the secondary level School-Based Nutrition Monitoring student questionnaire. *J Am Diet Assoc*. 2003;103:186-194.
50. Penkilo M, George GC, Hoelscher DM. Reproducibility of the School-Based Nutrition Monitoring Questionnaire among fourth-grade students in Texas. *J Nutr Educ Behav*. 2008;40:20-27.
51. Brunner E, Stallone D, Juneja M, Bingham S, Marmot M. Dietary assessment in Whitehall II: comparison of 7 d diet diary and food-frequency questionnaire and validity against biomarkers. *Br J Nutr*. 2001;86:405-414.
52. Basiotis PP, Welsh SO, Cronin FJ, Kelsay JL, Mertz W. Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *J Nutr*. 1987;117:1638-1641.

53. Nelson M, Black AE, Morris JA, Cole TJ. Between- and within-subject variation in nutrient intake from infancy to old age: estimating the number of days required to rank dietary intakes with desired precision. *Am J Clin Nutr.* 1989;50:155-167.
54. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol.* 1985;122:51-65.
55. Mark SD, Thomas DG, Decarli A. Measurement of exposure to nutrients: an approach to the selection of informative foods. *Am J Epidemiol.* 1996;143:514-521.
56. Moore C, Murphy MM, Keast DR, Holick MF. Vitamin D intake in the United States. *J Am Diet Assoc.* 2004;104:980-983.
57. Bailey RL, Dodd KW, Goldman JA, Gahche JJ, Dwyer JT, Moshfegh AJ, Sempos CT, Picciano MF. Estimation of total usual calcium and vitamin D intakes in the United States. *J Nutr.* 2010;140:817-822.
58. Marshall TA, Eichenberger Gilmore JM, Broffitt B, Stumbo PJ, Levy SM. Diet quality in young children is influenced by beverage consumption. *J Am Coll Nutr.* 2005;24:65-75.
59. Fulgoni V, 3rd, Nicholls J, Reed A, Buckley R, Kafer K, Huth P, DiRienzo D, Miller GD. Dairy consumption and related nutrient intake in African-American adults and children in the United States: continuing survey of food intakes by individuals 1994-1996, 1998, and the National Health And Nutrition Examination Survey 1999-2000. *J Am Diet Assoc.* 2007;107:256-264.

## CHAPTER III

Manuscript in style of Journal

1 **AUTHORS' PAGE**

2 The Effectiveness of a Short Food Frequency Questionnaire in Determining the

3 Adequacy of Vitamin D Intake in Children

4 Key Words: nutritional assessment; food frequency questionnaire; adolescents; vitamin D

5 Word Count – Abstract: 293

6 Word Count – Text: 4,893

7 Anita M. Nucci, PhD, RD, LD; Assistant Professor and Graduate Program Director;

8 Division of Nutrition, School of Health Professions, College of Health and Human

9 Sciences, Georgia State University; P.O. Box 3995, Atlanta, GA, 30302-3995; Phone:

10 (404) 413-1234; Fax: (404) 413-1228; [anucci@gsu.edu](mailto:anucci@gsu.edu)

11 Caitlin Sundby Russell, MS; 3978 Land O' Lakes Drive, Atlanta, GA, 30342; Phone:

12 (404) 816-3455; [cait@mindspring.com](mailto:cait@mindspring.com)

13 Barbara Hopkins, MMSc, RD, LD; Clinical Instructor and Didactic Program in Dietetics

14 Director; Division of Nutrition, School of Health Professions, College of Health and

15 Human Sciences, Georgia State University; P.O. Box 3995, Atlanta, GA, 30302-3995;

16 Phone: (404) 413-1238; Fax: (404) 413-1228; [bhopkins@gsu.edu](mailto:bhopkins@gsu.edu)

17 Kumaravel Rajakumar, MD; Associate Professor of Pediatrics; Children's Hospital of

18 Pittsburgh of UPMC; Children's Oakland Medical Building, 3420 Fifth Avenue,

19 Pittsburgh, PA, 15213; Phone: (412) 692-6000; Fax (412) 692-6660;

20 [kumaravel.rajakumar@chp.edu](mailto:kumaravel.rajakumar@chp.edu)

21 Corresponding Author: Anita M. Nucci

22 Author for Reader Requests for Reprints: Anita M. Nucci

23 **ABSTRACT**

24 **Background:** Studies have consistently found a high prevalence of vitamin D deficiency  
25 in adolescents. Few validated dietary intake assessment tools for vitamin D exist for  
26 adolescents.

27 **Objective:** The aim of this study was to determine if a short food frequency  
28 questionnaire (SFFQ) can be used to effectively assess vitamin D intake in adolescents  
29 compared to a previously validated long food frequency questionnaire (LFFQ).

30 **Participants/setting:** 140 healthy 6-12 year old (male  $n=81$ ) Caucasian and African  
31 American ( $n=94$ ) children from Pittsburgh, Pennsylvania completed a SFFQ and LFFQ at  
32 two time points 6 months apart.

33 **Main outcome measures:** Reliability and validity of a SFFQ by comparison with a  
34 previously validated LFFQ for children and adolescents.

35 **Statistical analysis:** Reliability, validity, sensitivity, specificity, positive, and negative  
36 predictive values were assessed using Pearson correlation coefficients.

37 **Results:** Mean vitamin D intake from the SFFQ (range, 434 to 485 IU) was higher than  
38 the LFFQ (range, 320 to 378 IU). Overall association between the SFFQ and the LFFQ  
39 for vitamin D intake was modest ( $r=0.36$ ,  $P<0.001$ ). When stratified by race, the overall  
40 degree of association was weak for African Americans ( $r=0.26$ ,  $P=0.001$ ) and moderate  
41 for Caucasians ( $r=0.57$ ,  $P<0.001$ ). Overall reliability testing results were modest and  
42 significant for the LFFQ ( $r=0.28$ ,  $P=0.002$ ) and SFFQ ( $r=0.33$ ,  $P<0.001$ ). Association  
43 between mean vitamin D intake from LFFQs and SFFQs was used to determine validity.  
44 The association for validity was found to be modest ( $r=0.51$ ,  $P<0.001$ ). Sensitivity,

45 specificity, positive predictive value, and negative predictive value for the SFFQ were  
46 90%, 64%, 0.78, and 0.58, respectively.

47 **Conclusion:** The SFFQ was found to be modestly valid and reliable in an early  
48 adolescent population. Associations between African Americans were not as strong as  
49 Caucasians which may be due to errors in reporting dietary consumption related to higher  
50 body weight.

51 The Effectiveness of a Short Food Frequency Questionnaire in Determining the  
52 Adequacy of Vitamin D Intake in Children

### 53 **INTRODUCTION**

54         Several studies have reported the excessive prevalence of vitamin D insufficiency  
55 among children (1-9). The primary role of vitamin D in humans is to maintain serum  
56 calcium and phosphorus concentrations at levels that support proper cell function and  
57 bone mineralization (10,11). In children, deficiency of vitamin D is associated with  
58 rickets, when growing bones fail to mineralize. When a subject is deficient in vitamin D,  
59 calcium absorption is only 10 – 15% of what is normally absorbed, resulting in decreased  
60 bone mineralization and secondary hyperparathyroidism (12). Maintaining adequate  
61 vitamin D and calcium status during childhood may reduce the risk of developing chronic  
62 diseases such as osteoporosis, multiple sclerosis, hypertension, osteomalacia, and certain  
63 cancers, as well as reduce the risk of bone fracture, falls, depression, certain autoimmune  
64 diseases, muscle weakness, and cardiovascular disease later in life (11,13-15). Immune  
65 function and inflammation reduction are also regulated by vitamin D (16). Several factors  
66 determine vitamin D status and these include dietary intake of vitamin D, skin  
67 pigmentation, season, and latitude of residence (14,16). Because vitamin D is found in  
68 very few foods, fortified food and beverages provide the U.S. population with most of its  
69 dietary vitamin D. Fish liver oil, fatty fish, and egg yolks are natural sources of vitamin D  
70 (15,16). Researchers have measured the prevalence of vitamin D deficiency in healthy  
71 adolescents using serum 25-hydroxyvitamin D (25(OH)D) as a marker and dietary  
72 assessment methods (2-4). These studies have consistently found a high prevalence of  
73 vitamin D deficiency or insufficiency in adolescents. Subjects with the lowest serum

74 25(OH)D tended to be African American, obese, and those who did not consume vitamin  
75 D-rich foods (2).

76         There is a range of dietary assessment methods that help determine individual and  
77 population nutritional intakes and habits. These methods play an essential role in  
78 establishing when a segment of the population is at risk for nutritional problems. There  
79 are limited data available on validated dietary assessment tools for adolescents (17-20).  
80 Previously validated tools include a food frequency questionnaire (FFQ) that provided  
81 reasonable estimates on vitamin D intake in children when compared to 3-day diaries  
82 (21), a calcium FFQ that was moderately associated with estimates from 24-hour dietary  
83 recalls in middle-school-aged children (22), a FFQ that showed good test-retest reliability  
84 for assessing food intake in schoolchildren when compared with observed food intake  
85 (23), and a FFQ that suitably ranked food and nutrient intake of adolescents when  
86 compared with food records (24). Fumagalli and colleagues (25) measured dietary intake  
87 in 5-10 year olds comparing 3-day dietary records with a FFQ that had previously been  
88 validated for use in adults. They found that the FFQ appears to overestimate usual energy  
89 and nutrient intakes in children and that portion sizes need to be adjusted before adoption  
90 of this instrument in children. When designing a valid assessment tool for this age group,  
91 consideration should be given to the populations' age, reading level, ability to understand  
92 questions, ability to understand conceptual thinking, and whether an adult caretaker will  
93 assist in completing the assessment (26).

94         At the Primary Care Center of the Children's Hospital of Pittsburgh of the  
95 University of Pittsburgh Medical Center (UPMC), Rajakumar and colleagues conducted a  
96 Vitamin D and Sunlight study between June 2006 and March 2008. This study was

97 designed to determine the seasonal variation of vitamin D insufficiency in healthy 6-12  
98 year old pre- and early adolescent African American and Caucasian children residing in  
99 Pittsburgh, PA. Included in this initial study was an evaluation of nutrition intake using  
100 both a long food frequency questionnaire (LFFQ) and a short food frequency  
101 questionnaire (SFFQ). For the current study, a secondary analysis was performed on the  
102 data generated from Rajakumar's initial investigation. By way of validity and  
103 reproducibility tests, the secondary analysis aimed to determine if the SFFQ could  
104 effectively assess vitamin D intake in adolescents compared to the LFFQ. It is important  
105 to use validity and reproducibility components because any interpretation of a FFQ  
106 should include the degree to which a questionnaire can truly measure dietary intake (17).  
107 We hypothesized that there will be no difference between adolescents' vitamin D intake  
108 calculated from a SFFQ when compared to a validated LFFQ.

## 109 **METHODS**

### 110 **Study Design**

111           The Vitamin D and Sunlight study was an observational study designed to  
112 determine the seasonal variation of vitamin D insufficiency, and to assess its risk factors  
113 and metabolic impact in healthy 6-12 year old pre-and early adolescent African American  
114 and Caucasian children residing in Pittsburgh, PA. A total of 140 participants were  
115 enrolled in the initial study between the summer months of June-September 2006 and the  
116 winter months of December 2007-March 2008. There was no intervention and  
117 participants were seen at 2 time points, with the initial visit being either in the summer or  
118 winter and the follow-up visit occurring 6 months later. Blood was drawn at both visits to  
119 determine serum concentrations of calcium, phosphorus, albumin, serum PTH, 25(OH)D  
120 and 1,25(OH)<sub>2</sub>D. Subjects completed a LFFQ and a SFFQ at both timepoints in order to  
121 assess sunlight exposure and dietary intake of vitamin D and calcium. The initial study  
122 was approved by the Institutional Review Board of the University of Pittsburgh. Written  
123 informed consent was obtained from parents at recruitment of the initial study and  
124 participants' assent was obtained prior to enrollment. The Institutional Review Board at  
125 Georgia State University approved the secondary analysis study.

126

### 127 **Subjects**

128           The initial study used a convenience sample of patients with a mean age of 9.1  
129 years at the time of recruitment. Participants were recruited from the Primary Care Center  
130 of the Children's Hospital of Pittsburgh of the University of Pittsburgh Medical Center  
131 (UPMC) and from the surrounding area. Inclusion criteria included healthy 6-12 year old

132 pre- and early adolescent African American and Caucasian children residing in  
133 Pittsburgh, PA. Children with hepatic or renal disease, metabolic rickets, malabsorptive  
134 disorders, cancer, or those on treatment with anticonvulsants or systemic glucocorticoids  
135 were excluded.

136

### 137 **Data Collection**

138 Study participants received a clinical exam at their initial and six month visit to  
139 gather anthropometric information including weight (kg), height (cm) and body mass  
140 index (BMI) ( $\text{kg}/\text{m}^2$ ). BMI was calculated from weight and height at baseline and follow-  
141 up. Dietary intake was recorded by participants at both time points using a SFFQ and a  
142 LFFQ. BMI was evaluated using the 2000 Centers for Disease Control and Prevention  
143 growth charts, where overweight is determined by a BMI value between the 85<sup>th</sup> and 95<sup>th</sup>  
144 percentile and obesity >95<sup>th</sup> percentile (27,28).

145

#### 146 *The SFFQ.*

147 Rajakumar and colleagues adapted a questionnaire, developed by Dr. Michael  
148 Holick (Boston University Medical Center) that assessed vitamin D intake in adults, to  
149 address vitamin D intake and sunlight exposure in pre- and early adolescents. The  
150 questionnaire included 21 questions for the patient/parent to complete. Subjects were  
151 asked if they take a multivitamin, vitamin D supplement, or cod liver oil, how many  
152 servings of milk, cheese, yogurt, and vitamin D-fortified orange juice they consume per  
153 day, and how often they consume fish and dried mushrooms per month. The remaining  
154 questions documented sun exposure and sunscreen use.

155

156 *The LFFQ.*

157           Subjects also completed a LFFQ at baseline and at the follow-up visit. The LFFQ  
158 is an eating survey designed by the Harvard Medical School, copyright ©1995 Brigham  
159 and Women’s Hospital. The LFFQ was used as the gold standard against the SFFQ. The  
160 LFFQ is a semiquantitative FFQ that consists of 7 food groups with 152 questions and  
161 requests information on dietary intake over the past year.

162

163 *Data Analyses.*

164           In order to quantify the dietary intake of vitamin D, completed SFFQs were  
165 analyzed using the Food Processor SQL (version 10.4.0, ESHA Research; Salem, OR) by  
166 the Clinical Nutrition Department at the Children’s Hospital of Pittsburgh of UPMC.  
167 LFFQ data forms were analyzed at Brigham and Women’s Hospital. Nutrient intake data  
168 comprised 17 nutrients including total energy, dietary calcium, and dietary vitamin D.

169           For the current study, variables and data distribution were analyzed for normality  
170 using the Kolmogorov-Smirnov test ( $P \leq 0.05$ ). Normality testing revealed that most  
171 anthropometric data were not normally distributed (Table 1). Mean values were used for  
172 age and vitamin D intake while median values were used for weight, height and BMI.  
173 The Wilcoxon Signed-Rank test was used to examine differences in BMI between the  
174 initial and 6 month visit. The Mann-Whitney U test was used to determine if there was a  
175 significant difference between BMI by race at each time point. A summary of SFFQ  
176 responses for multivitamin, calcium supplement, and vitamin D supplement use, as well  
177 as dairy, and vitamin D-fortified orange juice consumption was determined using

178 frequency statistics. Vitamin D intake (IU) was found to be normally distributed for both  
179 the LFFQ and SFFQ for most groups. Pearson's correlation was used to determine the  
180 correlation coefficient between the LFFQ and the SFFQ for vitamin D intake. The  
181 correlation coefficient "*r*" indicates the strength of the relationship and the value can  
182 range from -1.0 to +1.0. A correlation coefficient of near +1.0 implies perfect  
183 correlation. Correlations were interpreted using the following guidelines: 0.0–0.3 as  
184 weak, 0.3–0.6 as modest and 0.6–1.0 as good. Reliability testing was done using  
185 Pearson's correlation between the LFFQ and SFFQ completed at the initial visit and  
186 corresponding LFFQ and SFFQ completed 6 months later. Validity of the SFFQ and  
187 LFFQ was evaluated by comparing mean vitamin D intake from the two LFFQs with the  
188 two SFFQs. External validity of the SFFQ was evaluated via sensitivity, specificity,  
189 positive predictive value (PPV), and negative predictive value (NPV) equations.  
190 Sensitivity refers to the test instrument (SFFQ) and describes the proportion of  
191 participants who were correctly identified as having met the American Academy of  
192 Pediatrics' (AAP) recommendation for vitamin D (400 IU/day). Specificity also refers to  
193 the test instrument, but describes the proportion of participants who were correctly  
194 identified as not having met the AAP's recommendation for vitamin D. PPV measures  
195 the probability that a person who tests positive for meeting the AAP's recommendation  
196 truly consumed 400 IU/day. NPV measures the probability that a person who tests  
197 negative for meeting the AAP's recommendation actually consumed less vitamin D.  
198 Statistical analyses were done using IBM® SPSS® Statistics 18 (version 18, 2010, IBM  
199 Corp, Armonk, NY).

## 200 RESULTS

201 The study sample ( $n=140$ ) consisted of 67% African American ( $n=94$ ). At the 6  
202 month follow-up ( $n=122$ ), the sample consisted of 68% African American ( $n=83$ ). For  
203 the 6 month visit, 18 participants were lost due to follow-up. At both time points, 58% of  
204 the participants were boys. The majority of data is missing for one participant in the 6  
205 month follow-up group and has been excluded from the data analysis. In African  
206 Americans, BMI ranged from 13.8 to 35.3 kg/m<sup>2</sup> for the initial and 6 month visits, with  
207 46% of observations above the 85<sup>th</sup> percentile at the initial visit and 50% at the 6 month  
208 visit. For Caucasians, BMI ranged from 14.3 to 32.0 kg/m<sup>2</sup>, with 25% and 28% of  
209 observations above the 85<sup>th</sup> percentile at the initial visit and 6 month visit, respectively.  
210 At the initial visit, 24% of African Americans were obese compared with 12% of  
211 Caucasians. Six months later, 29% of African Americans were obese compared with 12%  
212 of Caucasians. A significant difference in BMI was found between the initial and 6  
213 month visits for Caucasians (16.3 vs. 17.1 kg/m<sup>2</sup>,  $P<0.01$ ) and African Americans (18.3  
214 vs. 19.1 kg/m<sup>2</sup>,  $P<0.001$ ) (Table 1). There was a significant difference between BMI by  
215 race at the initial and at the 6 month visit ( $P<0.05$ ).

216 At the initial visit, 24% ( $n=34$ ) of participants reported taking a multivitamin,  
217 while 33% ( $n=39$ ) reported multivitamin use at the follow-up visit. Only 1 participant  
218 reported taking a vitamin D supplement at the initial and 6 month visit. Reported intake  
219 of milk, dairy products, and vitamin D-fortified orange juice from the SFFQ is shown in  
220 Table 2. Median milk intake per day for participants at the initial and six month visit was  
221 two 8-oz servings of milk. Median daily cheese consumption was one 1-oz serving for  
222 the initial visit and one and one-half 1-oz servings at the follow up visit. Median daily

223 intake of other dairy products (i.e. Lactaid<sup>®</sup>, chocolate milk, yogurt) was minimal. Mean  
224 vitamin D intake stratified by visit and race is shown in Table 3. Mean values for vitamin  
225 D intake for the LFFQ were significantly lower than for the SFFQ at both timepoints  
226 ( $P<0.001$ ). There are intra-group differences between LFFQs and SFFQs at the initial and  
227 6 month follow-up visits.

228         Associations between the LFFQ and the SFFQ for vitamin D intake by race and  
229 visit timepoint are shown in Table 4. With the exception of African Americans at the 6  
230 month visit, the associations between the 2 FFQs for vitamin D intake were significant  
231 ( $P<0.01$ ). For all pairs, the degree of association between the LFFQ and the SFFQ was  
232 modest ( $r=0.36$ ). After stratifying the population by race, the degree of correlation  
233 between the LFFQ and the SFFQ was weak ( $r=0.26$ ) for African Americans and modest  
234 for Caucasians ( $r=0.57$ ).

235         Reliability test results for vitamin D intake between the two LFFQs administered  
236 6 months apart and the two SFFQs administered 6 months apart are shown in Table 5.  
237 With the exception of African Americans for the LFFQs, the reliability of the 2  
238 questionnaires is modest and is statistically significant ( $P<0.01$ ) for all groups.

239         Association or validity between the average vitamin D intake from the two LFFQs  
240 and the two SFFQs administered 6 months apart ( $n=119$ ) was modest ( $r=0.51$ ) and  
241 significant ( $P<0.001$ ). Probability statistics for predicting whether the SFFQ is able to  
242 identify participants who have or have not met the AAP's recommendation for vitamin D  
243 are shown in Table 6. The SFFQ was tested against the LFFQ for sensitivity, specificity,  
244 PPV, and NPV. In this evaluation, 90% of participants were correctly identified as having  
245 met the AAP's recommendation for vitamin D (400 IU/day). Of these participants, the

246 probability (PPV) that they actually consumed  $\geq 400$  IU/day was 0.78. This indicates that  
247 the SFFQ is very sensitive and that the probability of the result being true is high.  
248 According to the SFFQ, 64% of participants were correctly identified as not having met  
249 the AAP's recommendation; the probability that these participants actually consumed  
250  $< 400$  IU/day was 0.58. This indicates that the SFFQ is less specific than it is sensitive;  
251 however, out of those who did not consume 400 IU/day, the likelihood of this result  
252 being false is low.

253

## 254 **DISCUSSION**

255         This study was designed to evaluate the reliability and validity of a SFFQ in  
256 determining vitamin D intake among adolescents using a LFFQ as a reference. There are  
257 limited data available on adolescent behavior in regards to dietary consumption habits  
258 (18,19). The adolescents' level of independence and their caretakers' unawareness of the  
259 foods they consume further complicate assessing adolescent dietary consumption (29).  
260 Developing a valid tool that accurately measures vitamin D consumption in children is  
261 important in order to assess factors that optimize adequate vitamin D status. Shortening a  
262 long FFQ to a targeted short FFQ would provide a quick and accurate estimate of the  
263 nutrient of interest. Constructing and evaluating a less burdensome tool (the SFFQ) that  
264 specifically targets a certain nutrient (vitamin D) is important in order to assess the  
265 vitamin D status in a vulnerable segment of the population.

266         Rajakumar's initial study is significant because there are a limited number of  
267 dietary assessment instruments that are specifically designed for assessing vitamin D  
268 intake in adolescents (18,19,30). Accurately assessing the dietary intake of the adolescent

269 population is essential in order to monitor their nutritional status. Valid assessment is also  
270 crucial for determining links between diet and health (31). It is important to evaluate the  
271 effectiveness of Rajakumar's SFFQ because this instrument could be used in future large  
272 epidemiological studies to estimate the vitamin D intake of an adolescent population.  
273 Estimating vitamin D consumption in adolescents is important because of the significant  
274 role this nutrient plays in establishing peak bone mass, as well as potentially preventing  
275 or delaying adult-onset diseases (10,13). Since studies have shown that vitamin D  
276 deficient status in US children is prevalent, it is important to have an instrument that will  
277 quickly assess an adolescent's vitamin D intake. Subsequent intervention can take place  
278 if subjects are found to be deficient.

279         For the total population in our study, the mean vitamin D intake, assessed by the  
280 SFFQ, was 142 IU and 90 IU higher than the LFFQ at the initial visit and 6 month  
281 follow-up visit, respectively. This is not surprising given that our SFFQ's primary focus  
282 was on vitamin-D-rich foods, whereas the LFFQ assessed all foods eaten over a one year  
283 period, not concentrating specifically on one nutrient. Despite the higher reported vitamin  
284 D intake by the SFFQ, associations between the short and long FFQs ranged from modest  
285 for the entire population ( $r=0.36$ ) and weak for African Americans ( $r=0.26$ ) to modest for  
286 Caucasians ( $r=0.57$ ). Using a 3-day diary for validation purposes, Marshall and  
287 colleagues (21) found that vitamin D intake estimates were lower on targeted nutrient  
288 questionnaires compared to food diaries. Our findings differ perhaps due to the open-  
289 ended nature of food diaries, which allow a more comprehensive assessment of nutrient  
290 intake and thus accumulate more nutrient-specific data. FFQs assess specific foods or  
291 nutrients and may leave out a food that is routinely eaten by a participant.

292           At the 6 month visit, we found a weak relation ( $r=0.19$ ) between the LFFQ and  
293 SFFQ for African Americans. Kaaks and colleagues (32) report that, when compared to  
294 normal weight subjects, obese subjects usually underreported food intake regardless of  
295 dietary assessment technique used. Similarly, Bandini and colleagues (33) found that  
296 obese adolescents underreported food intake significantly more than nonobese  
297 adolescents. This could explain the weak relation seen in the African American segment  
298 of our study.

299           Reliability was assessed by comparing vitamin D intake data from the same  
300 questionnaire administered at two separate time points. Associations were weak for the  
301 entire population for the LFFQs administered six months apart ( $r=0.28$ ) and modest for  
302 the entire population for the SFFQs administered 6 months apart ( $r=0.33$ ). Validity was  
303 assessed by comparing vitamin D intake data collected using different methods (LFFQ  
304 and SFFQ) at two time points. Using unadjusted values of vitamin D intake, validity  
305 testing revealed modest association ( $r=0.51$ ). Based on sensitivity and specificity, it is  
306 deduced that the SFFQ is more accurate in measuring the proportion of people who met  
307 the AAP's recommendation for vitamin D than measuring the proportion of people who  
308 did not meet the AAP's recommendation.

309           The United States Department of Agriculture's (USDA) MyPyramid recommends  
310 2-3 cups from the milk group per day for children (34). In general, 1 cup in the milk  
311 group is defined as 1 cup of milk or yogurt, 1.5-oz of natural cheese, or 2-oz of processed  
312 cheese. The reported median intake our sample consumed from the milk group per day  
313 was 3.05 cups at the initial visit and 3.65 cups at the 6 month follow up visit, which  
314 slightly exceeds the USDA guideline.

315 This is the first study to assess vitamin D-only intake data from a SFFQ designed  
316 specifically to measure vitamin D consumption in adolescents. Other studies have  
317 assessed total macronutrient and select micronutrient intakes or types of foods eaten in  
318 children (23-25,35); however, only 4 studies included relations between mean daily  
319 vitamin D intake estimated from questionnaires and food records in children and young  
320 adults (21,36-38). Taylor and colleagues (37) performed a study to determine the validity  
321 of a FFQ for assessing calcium and vitamin D intake in 12-18 year old anorexic and  
322 healthy girls. Researchers validated the FFQ with 4-day food records. They found that  
323 subjects demonstrated greater compliance with the FFQ than with the food record. They  
324 also observed a strong association ( $r=0.78$ ) between daily vitamin D intake derived from  
325 the FFQ and the food record. The researchers concluded that the FFQ was useful in  
326 maximizing the information regarding calcium and vitamin D intake in a segment of the  
327 population that is vulnerable to nutrient deficiencies. Our SFFQ had similar or better  
328 validity than reported in other studies of one or more micronutrient-focused FFQs  
329 (22,38,39). Hacker-Thompson and colleagues (39) compared 2 calcium-focused  
330 questionnaires to 3-day diet records. They found modest association between the  
331 questionnaires and the diet records ( $r=0.37$ ). Harnack and colleagues (22) found good  
332 reliability ( $r=0.74$ ) of a calcium FFQ administered twice, with 1 week between  
333 administrations. Validity showed a modest association between the calcium FFQ and the  
334 dietary recall ( $r=0.43$ ). Wu and colleagues (38) evaluated the validity of a FFQ for  
335 assessing vitamin D and used a 7-day food diary for reference. Validity of the FFQ was  
336 good ( $r=0.60$ ).

337           Using a 3-day diary as a reference, Marshall and colleagues (21) investigated the  
338 relative validities of the Iowa Fluoride Study questionnaire and the Block Kids' Food  
339 Questionnaire in assessing vitamin D intakes. Children completed the Iowa Fluoride  
340 Study nutrient questionnaire ( $n=223$ ), the Block Kids' Food Questionnaire ( $n=129$ ), and  
341 3-day diaries during similar time periods. Associations between intakes estimated from 3-  
342 day diaries and the 2 questionnaires were modest for vitamin D intake and virtually  
343 identical to our validation findings ( $r=0.51$ ). Study authors concluded that a dietary  
344 assessment method that targets specific foods or nutrients can be as effective at  
345 estimating intakes as a more comprehensive assessment tool. They also stated that it was  
346 important to consider major dietary sources of the relevant nutrient and dietary habits of  
347 the population of interest when designing a targeted nutrient tool. Using a 3-day food  
348 record as reference, Marshall and colleagues (36) designed a study to evaluate the  
349 validity of a beverage frequency questionnaire in assessing calcium and vitamin D  
350 intakes. They found that their questionnaire could provide a relative estimate of vitamin  
351 D intake. The validity of our vitamin D-focused SFFQ compared favorably with the  
352 results of Marshall's beverage questionnaire. Researchers stated that as beverage intake  
353 patterns change, with decreased milk intake and increased soft drink and energy drink  
354 intake, this leaves fewer beverages contributing vital nutrients such as calcium and  
355 vitamin D to the total diet. A study by Araujo and colleagues (24) estimated the validity  
356 of dietary intake data from a FFQ designed specifically for adolescents living in Brazil  
357 and demonstrated that their FFQ was a suitable tool for ranking adolescent's energy and  
358 nutrient intake. Study authors stated that aiding tools such as food models and  
359 photographs should be used to reduce bias when reporting food intake portions.

360 Estimating vitamin D intake in children is important because of the role of  
361 vitamin D in various important biological functions that include accruing peak bone  
362 mass. Increased melanin causes decreased cutaneous production of previtamin D<sub>3</sub>, putting  
363 African Americans at higher risk of vitamin D deficiency (40). For those who live  $\geq 40$   
364 degrees north all year or  $\leq 40$  degrees from November to early March, there is  
365 insufficient ultraviolet B (UVB) radiation from the sun for vitamin D synthesis (11).  
366 Experts disagree about what constitutes vitamin D deficiency (1,3,9,10,13,14,16,41-42).  
367 Proposed cutoff values for deficient 25(OH)D range from 10 ng/mL to 30 ng/mL.

368 In November 2010, the Institute of Medicine (IOM) revised the Dietary Reference  
369 Intakes (DRIs) for vitamin D. The new recommendations include an EAR of 400 IU/day  
370 for all age groups. The Recommended Dietary Allowance (RDA) is 600 IUs for persons  
371 age 1-70 years (43). In a 2008 publication, the American Academy of Pediatrics (AAP)  
372 revised their recommendation that all infants, children, and adolescents have a minimum  
373 daily intake of 400 IU of vitamin D, up from 200 IU/day (13). In our total study  
374 population using the SFFQ, 54% reportedly met 400 IU/day and 30% met 600 IU/day at  
375 the initial visit. At the follow-up visit, these percentages reduced to 46% and 23%,  
376 respectively.

377

### 378 **Vitamin D Insufficiency in Children**

379 Among the scientific community, there is mounting consensus that vitamin D  
380 insufficiency is more widespread than originally thought (1-9). Research has shown that a  
381 large percentage of adolescents living in the northern hemisphere are frequently vitamin  
382 D deficient, particularly African Americans. Otherwise healthy children may not show

383 clinical signs of vitamin D deficiency, but when serum 25(OH)D concentration is  
384 measured, vitamin D deficiency is present in many US adolescents. Using data from the  
385 National Health and Nutrition Examination Survey (NHANES) 2001-2004, Kumar and  
386 colleagues (1) analyzed data on children and adolescents and found that vitamin D  
387 deficiency was common among 1-21 year old persons and was associated with adverse  
388 cardiovascular risks. In this nationally representative sample of the U.S. population, 61%  
389 (50.8 million) of the children had vitamin D concentrations between 15 and 29 ng/mL,  
390 considered vitamin D deficient in this study. Only 4% of the children had taken 400 IU of  
391 vitamin D/day for 30 days. In our study, only 1 child out of 140 reported taking a vitamin  
392 D supplement on a regular basis. In Kumar's study, children with lower vitamin D  
393 concentrations were girls, non-Hispanic blacks, obese, and those who spent more time in  
394 front of a television, playing video games, or using a computer. Deficiency was also  
395 associated with hypertension, low HDL concentration, and elevated PTH. Suboptimal  
396 vitamin D status was also common among healthy, low-income, minority children in  
397 Atlanta, GA (8). Cole and colleagues reported that 22% of the children in their study  
398 ( $n=290$ ) had vitamin D deficiency ( $\leq 20$  ng/mL) and 74% had insufficiency ( $\leq 30$  ng/mL).  
399 Fortified milk provided most dietary vitamin D (62%), comparing favorably with our  
400 study where milk was the most consumed source of dietary vitamin D. Cole and  
401 colleagues concluded that vitamin D intake did not influence vitamin D status. At a  
402 Boston primary care center, Gordon and colleagues (4) studied 380 pediatric patients and  
403 found that 12% were vitamin D deficient ( $\leq 20$  ng/mL) and 40% were below an accepted  
404 optimal threshold ( $\leq 30$  ng/mL). Multivitamin use was reported in 25% of subjects and  
405 69% of subjects reported occasional consumption of fortified milk. Among the

406 participants who had vitamin D deficiency, 32.5% showed evidence of demineralization  
407 after wrist and knee radiographic assessments. In light of growing data that support  
408 vitamin D's immunomodulatory effects, study authors stated that their findings support  
409 recommendations that vitamin D supplementation should be made available for all young  
410 children. The Atlanta and Boston studies contradict the assumption that children living at  
411 lower latitudes exhibit better vitamin D status than children living at higher latitudes.

412

### 413 **Food Frequency Questionnaires**

414 FFQs that assess nutrient intake in adolescents are uncommon. Although dietary  
415 recalls and food records allow for a comprehensive evaluation of dietary habits, they  
416 require a substantial commitment from the subjects which can lead to noncompliance and  
417 increased dropout rates (21). Therefore, the FFQ has become the primary method for  
418 assessing intake in large, population-based studies. The FFQ can also be specialized to  
419 identify specific nutrient intake (44). The FFQ estimates food intake over a period of time  
420 and is usually less expensive and takes less time to complete than a 24-hour recall and a  
421 food record (18). Nutrition assessment instruments like the SFFQ in our study are easy to  
422 administer, low-cost, simple to complete, reproducible, and accurate (19). The use of  
423 FFQs can be a consistent and precise method for describing habitual dietary intake.  
424 Before a FFQ is used in an epidemiological study, it should be validated to see if it  
425 actually measures the aspect of the diet it was intended to assess (17). Although Klohe  
426 and colleagues did not exclusively determine vitamin D status, they developed and  
427 validated a FFQ for low-income children, and found that the FFQ yielded excellent  
428 reliability and acceptable validity when compared to a 3-day diet record (45). This

429 demonstrates that FFQs can be reliable tools for assessing dietary consumption patterns  
430 in youth.

431         Due to variations in adolescents' literacy levels and their limited knowledge of  
432 foods, developing an assessment tool that accurately reports nutrient intake is challenging  
433 (46). We were unable to adjust the analysis for energy intake because energy intake was  
434 not mentioned in the SFFQ. However, no association was found between calories and  
435 vitamin D for the LFFQ; we felt using the unadjusted values for the SFFQ was justified.  
436 Another limitation was the relatively small sample size of our study, which may have  
437 affected the statistical power for vitamin D intake assessment. In addition to the SFFQ  
438 and the LFFQ, a 3-day food record was completed by fewer than half of the participants.  
439 The food record was to be used to further validate the SFFQ. However, the data from  
440 many of the food records were incomplete, and did not represent an accurate description  
441 of macro- and micronutrient consumption. Therefore, it was determined not to use the  
442 food record data in this secondary analysis.

443         Future studies should involve continued evaluation of this micronutrient  
444 assessment instrument, especially among populations vulnerable to vitamin D deficiency.  
445 Using biological markers such as 25(OH)D could improve the reliability and validity of  
446 this instrument. As FFQs can provide valuable feedback, which is useful for evaluating  
447 micronutrient policy initiatives, the SFFQ used in this study could be used in large-scale  
448 surveys to monitor vitamin D intake in segments of the population at risk for deficiency.  
449 Additional studies are needed to document the effects of screening otherwise healthy  
450 children for and treatment of deficient vitamin D status.

451 **Conclusion**

452           The SFFQ is a reasonably valid and reliable tool that can be used to assess  
453 vitamin D intake in adolescents. The SFFQ would serve well as a tool for identifying  
454 children with low vitamin D intake who may benefit from a vitamin D nutrition  
455 intervention designed for increasing vitamin D consumption.

456 **Tables**

457 Table 1. Mean age and median weight, height and body mass index measures by visit and  
 458 race

Initial Visit	Total (n=140)	African American (n=94)	Caucasian (n=46)
Age (y) <sup>a</sup>	9.1 ± 1.7	8.9 ± 1.7	9.5 ± 1.7
Weight (kg) <sup>b</sup>	32.1 (26.8-39.4)	34.5 (27.5-39.1)	31.2 (24.9-41.6)
Height (cm) <sup>b</sup>	133.5 (126.8-140.8)	131.9 (127.0-139.8)	135.4 (126.5-145.6)
Body mass index (kg/m <sup>2</sup> ) <sup>b</sup>	17.8 (16.2-20.4)	18.3 (16.6-21.0)*‡	16.3 (15.4-19.2)**
6 Month Visit	(n=122)	(n=83)	(n=39)
Age (y) <sup>a</sup>	9.6 ± 1.7	9.5 ± 1.7	10.0 ± 1.8
Weight (kg) <sup>b</sup>	34.1 (28.8-42.8)	35.0 (29.0-42.7)	33.4 (27.8-45.7)
Height (cm) <sup>b</sup>	136.3 (129.5-143.2)	135.5 (129.5-142.2)	137.9 (128.2-145.6)
Body mass index (kg/m <sup>2</sup> ) <sup>b</sup>	18.2 (16.6-21.6)	19.1 (16.7-22.4)‡	17.1 (15.8-20.0)

459 <sup>a</sup>Mean ± SD

460 <sup>b</sup>Median (Interquartile range of 25%-75% of the study population)

461 \*Comparison of median BMI between initial and 6 month visit ( $P < 0.001$ ): Wilcoxon

462 Signed-Rank test

463 \*\*Comparison of median BMI between initial and 6 month visit ( $P < 0.01$ ): Wilcoxon

464 Signed-Rank test

465 ‡Comparison of BMI by race ( $P < 0.05$ ): Mann Whitney U test

466 Table 2. Reported intake of milk, dairy products, and vitamin D-fortified orange juice  
 467 per day from the short food frequency questionnaire

	n	Range	Median (25%-75%) <sup>a</sup>
Initial visit			
Milk (8 oz)	137	0.0 – 10.0	2.0 (1.0-3.0)
Soy, Lactaid <sup>®</sup> or chocolate milk (8 oz)	106	0.0 – 4.5	0.0 (0.0-1.0)
Cheese (1 oz or 1 slice)	133	0.0 – 8.0	1.0 (1.0-2.0)
Yogurt (1 cup)	129	0.0 – 5.0	0.05 (0.0-1.0)
Fortified orange juice (8 oz)	107	0.0 – 3.0	0.20 (0.0-1.0)
6 month visit			
Milk (8 oz)	119	0.0 – 10.0	2.0 (1.0-3.0)
Soy, Lactaid <sup>®</sup> or chocolate milk (8 oz)	92	0.0 – 4.0	0.0 (0.0-1.0)
Cheese (1 oz or 1 slice)	110	0.0 – 8.0	1.5 (1.0-2.0)
Yogurt (1 cup)	106	0.0 – 4.0	0.15 (0.0-1.0)
Fortified orange juice (8 oz)	92	0.0 – 4.0	0.35 (0.0-1.0)

468 <sup>a</sup>Interquartile range of 25%-75% of the study population

469 Table 3. Vitamin D intake for the long (LFFQ) and short food frequency questionnaire  
 470 (SFFQ) by visit and race<sup>a</sup>

Initial Visit	Total (n=140) IU	African American (n=94) IU	Caucasian (n=46) IU
LFFQ	332 ±180*	320 ±168*	357 ±201*
SFFQ	474 ±288	485 ±319	452 ±215
6 Month Visit	(n=122)	(n=83)	(n=39)
LFFQ	349 ±200*	334 ±185*	378 ±227
SFFQ	439 ±234	440 ±222	434 ±263

471 <sup>a</sup>Values are mean ± SD

472 \*Mean values for the LFFQ were lower than for the SFFQ ( $P < 0.01$  to  $0.001$ ): Paired-  
 473 Samples *t*-Test

474 Conversion equation: 40 IU = 1 µg

475 Table 4. Association between the long (LFFQ) and short food frequency questionnaires  
 476 (SFFQ) for vitamin D intake by visit and race

	n	LFFQ and SFFQ	
		$r^a$	$P$ value <sup>b</sup>
Total	257	0.36	<0.001
African American	172	0.26	0.001
Caucasian	85	0.57	<0.001
Initial Visit			
Total	138	0.35	<0.001
African American	92	0.32	0.002
Caucasian	46	0.48	0.001
6 Month Visit			
Total	119	0.38	<0.001
African American	80	0.19	0.087
Caucasian	39	0.66	<0.001

477 <sup>a</sup>Pearson correlation coefficient

478 <sup>b</sup>Significance for  $r$  statistic

479 Table 5. Reliability test for vitamin D intake between two long food frequency  
 480 questionnaires (LFFQs) administered 6 months apart and two short food frequency  
 481 questionnaires (SFFQs) administered 6 months apart

	LFFQs <sup>a</sup>			SFFQs <sup>a</sup>		
	n	<i>r</i>	<i>P</i> value	n	<i>r</i>	<i>P</i> value
Total	119	0.28	0.002	120	0.33	<0.001
African American	79	0.09	0.428	80	0.31	0.006
Caucasian	40	0.52	0.001	40	0.41	0.008

482 <sup>a</sup>Test for reliability: Pearson correlation coefficient

483 Table 6. Probability statistics results for the short food frequency questionnaire (SFFQ)

	Total (258 pairs)
Sensitivity <sup>a</sup>	0.896
Specificity <sup>b</sup>	0.644
Positive Predictive Value <sup>c</sup>	0.776
Negative Predictive Value <sup>d</sup>	0.578

484 <sup>a</sup>Proportion of participants who were correctly identified as having met AAP vitamin D  
485 recommendation (400 IU/day)

486 <sup>b</sup>Proportion of participants who were correctly identified as not having met AAP  
487 recommendation

488 <sup>c</sup>Probability that a person who tested positive for meeting AAP recommendation truly  
489 consumed  $\geq 400$  IU/day

490 <sup>d</sup>Probability that a person who tested negative for meeting the AAP recommendation  
491 actually consumed  $< 400$  IU/day

492 **References**

- 493 **1.** Kumar J, Muntner P, Kaskel FJ, Hailpern SM, Melamed ML. Prevalence and  
494 associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001-  
495 2004. *Pediatrics*. 2009;124:e362-370.
- 496 **2.** Gordon CM, DePeter KC, Feldman HA, Grace E, Emans SJ. Prevalence of  
497 vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med*.  
498 2004;158:531-537.
- 499 **3.** Rajakumar K, Fernstrom JD, Janosky JE, Greenspan SL. Vitamin D insufficiency  
500 in preadolescent African-American children. *Clin Pediatr*. 2005;44:683-692.
- 501 **4.** Gordon CM, Feldman HA, Sinclair L, Williams AL, Kleinman PK, Perez-  
502 Rossello J, Cox JE. Prevalence of vitamin D deficiency among healthy infants and  
503 toddlers. *Arch Pediatr Adolesc Med*. 2008;162:505-512.
- 504 **5.** Dong Y, Pollock N, Stallmann-Jorgensen IS, Gutin B, Lan L, Chen TC, Keeton  
505 D, Petty K, Holick MF, Zhu H. Low 25-hydroxyvitamin D levels in adolescents:  
506 race, season, adiposity, physical activity, and fitness. *Pediatrics*. 2010;125:1104-  
507 1111.
- 508 **6.** Looker AC, Dawson-Hughes B, Calvo MS, Gunter EW, Sahyoun NR. Serum 25-  
509 hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations  
510 from NHANES III. *BONE*. 2002;30:771-777.
- 511 **7.** Willis CM, Laing EM, Hall DB, Hausman DB, Lewis RD. A prospective analysis  
512 of plasma 25-hydroxyvitamin D concentrations in white and black prepubertal  
513 females in the southeastern United States. *Am J Clin Nutr*. 2007;85:124-130.

- 514 **8.** Cole CR, Grant FK, Tangpricha V, Swaby-Ellis ED, Smith JL, Jacques A, Chen  
515 H, Schleicher RL, Ziegler TR. 25-hydroxyvitamin D status of healthy, low-  
516 income, minority children in Atlanta, Georgia. *Pediatrics*. 2010;125:633-639.
- 517 **9.** Weng FL, Shults J, Leonard MB, Stallings VA, Zemel BS. Risk factors for low  
518 serum 25-hydroxyvitamin D concentrations in otherwise healthy children and  
519 adolescents. *Am J Clin Nutr*. 2007;86:150-158.
- 520 **10.** Shils ME, Shike M, Ross C, Caballero B, Cousins R. *Modern Nutrition in Health  
521 and Disease*. 10th ed. Baltimore, MD: Lippincot Williams & Wilkins; 2005.
- 522 **11.** Linus Pauling Institute Micronutrient Research for Optimum Health: Vitamin D.  
523 <http://lpi.oregonstate.edu/infocenter/vitamins/vitaminD/>. Updated November 7,  
524 2008. Accessed November 22, 2009.
- 525 **12.** Misra M, Pacaud D, Petryk A, Collett-Solberg PF, Kappy M. Vitamin D  
526 deficiency in children and its management: review of current knowledge and  
527 recommendations. *Pediatrics*. 2008;122:398-417.
- 528 **13.** Wagner CL, Greer FR. Prevention of rickets and vitamin D deficiency in infants,  
529 children, and adolescents. *Pediatrics*. 2008;122:1142-1152.
- 530 **14.** Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357:266-281.
- 531 **15.** Holick MF. Vitamin D: importance in the prevention of cancers, type 1 diabetes,  
532 heart disease, and osteoporosis. *Am J Clin Nutr*. 2004;79:362-371.
- 533 **16.** National Institutes of Health: Dietary Supplement Fact Sheet: Vitamin D.  
534 <http://dietary-supplements.info.nih.gov/factsheets/vitamind.asp#h3>. Updated  
535 November 13, 2009. Accessed November 22, 2009.

- 536 **17.** Willet W. *Nutritional Epidemiology*. 2nd ed. New York, NY: Oxford University  
537 Press; 1998.
- 538 **18.** Rockett HR, Berkey CS, Colditz GA. Evaluation of dietary assessment  
539 instruments in adolescents. *Curr Opin Clin Nutr Metab Care*. 2003;6:557-562.
- 540 **19.** Rockett HR, Wolf AM, Colditz GA. Development and reproducibility of a food  
541 frequency questionnaire to assess diets of older children and adolescents. *J Am  
542 Diet Assoc*. 1995;95:336-340.
- 543 **20.** Rockett HR, Berkey CS, Colditz GA. Comparison of a short food frequency  
544 questionnaire with the Youth/Adolescent Questionnaire in the Growing Up Today  
545 Study. *Int J Pediatr Obes*. 2007;2:31-39.
- 546 **21.** Marshall TA, Eichenberger Gilmore JM, Broffitt B, Stumbo PJ, Levy SM.  
547 Relative validity of the Iowa Fluoride Study targeted nutrient semi-quantitative  
548 questionnaire and the block kids' food questionnaire for estimating beverage,  
549 calcium, and vitamin D intakes by children. *J Am Diet Assoc*. 2008;108:465-472.
- 550 **22.** Harnack LJ, Lytle LA, Story M, Galuska DA, Schmitz K, Jacobs DR, Gao S.  
551 Reliability and validity of a brief questionnaire to assess calcium intake of  
552 middle-school-aged children. *J Am Diet Assoc*. 2006;106:1790-1795.
- 553 **23.** de Assis MAA, Kupek E, Guimaraes D, Calvo MCM, de Andrade DF, Bellisle F.  
554 Test-retest reliability and external validity of the previous day food questionnaire  
555 for 7-10-year-old school children. *Appetite*. 2008;51:187-193.
- 556 **24.** Araujo MC, Yokoo EM, Pereira RA. Validation and calibration of a  
557 semiquantitative food frequency questionnaire designed for adolescents. *J Am  
558 Diet Assoc*. 2010;110:1170-1177.

- 559 **25.** Fumagalli F, Pontes Monteiro J, Sartorelli DS, Vieira MN, de Lourdes Pires  
560 Bianchi M. Validation of a food frequency questionnaire for assessing dietary  
561 nutrients in Brazilian children 5 to 10 years of age. *Nutrition*. 2008;24:427-432.
- 562 **26.** Ortiz-Andrellucchi A, Henriquez-Sanchez P, Sanchez-Villegas A, Pena-Quintana  
563 L, Mendez M, Serra-Majem L. Dietary assessment methods for micronutrient  
564 intake in infants, children and adolescents: a systematic review. *Br J Nutr*.  
565 2009;102 Suppl 1:S87-117.
- 566 **27.** Centers for Disease Control and Prevention. Overweight and Obesity. Defining  
567 Childhood Overweight and Obesity.  
568 <http://www.cdc.gov/obesity/childhood/defining.html>. Updated October 20, 2009.  
569 Accessed October 12, 2010.
- 570 **28.** Centers for Disease Control and Prevention. Clinical Growth Charts.  
571 [http://www.cdc.gov/growthcharts/clinical\\_charts.htm#Set1](http://www.cdc.gov/growthcharts/clinical_charts.htm#Set1). Updated August 4,  
572 2009. Accessed October 12, 2010.
- 573 **29.** Zemel BS, Carey LB, Paulhamus DR, Stallings VA, Ittenbach RF. Quantifying  
574 Calcium Intake in School Age Children: Development and Validation of the  
575 Calcium Counts!© Food Frequency Questionnaire. *Am J Hum Biol*. 2010;22:180-  
576 186.
- 577 **30.** Rockett HR, Breitenbach M, Frazier AL, Witschi J, Wolf AM, Field AE, Colditz  
578 GA. Validation of a youth/adolescent food frequency questionnaire. *Prev Med*.  
579 1997;26:808-816.
- 580 **31.** Livingstone MB, Robson PJ, Wallace JM. Issues in dietary intake assessment of  
581 children and adolescents. *Br J Nutr*. 2004;92 Suppl 2:S213-222.

- 582 **32.** Kaaks R, Ferrari P, Ciampi A, Plummer M, Riboli E. Part H. Advances in the  
583 statistical evaluations and interpretation of dietary data. Uses and limitations of  
584 statistical accounting for random error correlations, in the validation of dietary  
585 questionnaire assessments. *Public Health Nutr.* 2002;5:969-976.
- 586 **33.** Bandini LG, Schoeller DA, Cyr HN, Dietz WH. Validity of reported energy  
587 intake in obese and nonobese adolescents. *Am J Clin Nutr.* 1990;52:421-425.
- 588 **34.** United States Department of Agriculture. MyPyramid.gov. Inside the Pyramid.  
589 [http://www.mypyramid.gov/pyramid/milk\\_amount.aspx#](http://www.mypyramid.gov/pyramid/milk_amount.aspx#). Updated September 28,  
590 2010. Accessed November 8, 2010.
- 591 **35.** Bellu R, Ortisi MT, Riva E, Banderali G, Cucco I, Giovannini M. Validity  
592 assessment of a food frequency questionnaire for school-age children in northern  
593 Italy. *Nutr Res.* 1995;15:1121-1128.
- 594 **36.** Marshall TA, Eichenberger Gilmore JM, Broffitt B, Levy SM, Stumbo PJ.  
595 Relative validation of a beverage frequency questionnaire in children ages 6  
596 months through 5 years using 3-day food and beverage diaries. *J Am Diet Assoc.*  
597 2003;103:714-720; discussion 720.
- 598 **37.** Taylor C, Lamparello B, Kruczek K, Anderson EJ, Hubbard J, Misra M.  
599 Validation of a food frequency questionnaire for determining calcium and vitamin  
600 D intake by adolescent girls with anorexia nervosa. *J Am Diet Assoc.*  
601 2009;109:479-485, 485 e471-473.
- 602 **38.** Wu H, Gozdzik A, Barta JL, Wagner D, Cole DE, Vieth R, Parra EJ, Whiting SJ.  
603 The development and evaluation of a food frequency questionnaire used in

- 604 assessing vitamin D intake in a sample of healthy young Canadian adults of  
605 diverse ancestry. *Nutr Res.* 2009;29:255-261.
- 606 **39.** Hacker-Thompson A, Robertson TP, Sellmeyer DE. Validation of two food  
607 frequency questionnaires for dietary calcium assessment. *J Am Diet Assoc.*  
608 2009;109:1237-1240.
- 609 **40.** Chen TC, Chimeh F, Lu Z, Mathieu J, Person KS, Zhang A, Kohn N, Martinello  
610 S, Berkowitz R, Holick MF. Factors that influence the cutaneous synthesis and  
611 dietary sources of vitamin D. *Arch Biochem Biophys.* 2007;460:213-217.
- 612 **41.** Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B.  
613 Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple  
614 health outcomes. *Am J Clin Nutr.* 2006;84:18-28.
- 615 **42.** Moore CE, Murphy MM, Holick MF. Vitamin D intakes by children and adults in  
616 the United States differ among ethnic groups. *J Nutr.* 2005;135:2478-2485.
- 617 **43.** Institute of Medicine of the National Academies. Dietary Reference Intakes for  
618 Calcium and Vitamin D. Washington, DC. National Academy of Sciences,  
619 November 2010.
- 620 **44.** Nelms M, Sucher K, Long S. *Nutrition Therapy and Pathophysiology.* Belmont,  
621 CA: Thomson Brooks/Cole; 2007.
- 622 **45.** Klohe DM, Clarke KK, George GC, Milani TJ, Hanss-Nuss H, Freeland-Graves J.  
623 Relative validity and reliability of a food frequency questionnaire for a triethnic  
624 population of 1-year-old to 3-year-old children from low-income families. *J Am*  
625 *Diet Assoc.* 2005;105:727-734.

- 626 **46.** Buzzard IM, Stanton CA, Figueiredo M, Fries EA, Nicholson R, Hogan CJ,  
627 Danish SJ. Development and reproducibility of a brief food frequency  
628 questionnaire for assessing the fat, fiber, and fruit and vegetable intakes of rural  
629 adolescents. *J Am Diet Assoc.* 2001;101:1438-1446.