

2012

Voltage Oscillations and Response Dynamics in a Model of Sensory Hair Cells

Alexander B. Neiman

Kai Dierkes

Benjamin Lindner

Humboldt-University, Berlin, benjamin.lindner@physik.hu-berlin.de

Andrey Shilnikov

Georgia State University, ashilnikov@gsu.edu

Follow this and additional works at: https://scholarworks.gsu.edu/neurosci_facpub



Part of the [Mathematics Commons](#), and the [Neuroscience and Neurobiology Commons](#)

Recommended Citation

Neiman *et al.*: Voltage oscillations and response dynamics in a model of sensory hair cells. *BMC Neuroscience* 2012 13 (Suppl 1):P186.
doi: 10.1186/1471-2202-13-S1-P186

This Presentation is brought to you for free and open access by the Neuroscience Institute at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Neuroscience Institute Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

POSTER PRESENTATION

Open Access

Voltage oscillations and response dynamics in a model of sensory hair cells

Alexander B Neiman¹, Kai Dierkes², Benjamin Lindner^{2,3}, Andrey L Shilnikov^{4,5*}

From Twenty First Annual Computational Neuroscience Meeting: CNS*2012
Decatur, GA, USA. 21-26 July 2012

Sensory hair cells in auditory and vestibular organs rely on active mechanisms to achieve high sensitivity and frequency selectivity with respect to weak stimuli. Self-sustained oscillations in hair cells occur on two very different levels. First, the mechano-sensory hair bundle itself can undergo spontaneous mechanical oscillations. Second, self-sustained electric voltage oscillations across the membrane of the hair cell have been documented in the inner ear of lower vertebrates. The functional significance of these self-sustained voltage oscillations is currently unknown.

We used a Hodgkin-Huxley type model of the basolateral ionic currents of bullfrog sacculus to study genesis of spontaneous voltage oscillation patterns and how the spontaneous oscillations shape the response of the hair cell to external mechanical stimuli. To examine the influence of inevitable fluctuations on the dynamical regimes, we included a stochastic transduction current originating in the Brownian motion of the hair bundle and channel noise arising due to the finite number of mechano-electrical transduction channels [1].

We determined the bifurcation structure of the model in terms of two important ionic conductances, associated with the inwardly rectifier (K1) and Ca²⁺-activated (BK) potassium currents. We found that for large values of BK conductance the system is either at equilibrium or exhibit tonic oscillations. For small values of BK and large values of K1 conductances the dynamics of the model shows diverse patterns of activity including quasi-periodic oscillations, large-amplitude periodic spikes, and bursts of spikes. In particular we found a peculiar transition to bursting through quasiperiodic oscillations with two independent frequencies corresponding to a 2D torus in the phase space of the system. Within small patches of parameter

space at the transition from spiking to bursting and at the spike adding transition, voltage dynamics are chaotic. Furthermore, we showed that thermal fluctuations of mechano-electrical transduction current can lead to chaos in a wide area of parameter space.

We found a high sensitivity and frequency selectivity for the regime of regular spontaneous oscillations in response to sinusoidal stimuli with frequencies $f > 5$ Hz. Hence, an oscillatory voltage compartment might constitute a biophysical implementation of a high-gain amplifier. Cells poised in the chaotic regime of low BK and high K1 conductances showed poor tuning, but provided a high sensitivity to low-frequency variations of external stimuli. In summary, this study shows that the electrical oscillator found in saccular hair cells contributes significantly to nonlinear amplification of external mechanical stimuli. This further supports the idea of nonlinear oscillators playing a crucial role in the operation of the inner ear.

Acknowledgements

Supported by the National Institutes of Health under Grant No. DC05063 (AN), by the National Science Foundation under Grant No. DMS-1009591 (AS), RFFI Grant No. 08-01-00083 (AS), by the GSU Brains & Behavior program (AS) and MESRF "Attracting leading scientists to Russian universities" project 14.740.11.0919 (AS).

Author details

¹Department of Physics and Astronomy, Ohio University, Athens, Ohio 45701, USA. ²Max Planck Institute for the Physics of Complex Systems, Dresden 01187, Germany. ³Bernstein Center for Computational Neuroscience, Humboldt University Berlin, Berlin 10115, Germany. ⁴Neuroscience Institute, Georgia State University, Atlanta, Georgia 30303, USA. ⁵Department of Mathematics and Statistics, Georgia State University, Atlanta, Georgia 30303, USA.

Published: 16 July 2012

Reference

1. Neiman AB, Dierkes K, Lindner L, Han L, Shilnikov AL: **Spontaneous voltage oscillations and response dynamics of a Hodgkin-Huxley type model of sensory hair cells.** *The Journal of Mathematical Neuroscience* 2011, **1**:11, doi:10.1186/2190-8567-1-12.

* Correspondence: ashilnikov@gsu.edu

⁴Neuroscience Institute, Georgia State University, Atlanta, Georgia 30303, USA
Full list of author information is available at the end of the article

doi:10.1186/1471-2202-13-S1-P186

Cite this article as: Neiman *et al.*: Voltage oscillations and response dynamics in a model of sensory hair cells. *BMC Neuroscience* 2012 13 (Suppl 1):P186.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

