

5-9-2015

The Survival and Recovery of $\phi 6$ Virus from Fomites

Richard L. Bearden II

Follow this and additional works at: https://scholarworks.gsu.edu/biology_theses

Recommended Citation

Bearden, Richard L. II, "The Survival and Recovery of $\phi 6$ Virus from Fomites." Thesis, Georgia State University, 2015.
https://scholarworks.gsu.edu/biology_theses/62

This Thesis is brought to you for free and open access by the Department of Biology at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Biology Theses by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

THE SURVIVAL AND RECOVERY OF $\phi 6$ VIRUS FROM FOMITES

by

RICHARD L. BEARDEN II

Under the Direction of Lisa Casanova, PhD

ABSTRACT

Viral transmission from the environment can occur via fomites, but there is uncertainty about which factors most affect viral persistence on fomites. Children are a population highly susceptible to viral infection, and sharing common fomites like toys may spread infection. The objective of this research was to assess the survival of enveloped viruses on the surfaces of children's toys, using bacteriophage $\phi 6$ as a surrogate for enveloped human viruses. The survival of infectious $\phi 6$ virions was observed over a 24 hour period at 22°C and relative humidities of 40% & 60%. On the surface of children's toys, $\phi 6$ was better able to persist at 60% RH (\log_{10} reduction $< 2 \log_{10}$) over a 24 hour period than it was at 40% RH (\log_{10} reduction $> 6 \log_{10}$). If $\phi 6$ virus persists on toy material for up to 24 hours, then viral transmission via shared fomites is certainly significant.

INDEX WORDS: Viral inactivation, Enveloped virus, Transmission, Surrogate, Relative humidity, Temperature

THE SURVIVAL AND RECOVERY OF $\phi 6$ VIRUS FROM FOMITES

by

RICHARD L. BEARDEN II

A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of

Master of Science

in the College of Arts and Sciences

Georgia State University

2015

Copyright by
Richard Louis Bearden II
2015

THE SURVIVAL AND RECOVERY OF $\phi 6$ VIRUS FROM FOMITES

by

RICHARD L. BEARDEN II

Committee Chair: Lisa Casanova

Committee: Sidney Crow

Robert Simmons

Electronic Version Approved:

Office of Graduate Studies

College of Arts and Sciences

Georgia State University

May 2015

DEDICATION

I would like to dedicate this body of work to my family. I'm forever grateful for their encouragement and undying support throughout this process.

ACKNOWLEDGEMENTS

I would like to thank Dr. Casanova for her mentorship and eagerness to help me achieve my goals throughout my graduate studies. Without her wisdom, guidance, and support this project would not have been possible. I would also like to thank Dr. Crow and Dr. Simmons for taking the time to be a part of the committee. Their knowledge and expertise have been invaluable throughout this process. Last, but certainly not least, I would like to thank Dr. Luo for providing me with the skills to use the data to model virus survival beyond the parameters used for the experiments. All of you were essential to this project and I am eternally grateful for your dedication to the completion of it.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	v
LIST OF FIGURES.....	vii
LIST OF TABLES.....	viii
LIST OF ABBREVIATIONS.....	ix
1 INTRODUCTION.....	1
2 Research Design & Methods	4
2.1 Propagation of Virus Stock.....	4
2.2 Suspension Media Experiments.....	4
2.3 Core Survival Experiments.....	5
3 RESULTS.....	7
3.1 Suspension Media Experiments.....	7
3.2 Core Survival Experiments.....	8
3.3 Linear Regression Analysis Fitting the Weibull Model	10
4 Discussion.....	12
REFERENCES	17

LIST OF FIGURES

<p>Figure 1. $\phi 6$ Survival in 1X PBS and DIW. Survival of $\phi 6$ virus suspended in either deionized water or 1X PBS solution for $t=2$ hours at 22°C, 60%RH. Gray column, survival in DIW; white column, survival in 1X PBS. Bars, 95% confidence interval.....</p>	7
<p>Figure 2. $\phi 6$ Survival at 22°C over a 24 hour period. Circles, 60%RH. Squares, 40%RH. Bars, 95% confidence intervals.</p>	8
<p>Figure 3. $\phi 6$ Survival at 22°C, 60%RH. Circles, second aliquots of virus stock. Triangles, first aliquots of virus stock. Bars, 95% confidence intervals.</p>	9
<p>Figure 4. Linear Regression Analysis Fitting the Weibull Model. (a) Log transformations of survival of the second aliquots of virus stock at 40 & 60% RH. Squares, log transformations of inactivation of second aliquots of virus stock at 40% RH; circles, log transformations of inactivation at 60% RH. (b) Log transformations of inactivation of the first and second aliquots of virus stock at 60% RH. Triangles, log transformations of inactivation of the first aliquots of virus stock; circles, log transformations of inactivation of the second aliquots of virus stock.</p>	10

LIST OF TABLES

Table 1. Linear Regression Analysis of the Weibull Function. Linear model fitted with $\ln(\log_{10} N_T/N_0)$ as the response variable and $\ln(t)$, time, as the predictor to get estimations of $n =$ slope of the log time.	11
---	----

LIST OF ABBREVIATIONS

°C	Degrees Celsius
φ6	Bacteriophage phi 6
BE	1.5% Beef Extract
DIW	Deionized Water
Hr	Hour
mL	Milliliter
μL	Microliter
PBS	Phosphate Buffer Solution
PFU	Plaque Forming Units
RH	Relative Humidity
rpm	Rotations Per Minute
TA	Top Agar
TSA	Tryptic Soy Agar
TSB	Tryptic Soy Broth

1 INTRODUCTION

Viral particles consisting of an outer layer of lipids in addition to their protein capsid are called enveloped viruses. Orthomyxoviruses such as influenza viruses H1N1 and H5N1 as well as Coronavirus SARS-CoV and other coronaviruses (CoV) are examples of very pathogenic strains of enveloped viruses [13]. These viruses are responsible for causing many respiratory tract infections that often result in fatality in humans. The WHO estimates that seasonal influenza epidemics alone will result in about 3-5 million cases of severe illness, and about 250,000 to 500,000 deaths globally [21]. Respiratory viruses can be transmitted from person to person through a variety of modes. For instance, influenza can be transmitted via contaminated fomites or inanimate objects, droplets from infected persons, and persistent droplet nuclei suspended in aerosols [5]. The efficiency of various modes of transmission depends partly on the survival of the virus in the environment before it interacts with its next host, and the efficiency with which enveloped viruses spread from one host to the next depends partly on which mode of transmission leaves viruses most vulnerable to inactivation [10]. In environments where a large number of people interact with shared surfaces, there could be continuous contamination of those surfaces with virus and subsequent spread of the virus throughout that population [5]. If viral infection stemming from interaction with contaminated fomites is a major source of viral persistence and spread within a population, then it is extremely important to understand which type of environmental conditions are conducive to a virus's survival on shared surfaces and the risks associated with those surfaces [5]. This knowledge could better equip agencies to foster indoor environmental standards specifically targeted at the inactivation of enveloped viruses using a variety of strategies – controlling/maintaining relative humidity, maintaining temperature, fresh air ventilation within buildings, or surface decontamination/disinfection

protocols [13]. Such standards could prove effective at preventing illness from viruses, especially for vulnerable populations such as children.

It has generally been accepted that enveloped viruses are more sensitive to environmental conditions and have limited survival outside of their host when compared to non-enveloped viruses. Many studies have concluded that humidity is an extremely important factor impacting virus survival but there is a large amount of variation in the findings of these studies [13]. Some studies suggest that as humidity increases, viral inactivation increases [2, 8, 9]. Others suggest that lower humidity levels increase viral inactivation [2, 6]. Currently there is no consensus as to a defined minimum or maximum relative humidity that reduces a virus' survival or ability to infect a new host [13]. Another, often missing, component in some of the existing literature is the significance of the amount of time the virus is exposed to various environmental conditions and how time impacts survival under these conditions [13]. There are also studies that assess virus survival on household surfaces [5]. Virus survival on fomites or surfaces is dependent upon several factors, e.g. the fomite or surface characteristics (porosity, chemical residue, etc.), the matrix surrounding the virus, and the environmental conditions.

If viruses can survive on a fomite for a period of time, than it is probable that viral transmission through direct contact with the contaminated fomite can occur [5]. Virus survival and recovery from children's toys are an exceptional choice because toys are often communal objects. In daycare facilities, schools, or even in homes with multiple children close in age, toys are often shared and circulated from child to child. Children also represent a susceptible population to viral infection. Understanding the factors influencing the persistence of viruses on fomites under varying simulated indoor environments could have a profound impact on how we approach creating environments conducive to viral inactivation, how we reduce the number of

cases of disease, particularly amongst susceptible populations, and how we evaluate the risk posed by contaminated objects that are shared, like children's toys. Therefore, the objective of this experiment is to assess how relative humidity, temperature, and time influence survival of an enveloped virus surrogate, bacteriophage $\phi 6$, on plastic children's toys.

2 Research Design & Methods

2.1 Propagation of Virus Stock

Bacteriophage $\phi 6$ was propagated in *Pseudomonas syringae* (*host*) using the soft agar preparation method [3]. 30mL of host bacterial culture was grown for 24 hours with shaking at 100rpm at room temperature (22°C). 2mL of $\phi 6$ virus stock was added and incubated with shaking for an additional 24 hours. 0.5mL of this virus culture and 0.5mL of host culture were added to 30mL of soft agar (0.7% agar), dispensed into tryptic soy agar (TSA) plates, and incubated at room temperature for 24 hours. The top layer was then harvested, pooled and centrifuged (5900g, 30 minutes at 4°C), and stored as stock in tryptic soy broth (TSB) with 20% glycerol at -80°C.

2.2 Suspension Media Experiments

Host was prepared by adding a 1.5mL volume of host to 150mL of TSB and incubating with shaking at 22°C for 24 hours. 100 μ L of $\phi 6$ virus stock (stored at -80C in 20% glycerol) was diluted into 900 μ L of deionized water (DIW) and 900 μ L of 1X PBS (8.0g NaCl₂, 0.2g KCl, 0.12g KH₂PO₄, 0.91g Na₂HPO₄ /liter of DIW). 10 μ L of virus suspended in DIW (target concentration $\sim 10^7$ plaque-forming units (PFU)) was added to 6 toy coupons (UV sterilized 2cm x 2cm pieces of a child's toy). 10 μ L of virus suspended in 1X PBS was added to an additional 6 toy coupons. 3 coupons from each group of 6 were immediately placed in tubes containing 5mL of 1.5% beef extract (BE) (~ 7.5 pH) using sterile forceps and placed on a shaker (220rpm) at 22°C for 20 minutes. These coupons represented the concentration of virus at t=0hr in both DIW and 1X PBS. Samples were serially diluted in TSB and assayed using the double agar layer method. Plates were incubated at room temperature $\sim 22^\circ\text{C}$ for 24 hours. The remaining 3 coupons for each group were placed in 60% humidity chamber at $\sim 22^\circ\text{C}$. Controlled humidity

and temperature environments were created using sealed glass tanks containing saturated salt solutions (40% - magnesium chloride, 60% - magnesium nitrate). After 2 hours, each coupon was placed into a tube containing 5mL of 1.5% beef extract and placed on the shaker (220rpm) at 22°C for 20 minutes. These coupons represent infectious virus concentration at $t=2\text{hr}$ in DIW and 1X PBS. After 24 hours incubation, the number of plaques on each plate were counted to quantify the number of infectious viruses remaining at after 2 hours in DIW and 1X PBS and the log reduction of viruses was calculated $\log_{10} (N_t/N_0)$, where N_t is the number of viruses at time t and N_0 is the number of viruses at $t=0$.

2.3 Core Survival Experiments

The core survival experiments were carried out in a similar fashion as the suspension media experiments. 100 μL of $\phi 6$ virus stock was diluted into 900 μL of DIW. 10 μL of virus suspended in DIW (target concentration $\sim 10^7$ plaque-forming units (PFU)) was then added 12 coupons. 3 coupons were immediately added to tubes containing 5mL of 1.5% BE (~ 7.5 pH) using sterile forceps and placed on a shaker at (220rpm) for 20 minutes. These coupons represented the concentration of virus at $t=0\text{hr}$. Samples were serially diluted in TSB and assayed using the double agar layer method. Plates were allowed to incubate at room temperature $\sim 22^\circ\text{C}$ for 24 hours. The remaining BE eluent was stored at -80°C . The 9 remaining coupons were grouped into groups of 3 and were placed in either the 40% or 60% humidity chamber at $\sim 22^\circ\text{C}$. Every 2 hours, 1 group of coupons was removed from the humidity chamber and the above procedure was repeated. These experiments were completed at 6-hour intervals ($t=2\text{hr}$, $t=4\text{hr}$, and $t=6\text{hr}$) up to 24 hours for both relative humidity levels at $\sim 22^\circ\text{C}$. After 24 hours of incubation, the number of plaques on each plate were counted to quantify the number of infectious viruses remaining at each time point and the log reduction of viruses was calculated

$\log_{10} (N_t/N_0)$. When $\phi 6$ virus could not be recovered from the plaque assay for the 7-dilution titer, the remaining volume of viral eluent (recovered virus in 1.5% BE) was divided into equal volumes and plated onto TSA plates with 1mL of host and 5mL of TA. The data was analyzed using Excel 2011 (Mac) and GraphPad Prism 5 (GraphPad). The data was fitted to the Weibull model by completing log transformations of the parameter $\log N_t/N_0$ and plotting them against the log transformation of time, t , for survival at each relative humidity [1,7]. Linear regression analysis was conducted to determine the slope of the inactivation line and to predict the shape of the survival curve at each set point of relative humidity.

3 RESULTS

3.1 Suspension Media Experiments

To determine whether the choice of suspension media (PBS vs. deionized water) affected virus survival, virus survival in PBS was compared to deionized water (Figure 1).




Figure 1. $\phi 6$ Survival in 1X PBS and DIW. Survival of $\phi 6$ virus suspended in either deionized water or 1X PBS solution for $t=2$ hours at 22°C , 60%RH. Gray column, survival in DIW; white column, survival in 1X PBS. Bars, 95% confidence interval.

A t test was used to compare virus survival ($\log N_T/N_0$) in deionized water ($n=9$) as a suspension media to 1X PBS ($n=6$). Experiments were conducted at 22°C in a controlled chamber at 60% relative humidity. There was not a statistically significant difference in virus inactivation at $t=2$ hours between the two types of suspension media ($p=0.1564$). PBS was used as the suspension media for subsequent survival experiments.

3.2 Core Survival Experiments




Figure 2. $\phi 6$ Survival at 22°C over a 24 hour period. Circles, 60%RH. Squares, 40%RH. Bars, 95% confidence intervals.

Over a 24 hour period, there was a $\sim 2 \log_{10}$ reduction (99% inactivation) in the number of infectious viruses recovered at 60% RH (Fig.2). At $t=8$ hours post application to the coupon, the number of infectious viral particles declines by $\sim 1 \log_{10}$ and then remains somewhat stable (between 1-2 \log_{10} reduction) up to 24 hours at 60%RH.

At 40%RH, there was a more rapid decline ($\sim 3 \log_{10}$ reduction, 99.9% inactivation) in the number of infectious viral particles recovered after $t=2$ hours post application of the viruses to the coupon, and the level of the inactivation was considerably greater at 40%RH (Fig.2). At $t=8$ to 10 hours post-application most trials were below the detection limit for the double agar layer plaque assay ($>6.5 \log_{10}$ reduction, 99.9999% inactivation) ($n=1$, $n=1$).




Figure 3. $\phi 6$ Survival at 22°C, 60%RH. Circles, second aliquots of virus stock. Triangles, first aliquots of virus stock. Bars, 95% confidence intervals.

In initial survival experiments conducted at 60% RH, two different pools of virus stock were used that were propagated at different times using identical methods from the same initial seed stock. The survival experiments using the first set of aliquots of virus stock (Fig.3), yielded a $\sim 2.5 \log_{10}$ reduction (99% inactivation) in the number of infectious viruses recovered from the coupons after $t=4$ hours of exposure, reaching $\sim 5 \log_{10}$ reduction (99.999% inactivation) after $t=12$ hours. However, completing the double agar layer plaque assay under the same conditions with aliquots of virus stock created at a later time revealed dissimilar results. Survival experiments using the second set of aliquots never reached $>2 \log_{10}$ reduction (99% inactivation) in the number of infectious viruses recovered from the coupons even after $t=24$ hours of exposure. It is important to note that the 40% RH survival experiments were also carried out using the second set of aliquots (Figure 2).

3.3 Linear Regression Analysis Fitting the Weibull Model




Figure 4. Linear Regression Analysis Fitting the Weibull Model. (a) Log transformations of survival of the second aliquots of virus stock at 40 & 60% RH. Squares, log transformations of inactivation of second aliquots of virus stock at 40% RH; circles, log transformations of inactivation at 60% RH. (b) Log transformations of inactivation of the first and second aliquots of virus stock at 60% RH. Triangles, log transformations of inactivation of the first aliquots of virus stock; circles, log transformations of inactivation of the second aliquots of virus stock.

After fitting the data to the Weibull model, a linear regression analysis was conducted for each set point of relative humidity. The log transformations of average inactivation ($\ln(-\log_{10}(N_T/N_0))$) (response variable) at each time point were plotted against the log transformation of the time ($\ln(t)$) at which survival was being evaluated. For the second set of aliquots of virus stock, the slopes for the survival plot at 40 and 60% RH are significantly different ($p=0.0448$).

However, the slopes of the survival plot for both sets of aliquots of virus stock at 60% RH were not significantly different ($p=0.3002$).

Variable	n	Standard Error	p -value	R^2
40% RH	0.4772	0.0914	0.0070	0.9362
60% RH 1	1.385	0.3512	0.0043	0.8951
60% RH 2	1.088	0.3519	<0.0001	0.8573

Table 1. Linear Regression Analysis of the Weibull Function. Linear model fitted with $\ln(\log_{10} N_T/N_0)$ as the response variable and $\ln(t)$, time, as the predictor to get estimations of n = slope of the log time.

Modeling this experimental data with the Weibull model and comparing the slopes of the inactivation plots at each set point of relative humidity, allows for extrapolation of what the actual shape of the survival curve may be. Linear regression analysis of this model reveals that the model fits the expectation of the data. At 40% RH, the slope (n) of the survival plot = 0.4772. When $n < 1$, the rate of inactivation will decrease over time due to significant inactivation at early time points. At 60% RH, $n > 1$, indicating that the inactivation of $\phi 6$ will increase over time at this set point of temperature and relative humidity.

4 Discussion

These experiments show that $\phi 6$ virus is able to persist at 22°C and 60% RH for up to 24 hours, and potentially longer. The rate of inactivation appears to be slower at 60% RH than at 40% over a 24 hour period. The rate of inactivation of infectious virions is much more pronounced at 40% RH, with virus reaching undetectable levels 8 hours post application ($>6 \log_{10}$ reduction, $\sim 99.9999\%$ inactivation). However, at 60% RH, the amount of infectious virus recovered from the toy coupons declines most around $t=8$ hours post application, and then appears to remain relatively stable up to $t=24$ hours ($\sim 1-2 \log_{10}$ reduction, $\sim 90-99\%$ inactivation). This data suggests that $\phi 6$ is better able to resist inactivation at 22°C and 60% RH on the surface of the toy coupon. This is contrary to some of the existing literature centered on the survival of enveloped viruses, where virus survival appears to be more stable at lower relative humidities [2, 9, 10].

There is 99.9999% inactivation of $\phi 6$ within the first 10 hours of exposure to 40% RH. It is expected that the rate of inactivation will decrease over time due to the vast majority of virions becoming inactivated early on in their exposure to this relative humidity. The plot of survival at 40% RH supports this prediction. In comparing n at both 40 and 60% RH, the opposite is true for the rate of inactivation at 60% RH. Although inactivation of the second aliquots of $\phi 6$ was no greater than 99% inactivation after 24 hours of exposure to the simulated environment, the slope calculated from the linear regression analysis suggests that the rate of inactivation will increase over time. However, the plot of survival at 60% relative humidity for the second aliquots of virus stock does not necessarily support this. Additional experiments carried out beyond 24 hours at 60% relative humidity are necessary to validate the model prediction.

Toys were chosen as a fomite for these experiments because they have been shown to be significant sources of viral and bacterial contamination in healthcare settings and nurseries [7, 11, 16]. Due to their age, children are often much more susceptible to viral and bacterial infections. They often lack the immunity provided by previous exposure or vaccination, especially when very young [16]. They also exhibit behavior conducive to contracting infections, such as putting foreign objects in their mouths or failure to wash their hands. In healthcare and nursery environments, toys are communal objects and often shared between children [11]. As a result, they are likely significant sources of the spread of infection throughout populations in those environments. It has been previously shown that respiratory syncytial virus, an enveloped virus, can survive on the surface of toys for up to 6 hours [11]. At 22°C and 60%RH, $\phi 6$ inactivation was between 90-99% 8 hours post application to the surface and never reached greater than 99% inactivation over a 24 hour period. Although $\phi 6$ virus is non-pathogenic to humans, its survival on the surface of the toy coupon highlights the necessity of examining how enveloped viruses become inactivated in indoor environments and the importance of fostering effective decontamination protocols for communal objects, like toys, that children interact with in these environments.

Virus survival and persistence on surfaces is influenced by several factors, whether it be the viral species, the surface the virus is applied to, the temperature and humidity of the environment, or the media the virus is suspended in. It is somewhat difficult to compare the findings of this paper with those of the existing literature because each set of experiments differs by 1 or more of these factors. For instance, Casanova and Waka completed survival experiments using $\phi 6$ virus and identical research methods [3]. The only major differences between the experiments within this paper and the ones used for their experiments was the suspension media

used (1X PBS) and the surface to which the virus was applied; coupons of N95 respirator material. $\phi 6$ virus survival at 22°C appeared to be much more stable at 40%RH (<2 log₁₀ reduction, 90-99% inactivation) on N95 material then at 60% RH (>3 log₁₀ reduction, 99.9% inactivation) over a 24 hour time period [3]. For the survival experiments carried out for this paper at 22°C and 40% RH, >3 log₁₀ reduction (>99.9% inactivation) of $\phi 6$ was achieved after 2 hours post application of the virus to the coupon. At 60% RH, the level of inactivation (~5 log₁₀ reduction, 99.999% inactivation) achieved at t=12hr using the first set of aliquots seemed to be somewhat comparable to the inactivation achieved by Casanova and Waka (~4 log₁₀ reduction, 99.99% inactivation). However, the second set of aliquots did not reach >2 log₁₀ reduction (99% inactivation) until 24 hours after application of the virus to the coupon. That degree of $\phi 6$ inactivation is reached by 6 hours post application by Casanova and Waka. It is also important to note that virus stock propagated at different times from the same seed stock appear to have varying rates of inactivation ($n=1.385$, $n=1.088$) when exposed to identical environmental conditions (22°C, 60% RH). The survival plots for the first set of aliquots of viral stock exhibit a significantly greater rate of inactivation than the survival plots of the second set of aliquots at 60%RH. At t=2 hours post application, both populations have similar rates of inactivation, but at t=4 hours post application, the inactivation diverges for the two populations of virus. Although the rate of inactivation for the first set of aliquots at 60% RH is higher than that of the second set, the slopes from the linear regression analysis were not significantly different ($p=0.3002$).

In a recent paper, filoviruses ZEBOV (Zaire Ebola virus) and MARV (Lake Victoria Marburg virus) were used for long-term survival experiments. Neither virus could be recovered from glass, plastic, or metal when applied to the surface at ~22°C, 55±5% RH suspended in guinea pig sera or tissue culture media. Virus was only detectable when applied to glass or

plastic at 4°C and 55±5% RH, and remained detectable 14 days post application [15]. A juxtaposition of this discovery with the results of the experiments using $\phi 6$ within this paper reveals a dichotomy. Although $\phi 6$ and filoviruses share structural similarities (envelope and nucleocapsid), they do not have the same resistance to similar environmental conditions. This is also true for viruses of the same species. In another paper, the survival of 2 strains of coronavirus in suspension and dried onto various surfaces to estimate the risk of viral transmission on items found in hospital settings [17]. They found that when HCV-229E and HCV-OC43 were dried onto surfaces, infectious HCV-229E was still detectable up to 6 hours post drying on sterile sponges (4% recovery) and aluminum (8% recovery) whereas, HCV-OC43 was undetectable (0% recovery) on any surface after 3 hours post drying on those surfaces.

One interesting explanation for the variation in survival of viruses is what happens at the air-water interface (AWI) when viruses are suspended in water on surfaces. The theory is that the proteins of virions are strongly attracted to the AWI, especially when suspended in liquids of high ionic strength [19, 20]. These proteins are often embedded in the envelope of the virion. When these proteins interact with the AWI, they cause the loss of the envelope due to hydrophobic interactions and rearrangement of the viral capsid. The rearrangement of the capsid, which is a highly organized structure in many viruses, would lead to the release of the viral genome and subsequent inactivation of the virus [12]. Other experiments have conclusively shown that viruses meeting the AIW become inactivated but it is not the sole reason viruses become inactivated [19, 20]. If multiple virions within a suspension become attracted to the AWI and form an aggregate of virions, it is possible that those virions at the AWI will become inactivated but will also form a barrier, blocking additional virions from reaching the AWI [12, 19, 20]. At low RH, or over a long period of time at any humidity, it would be expected that the

liquid volume would decrease in the suspension due to evaporation, and the viruses will become larger aggregates over time (more concentrated). This aggregation could have a protective effect for viruses at the interior of the viral suspension [12]. However, at 40% RH, perhaps the benefits of aggregation are outweighed by the increased rate of the disappearance of water, leading to desiccation.

Although virally contaminated environmental surfaces can contribute to the transmission of virus from one person to another, it is unclear how significant this mode of transmission is. This uncertainty stems from the fact that virus survival appears to be quite dynamic, in that it is impacted by a number of factors including but not limited to temperature, humidity, surface material, and the intrinsic properties of the virus. The impact of one particular environmental factor is hard to estimate because each species and/or strain of virus appears to respond to similar environmental conditions in a unique way. More investigation is needed to better understand the survival kinetics of viruses under various environmental conditions and the factors that confound them, particularly at the molecular level.

REFERENCES

1. Berger, R. W., & Lawrence, K. (1974). Estimating Weibull Parameters by Linear and Nonlinear Regression. *Technometrics*, 16(4), 617-619.
doi:10.1080/00401706.1974.10489245
2. Casanova, L., et al. Effects of air temperature and relative humidity on coronavirus survival on surfaces. *Applied and Environmental Microbiology*, 76 (9):712–2717.
3. Casanova, L., & Waka, B. Survival of a Surrogate Virus on N95 Respirator Material. *Infection Control and Hospital Epidemiology*, 34, 1334-1335.
4. Enstone J. Influenza and related infection control issues. In: Van-Tam J, Sellwood C, eds. Introduction to pandemic influenza. Wallingford: CAB International, 57–72.
5. Greatorex JS, Digard P, Curran MD, Moynihan R, Wensley H, et al. (2011) Survival of Influenza A(H1N1) on Materials Found in Households: Implications for Infection Control. *PLoS ONE* 6(11): e27932. doi:10.1371/journal.pone.0027932
6. Harper, G.J. Airborne Micro-organisms: Survival Tests with Four Viruses. Microbiological Research Establishment. *Journal of Hygiene*, 59, 479-486.
7. Ibfelt, T., E.H. Engelund, A.C. Schultz, and L.P. Andersen. "Effect of cleaning and disinfection of toys on infectious diseases and micro-organisms in daycare nurseries." *Journal of Hospital Infection* 89 (2015): 109-115. Print.
8. Kim, S. J., Jiyeon, S., Lee, J. E., & Ko, G. (2012). Temperature and Humidity Influences on Inactivation Kinetics of Enteric Viruses on Surfaces. *Environmental Science & Technology*, 46, 13303-13310.
9. Koep, T., et al. Predictors of indoor absolute humidity and estimated effects on influenza virus survival in grade schools. *BMC Infectious Diseases*, 13, 1471-2334.
10. Kohn, M., & Shaman, J. Absolute humidity modulates influenza survival, transmission, and seasonality. *Proceedings of the National Academy of Sciences*, 106, 3243-3248.
11. Little, Kathryn, and Stanley Cutcliffe. "The safe use of children's toys within the healthcare setting." *Nursing Times* 102.38 (2006): 34-37. Print.
12. Mbithi, J. N., Springthorpe, V. S., & Sattar, S. (1991). Effect of Relative Humidity and Air Temperature on Survival of Hepatitis A Virus on Environmental Surfaces. *Journal of Hospital Infection*.

13. Memarzadeh, F. Literature Review of the Effect of Temperature and Humidity on Viruses. *ASHRAE*, 12, 1049-1060.
14. Myatt, T., et al. Modeling the airborne survival of influenza virus in a residential setting: the impacts of home humidification. *Environmental Health*, 9, 1-7.
15. Piercy, T. J., Smither, S. J., Steward, J. A., Eastaugh, L., & Lever, M. S. (2010). The survival of filoviruses in liquids, on solid substrates and in a dynamic aerosol: Survival of filoviruses. *Journal of Applied Microbiology*. doi:10.1111/j.1365-2672.2010.04778.x
16. Posfay-Barbe, Klara M., Danielle M. Zerr, and Didier Pittet. "Infection Control in Paediatrics." *Lancet Infectious Diseases* 8 (2008): 19-31. Print.
17. Sizun, J., Yu, M. W., & Talbot, P. J. (2000). Survival of human coronaviruses 229E and OC43 in suspension and after drying on surfaces: a possible source of hospital-acquired infections. *Journal of Hospital Infection*. doi:10.1053/jhin.2000.0795
18. Thomas, Y. et al. Survival of influenza virus on human fingers. *Clinical Microbiology and Infection*, 20, O58-O64.
19. Thompson, S. S., & Yates, M. V. (1999). Bacteriophage Inactivation at the Air-Water-Solid Interface in Dynamic Batch Systems. *Journal of Hospital Infection*.
20. Thompson, S. S., Flury, M., Yates, M. V., & Jury, W. A. (1998). Role of the Air-Water-Solid Interface in Bacteriophage Sorption Experiments. *Journal of Applied Microbiology*.
21. World Health Organization. (2014, March). WHO | Influenza (Seasonal). Retrieved from <http://www.who.int/mediacentre/factsheets/fs211/en/>