

5-17-2019

# An Evaluation Study Examining the Understandability of STI/HIV Health Education Print Materials Provided by Health Clinics Serving Hispanic and Latino Populations of DeKalb County, Georgia

Rebecca Butler

Follow this and additional works at: [https://scholarworks.gsu.edu/iph\\_theses](https://scholarworks.gsu.edu/iph_theses)

---

## Recommended Citation

Butler, Rebecca, "An Evaluation Study Examining the Understandability of STI/HIV Health Education Print Materials Provided by Health Clinics Serving Hispanic and Latino Populations of DeKalb County, Georgia." Thesis, Georgia State University, 2019.  
[https://scholarworks.gsu.edu/iph\\_theses/655](https://scholarworks.gsu.edu/iph_theses/655)

This Thesis is brought to you for free and open access by the School of Public Health at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Public Health Theses by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

An Evaluation Study Examining the Understandability of STI/HIV Health Education Print  
Materials Provided by Health Clinics Serving Hispanic and Latino Populations of DeKalb  
County, Georgia

By

Rebecca L Butler

Health Promotion and Behavior

A Thesis Submitted to the Graduate Faculty  
of Georgia State University in Partial Fulfillment  
of the  
Requirements for the Degree

MASTER OF PUBLIC HEALTH

ATLANTA, GEORGIA  
30303

## TABLE OF CONTENTS

ACKNOWLEDGMENTS	iii
LIST OF TABLES	4
Chapter 1 BACKGROUND	
1.1 Abstract	7
1.2 Research Questions	8
1.3 Terminology	11
Chapter 2 REVIEW OF THE LITERATURE	
2.1 Transmission Categories and Health Outcomes for Hispanic and Latinos with HIV in the United States	12
2.2 Social Determinants of HIV in Hispanic and Latino populations	13
2.3 Previous Health Literacy Research	15
2.4 The Utilization of Health Literacy Assessment Tools in Previous Research	17
Chapter 3 METHODS AND PROCEDURES	
3.1 Study Design	19
3.2 Selection of the Evaluation Tool	19
3.3 Raters	20
3.4 Scoring	20
3.5 Evaluation Plan	21
Chapter 4 RESULTS	
4.1 Prevalence of qualifying data for evaluation	22
4.2 Rater agreed scores of the PEMAT-P questionnaire	23
Chapter 5 DISCUSSION AND CONCLUSION	
5.1 Discussion of Research Questions	24
5.2 Study Strengths and Limitations	25
5.3 Conclusion	26
REFERENCES	27
APPENDIX A	28

## Acknowledgements

The development, execution, and presentation of this research would not be possible without the guidance and expertise of my thesis committee. Dr. Sheryl Strasser, thank you for your willingness to chair my committee and for all of your encouragement and feedback. I hope that I made you proud as your 92<sup>nd</sup> thesis student! Thank you to Johnny Gossett for helping me to navigate local HIV surveillance data and for being a great example of how to talk to people so that they will listen whilst educating in the field. Thank you to my honorary thesis committee. Thank you to my Student Health Promotion colleagues and supervisors. Thank you to my mom and family for instilling the expectation of finding meaningful work and finding the courage and motivation to master it.

List of Tables

Table 1.1 Simple Statistical Summary of PEMAT-P Section Scores for Print Materials (n=11)

	Mean	Std. Deviation	Min	Max
Understandability Score (max =17)	12	2.56	7	15
Actionability Score (max=7)	3.54	1.03	1	5

Table 1.2 Rater Agreed PEMAT-P Scores (%) for All Items About the Use of Visual Aids

Item #	Agree=1 point (frequency)	Disagree=0 points (frequency)	N/A=0 points (frequency)	n=
15. The material uses visual aids whenever they could make content more easily understood (e.g., illustration of healthy portion size).	3	8		11
16. The material’s visual aids reinforce rather than distract from the content.	9	1	1	11
17. The material’s visual aids have clear titles or captions.	5	5	1	11
18. The material uses illustrations and photographs that are clear and uncluttered.	9	1	1	11
19. The material uses simple tables with short and clear row and column headings.	7	1	3	11

Table 1.4 Adapted From: Shoemaker SJ, Wolf MS, Brach C. Patient Education Materials Assessment Tool for Printable Materials (PEMAT- P). (Prepared by Abt Associates under Contract No. HHS290200900012I, TO 4). Rockville, MD: Agency for Healthcare Research and Quality

## List of Figures

Figure 1.1 Histogram of PEMAT-P scores of evaluated materials





Figure 1.2 Comparison of PEMAT-P section scores (%) for Understandability and Actionability



An Evaluation Study Examining the Understandability of STI/HIV Health Education Print  
Materials Provided by Health Clinics Serving Hispanic and Latino Populations of DeKalb  
County, Georgia

By

Rebecca L Butler

Approved:

April 18, 2019

Committee Chair

Dr. Sheryl Strasser PhD, MPH, MSW, MCHES

Committee Member

Johnny Gossett Jr.

## **ABSTRACT**

The prevalence of Human Immunodeficiency Virus (HIV) in Hispanic and Latino populations in the United States is on the rise. In 2015, 26% of all HIV diagnoses in the United States were among Hispanic and Latinos. To effectively combat the prevalence of HIV in this population, it is necessary to examine the efficacy and accessibility of current interventions. Printed educational materials (i.e. pamphlets, brochures, and fact sheets) are low-cost and effective intervention strategies if the material is clear to the reader. This study aims to provide further evidence of the understandability of printed educational resources on safe sex practices and HIV in health clinics



## CHAPTER 1

### An Evaluation Study Examining the Understandability of STI/HIV Health Education Print Materials Provided by Health Clinics Serving Hispanic and Latino Populations of DeKalb County, Georgia

Great strides have been made to improve health-related outcomes for patients receiving early treatment for HIV. However, the rate of transmission and prevalence of diagnosis remains a major public health concern. Although risk factors for the virus are the same across all populations, HIV disproportionately affects racial and ethnic minority populations (CDC, 2017). In 2016, the Hispanic and Latino population accounted for 25% (n=9813) of all new HIV diagnoses in the United States (CDC, 2017). If the Centers for Disease Control and Prevention is to achieve their primary goal of a future free from HIV (CDC, 2018), as outlined in the Division of HIV/AIDS Prevention's (DHAP) Strategic Plan (CDC, 2017), research and resources must specifically address HIV-related disparities in Hispanic and Latinos.

*Health literacy* refers to the ability of an individual to obtain and understand health information as part of the process of making informed, appropriate healthcare decisions (U.S. Department of Health and Human Services, 2000). Individual and systemic factors influence health literacy including,

- Environmental and economic access to care
- Cultural attitudes towards health and disease prevention
- The efficacy of communication between the patient and healthcare provider
- The structure of the healthcare system (U.S. Department of Health and Human Services, 2001)

The ability of an individual or group to access, understand, and act on evidence-based information yields positive health outcomes. Conversely, published studies across disciplines find that low health literacy yields negative health outcomes (Berkman et al., 2011). In 2016, the World Health Organization prioritized health literacy as a cornerstone goal of their *Moving Forward, A Plan for the Next Fifteen Years* agenda as part of the 9<sup>th</sup> Global Conference on Health Promotion (WHO, 2018). There is a need for research that examines the impact and efficacy of HIV prevention communication materials for Hispanic and Latino populations living in the United States.

### **Prevalence of HIV in southeastern United States**

HIV is not evenly distributed across the United States (CDC, 2018). Per 100,000 people, the rate of diagnosis in the southern region (i.e. AL, AR, DE, DC, FL, GA, KY, LA, MD, MS, NC, OK, SC, TN, TX, VA, WV) is the highest in the country at 24.9 (CDC, 2018). Surveillance data indicates that,

1. More than half of all HIV diagnoses (n=38739) in 2017 were distributed across the southern states. CDC surveillance for the same year indicates that Hispanics and Latinos accounted for 21% of new HIV diagnosis by race in the south (CDC, 2016).
2. Southern states have the greatest number of individuals estimated to be living with HIV (CDC, 2015)
3. In 2016, the CDC's *Morbidity and Mortality Weekly Report* indicated that just 60% of Hispanics/Latinos were linked to medical care within 90 days

of diagnosis. This is well below the national benchmark of 85% set forth by the U.S. Department of Health and Human Services.

United States federal health agencies have urged state health departments, policymakers, and prevention partners to prioritize HIV prevention efforts in regions and populations with the greatest disparities. In 2018, the CDC allocated \$216 million dollars to the southern United States for the purpose of improving HIV prevention and care outcomes (CDC, 2018). The *National HIV/AIDS Strategy for the United States* (2015) sets forth specific prevention guidelines for Hispanic and Latino populations in southern states including, “Requiring partners to use the latest, most cost-effective interventions and increasing cultural competency and tailoring prevention programs” (HIV.gov, 2017).

Print materials, such as brochures and pamphlets, are a commonly utilized, low-cost prevention strategy for disseminating health information. The aim of this study is to provide further evidence of the understandability of printed educational resources on HIV in health clinics in northeast Dekalb County, Georgia. This county was selected for its high rate of Hispanic and Latino residents within its municipalities including the cities Doraville and Chamblee (U.S. Census Bureau, 2018). The study methodology is designed to answer the following questions,

1. Do free and low-cost clinics that offer HIV testing in within a 10-mile radius of the Dekalb County zip code, 30341 provide educational print materials about HIV and safe-sex practices?
2. Do the evaluated print materials promote understandability and actionability when they are evaluated via a health literacy assessment tool?

3. How do the evaluated print materials make use of visual aides to promote understandability?

### **Terminology**

*Hispanic* or *Latino* are identifiers that are assigned to an ethnic category of people who are of a Spanish or Latin American country of origin. To refer to a population as *Hispanic* or *Latino* does not refer to a racial category (OMB, 1997). *Hispanic* and *Latino* are often referred to together or interchangeably when denoting a demographic category in published research. In 1997, the Federal Office of Management and Budget revised the standards for the classification of federal data on race and ethnicity to include, “Hispanic or Latino” (OMB, 1997). The classification, *Hispanic and Latino* will be used henceforth to describe the population of interest for this research.

## **Chapter II** **REVIEW OF THE LITERATURE**

### **Transmission Categories and Associated Health Outcomes for Hispanic/Latinos with HIV in the United States**

Transmission category statistics are paramount to identifying subpopulations who are particularly at-risk for HIV for the purpose of developing tailored prevention plans. The epidemiological mode of transmission for HIV is distributed amongst 6 categories. In adolescent and adult males, HIV acquisition is most frequently attributed to male-to-male sexual contact. In adolescent and adult females, HIV is most frequently attributed to heterosexual contact with a person known to have, or be at risk for HIV (CDC, 2017). In 2016, 85.6% of all Hispanic or Latinos diagnosed with HIV attributed their infection to sexual contact (n=8810).

Espinoza et al. (2008) looked for trends in mode of transmission and illness progression of HIV/AIDS in foreign-born Latino and Hispanic individuals residing in the United States. Espinoza et al. (2008) analyzed secondary CDC surveillance data to identify trends in incidence of HIV diagnosis, the progression of HIV to AIDS, and mortality of individuals who received an AIDS diagnosis based on country of birth. Data was analyzed by applying Poisson regression “to calculate the estimated annual percentage change in the number and rate of HIV diagnoses” (Espinoza et al., 2008). A logistic regression model was used to examine possible associations between birthplace and HIV-to-AIDS intervals that occurred in under 12 months. Qualifying criteria for inclusion in this study were reported diagnoses of HIV among Hispanics in the United States from 2003-2006 who reported a birthplace. Results indicate that 60% (n=14,954) of participants reported being born outside the United States. Additionally, Espinoza et al., found significant characteristic differences in HIV acquisition based on birthplace,

The transmission categories for HIV infection in US-born Hispanics were distributed as follows: male-to-male sexual contact (50%), followed by high-risk heterosexual contact (26%), IDU (18%), and male-to-male sexual contact and IDU (4%). Among persons born in Puerto Rico, larger proportions had been infected through high-risk heterosexual contact (40%) and IDU (33%) than through male-to-male sexual contact (23%)

The findings of this study highlight the number of foreign-born Hispanic and Latino individuals in the United States who have been diagnosed with HIV and inspires further inquiry about what HIV the prevention challenges that immigrant populations face. Furthermore, the categorial breakdown of qualifying participants by birthplace highlights the demographic diversity of Hispanics and Latinos with HIV living in the United States. Because the mode of transmission and rates of survival significantly vary based on Hispanic subgroups, our interventions should not be *one size fits all* (Freiman, 1998). The findings of this study encourage further investigation into what types of HIV educational materials are available to Latino and Hispanics seeking testing and treatment in a healthcare setting.

### **Determinants of HIV in Hispanic and Latino populations in the United States**

Ryan et al. (2009) conducted a qualitative study to examine the impact of familial rejection on predicting of negative health outcomes in self-identifying lesbian, gay, and bisexual young adults. The authors of this study developed quantitative scales to assess the associated frequency of family and caregiver rejection of participants' sexual orientation based on a previous qualitative study of in-depth interviews that were thematically coded for behaviors of

acceptance and rejection. The survey also contained indicators of negative health outcomes related to mental health and high-risk sexual behaviors. The population of interest for this study was n=244 white and Latino gay, lesbian, and bisexual young adults between the ages of 21 and 25. Participants completed a self-reported computer survey. Survey outcomes were combined with a dichotomous scoring system to create a Family Rejection Scale, with values ranging from 0 to 51. High rates of family rejection were positively associated with negative health outcomes. Latino men reported the highest rate of family rejection (mean=24.52) during adolescence of a gay, lesbian, or bisexual sexual orientation. LGB adults of all races were 3.4 times more likely to engage in unprotected sex if they reported a high level (>35.2) of family rejection in adolescence. This study exemplifies possible cultural barriers for familial support for minority LGBTQ populations' ability to be open about their sexual orientation. Furthermore, this study suggests a possible relationship between family rejection, unprotected sex, and negative health outcomes. When considered in a larger context, one such recommendation that could be made with regard to health literacy about HIV and safe sex practices may be to include resources for counseling and mental health services in communications literature that is consumed by minority, LGBTQ populations.

### **Previous Health Literacy Research**

In 2003, the National Assessment of Adult Literacy (NAAL) published its first national survey of estimated health literacy by state. The data indicates that 12% of U.S. adults scored at or above the NAAL's definition of proficient health literacy. Task examples for this category include, the ability to define a medical term within a complex prose document and the ability to use a table to calculate a share of employee health

insurance costs (Kutner et al., 2006). The report also indicates that 35% of adults that fell below the group of *basic health literacy* identified as Hispanic. One such task example for this category is, “read a short set of instructions and identify what is permissible to drink before a medical test” (Kutner et al.). The most significantly associated variables for health literacy across all racial categories were highest level of educational attainment and insurance status (CDC, 2017).

Edward et al. (2018) assessed the impact of health literacy in Hispanic and Latino populations on their ability to access and utilize health insurance. This study measured health literacy (HL) and health insurance literacy (HIL) on healthcare enrollment of minority populations in Massachusetts. The study design utilized a community engagement model to enroll participants and conduct activities in Spanish-speaking churches. N=239 participants 18 years of age and older, who self-identified as Hispanic or Latino, who met the enrollment criteria, participated in this study. Participants completed a detailed demographic survey that included questions about insurance status, general health status, and the frequency of utilizing healthcare. (Edward et al., 2018).

Health insurance literacy was assessed through the use of a knowledge-based survey that asked participants to define words such as, “copay”, “premium”, and “deductible”. Participants were scored as having “adequate” HIL if they were able to define at least one of these terms and were scored as having “inadequate” HIL. Health literacy was measured by using an 18-item Short Assessment of Health Literacy Scale in Spanish (SAHL-S). Scores for the SAHL-S range from 0 to 18, with a score between 0 to 14 suggesting inadequate health literacy and scores between 15 and 18 indicating adequate health literacy (Edward et al., 2018). Edward et al. (2018)



found that 60.9% of participants had health insurance at the time of the study, and 70% of participants reported having accessed health care in the United States in the last year. 56% (n=77) participants scored low on the Health Literacy Scale in Spanish. 93% of participants (n=129) participants and were unable to define at least one basic health insurance term and were designated as having “inadequate” health insurance literacy. This study emphasizes a lack of health insurance literacy in Spanish and Latino populations and highlights a barrier to care to the utilization of health services.

In an effort to describe the impact of health literacy on chronic disease management in the Hispanic and Latino population, Williams and Parker, (1998) conducted a cross-sectional survey to examine the literacy of hypertension and diabetes patients. Participants (n=516) were recruited during a three-month period from two public hospitals; Harbor-UCLA Medical Center in Los Angeles, California and Grady Memorial Hospital in Atlanta, Georgia. Eligible participants for the study were individuals over 18 years of age who had no psychiatric illness and who had a confirmed diagnosis of hypertension or Type I and Type II diabetes. Participants gave relevant demographic information and completed a 50-question and 17-question numeracy survey called, the Test of Functional Health Literacy in Adults<sup>1</sup> (TOFHLA) (eg. Reading prescription bottles, appointment slips, and instructions for diagnostic tests [Williams & Parker, 1998]). The TOFHLA was administered in the preferred language for each participant. Scores for this survey range from 0-100 and are classified into categories (*0-59 inadequate score, 60-75 marginal, and 76-100 adequate for literacy*). Additionally, each participant was surveyed about specific knowledge related to his or her disease. 69% of Spanish-speaking participants were classified as having inadequate and marginal health literacy. Williams and Parker (1998) also

---

established a correlation between health literacy and patients' knowledge about their illness. For example, the mean diabetes knowledge of patients who were classified as having inadequate health literacy (n=50) was  $5.2 \pm 6.4$  (CI=95%).

Williams and Parker, (1998) effectively identified a relationship between low health literacy and low knowledge about the diseases included in the study design for Hispanic and Latino participants. This study also highlights the challenge of how to educate low-literacy patients with quality and appropriate print materials that are understandable. Similar to HIV, hypertension is a chronic illness that requires early intervention and home management for favorable health outcomes.

### **The Utilization of Health Literacy Assessment Tools in Previous Research**

The *National Action Plan to Improve Health Literacy* (HHS, 2010) cites evidence-based research that emphasizes the importance of plain language health communications. Such literature most be written in a manner that promotes *understandability* and *actionability* in its content. Additionally, this report cites evidence that established a link between limited health literacy and limited English proficiency (LEP). The report states, "Therefore, interventions for vulnerable populations, such as those with LEP, should focus on health literacy *and* language to promote two-way, interactive communication" (HHS, 2010).

Shoemaker et al. developed the questionnaire in response to this report as a "new measure of understandability and actionability" for patient information (Shoemaker et al., 2014) that is comprehensive, yet simple enough to be reliably applied by health professionals and lay people alike. To develop the PEMAT, Shoemaker et al. reviewed existing health literacy instruments for items and concepts to be included. The methodology for the development of the

PEMAT was to first, garner ideas from a panel of experts in “health literacy; health communications; content creation, patient engagement and health information technology” (Shoemaker et al., 2014). Together, the panel and research team worked to develop assessment items for each scale of the questionnaire. The final item pool was refined and decided upon by conducting four rounds of reliability testing using a total of 22 raters. The authors of this study found,

For construct validation with consumers ( $n = 47$ ), we found significant differences between actionable and poorly-actionable materials in comprehension scores (76% vs. 63%,  $p < 0.05$ ) and ratings (8.9 vs. 7.7,  $p < 0.05$ ). For understandability, there was a significant difference for only one of two topics on consumer numeric scores.

## Chapter III

### METHODS AND PROCEDURES

The study design is a qualitative assessment of the health literacy of primary data of print health communications materials from clinics and healthcare providers. Inclusion criteria clinic selection stipulates that,

Clinic results were filtered and refined under the following criteria:

1. The provider is located within a ten-mile radius of the zip code 30341
2. The provider is free or low-cost
3. The provider offers a free HIV test *or* a Conventional Blood HIV Test

Print materials were collected from qualifying clinics under the following criteria,

1. The data collected must be a print material defined in *The Patient Education Materials Assessment Tool (PEMAT) and User's Guide* as being a “brochure, pamphlet, or materials that can be printed from websites like PDFs”
2. The print material must be free and visibly on display in the waiting room area of each clinic
3. The print materials collected must contain the language, “Human Immunodeficiency Virus” (HIV) or “PrEP” in the title
4. The language of the print materials collected may be English or Spanish

All print materials with language that specifically contain *Human immunodeficiency virus (HIV)*, *Acquired immunodeficiency syndrome (AIDS)*, or *Pre-exposure prophylaxis (PrEP)* were collected from the designated public waiting area of each clinic that met the inclusion criteria.

A quality assessment of the print materials gathered was conducted using the Patient Education Materials Assessment Tool for Printable Materials (PEMAT-P) evaluation tool.

#### **Selection of the Evaluation Tool**

The PEMAT-P is a 17-item questionnaire that produces a numeric score for understandability and actionability of the content of each print material examined. The PEMAT-P was selected as an evaluation tool for this study because it is an approved evaluation method

by the Agency for Healthcare and Research Quality, and because it is designed to be administered by health professionals and lay people (Shoemaker & Brach, 2013).

### **Raters**

Because the assessment tool is scored by an objective rater assessment of each material, and an inter-rater agreement score was determined for each item. Two additional raters were recruited from Georgia State University's School of Public Health. Each rater was required to read *The Patient Education Materials Assessment Tool (PEMAT) and User's Guide* prior to viewing the questionnaire or viewing collected data.

For the evaluation session, each rater was independently given a notebook with each print material collected in a sheet protector, for ease of the front and back of each item. To improve ease of rating, scoring, and data analyses, the PEMAT-P questionnaire was reproduced from its original published PDF form as an online survey using the Qualtrics platform. The section headings and question items were directly copied from the PEMAT-P assessment tool originally published to the Agency for Healthcare Research and quality website<sup>2</sup>. Responses were coded to auto sum a 1 or 0 for each item. Each rater was given a unique link to the survey via email. Each print material was assigned a letter that indicated to the rater which material to enter into the Qualtrics survey.

### **Scoring**

Raters were instructed to complete a PEMAT-P questionnaire for each print material (n=11) to be assessed. Raters were asked not to rely on prior knowledge when answering each questionnaire statement and to refer only to the content of the print material being evaluated to

---

2

answer the PEMAT-P questionnaire. Raters are instructed to mark each item as, “Agree”, “Disagree”, or “Not Applicable” for a 17-item section for *Understandability* and a 7-item section for *Actionability*. Example PEMAT-P items include (Shoemaker & Brach, 2013),

“The material uses common, everyday language.”

“The material’s visual aids have clear titles or captions”

“The material uses the active voice”

“The material uses visual aids whenever they could make it easier to act on the instructions”

The user is instructed to rate an item as, “Agree” if the question characteristic true of the content in the printed material being assessed between 80-100% of the time. The questionnaire utilizes a binary scoring system wherein “Agree” gets a score of 1, “Disagree” gets a score of 0, and “Not Applicable” is not scored. The material’s scores are calculated by calculating the percentage score of “Applicable” questions for the understandability and actionability sections of the PEMAT-P, respectively.

### **Evaluation plan**

In order to answer the research questions,

- A simple statistical analysis will be conducted to determine the prevalence of free, printed literature with information about HIV, AIDS, STIs and PrEP made available in clinics that meet the inclusion criteria
- The PEMAT-P assessment tool will be used to generate an *understandability score* and an *actionability score* for each print material. Three raters will independently complete a PEMAT-P survey for n=11 print materials
- Independent raters will come together to review each response as a group to determine an inter-rater agreement response for each item of the PEMAT-P

- Data analysis of user-agreed response frequency of all survey items relating to *The Use of Visual Aids* in the Understandability section of the PEMAT-P will be conducted

## CHAPTER IV RESULTS

### **Prevalence of qualifying data for evaluation**

A total of n=11 print materials were collected from 8 clinics in Dekalb County, Georgia. Clinics were selected from the Center for Disease Control's online *Get Tested: National HIV and STD Testing* directory (CDC, 2018). Of the 10 clinics and healthcare providers that met the inclusion criteria at the time of data collection, 8 had at least one educational pamphlet, brochure, or post card that displayed qualifying content for analysis. Two clinics did not have print literature that met the inclusion criteria.

For each evaluated print material, the PEMAT-P rater-agreed Understandability section scores ranked higher than rater-agreed Actionability section scores. On a scale of 0-17, the mean Understandability section score was 12 (70.5%). On a scale of 0-7, the mean Actionability score was 3.54 (50.6%). There were no statistical outliers for the frequency of "Agree" or "Disagree" response categories for any of the 24 PEMAT-P questionnaire items.



## **CHAPTER V DISCUSSION AND CONCLUSION**

### **Discussion of Research Questions**

The findings of this study highlight the need for an intensive focus on what kind of health communications content is most effective for Hispanic and Latino individuals in the United States. The CDC's directory database indicates that there is sufficient access to low-cost and free clinics within a 10-mile radius of the Doraville, GA zip code, 30341. Furthermore, 80% of the clinics that met the selection criteria had literature about HIV and safe-sex practices. For each evaluated print material, the PEMAT-P Understandability section score ranked higher than the Actionability section score. On a scale of 0-17, the mean Understandability section score was 12 (70.5%). On a scale of 0-7, the mean Actionability score was 3.54 (50.6%). Thus, it cannot be asserted that the evaluated print materials promote understandability and actionability in content and design. Congruent scores between raters that met the 80% threshold indicative of understandability and actionability of the printed materials were 27% and 0%, respectively. 3 brochures contained Spanish translations on the back of English content, however these materials did not rate significantly higher for understandability and actionability when compared to English-only materials.

### **Study Strengths and Limitations**

Using a dual rater approach, this study successfully utilized the Patient Education Materials Assessment Tool for Print Materials (PEMAT-P) to assess the constructs of *understandability* and *actionability* of health information resources developed for public audiences. Rater feedback indicated that the PEMAT-P user's manual and the scoring form were user-friendly and intuitive. Furthermore, the CDC's *Get Tested* database directory was a ready-made database from which to identify free and low-cost clinics for inclusion in the study.

The limitations of a very small sample size (n=11) yielded limited options for statistical analysis. Though simple statistical analysis was conducted to find the minimum, maximum, and mean scores for each summed section of the PEMAT-P assessment tool, non-parametric statistical analysis such as the Mann-Whitney U Test could have been employed to examine the difference in means between the Understandability and Actionability scores for evaluated materials. Furthermore, the experience of training and utilizing raters was time consuming. Coming to absolute agreement in inter-rater reliability scores for each PEMAT-P item for each material set back the timeline of this study by three weeks.

### **Recommendations for Future Research**

Because the mode of transmission and rates of survival significantly vary based on Hispanic subgroups, our interventions should not be *one size fits all* (Freiman, 1998). The findings of this study encourage further investigation into what types of HIV educational materials are available to Latino and Hispanics seeking testing and treatment in a healthcare setting.

Educational print materials that promote HIV prevention strategies represent low-cost resources with potential to reach Hispanic/Latino populations. However, careful review of the content's understandability and potential to trigger protective behavior among recipients in the intended populations is critically important. The results of this research reveal that there is vast potential for public health professionals improve effectiveness of educational safe sex promotional materials intended to reach vulnerable populations. There is a need for future research that assesses motivating factors towards actionability for this population as well as the impact of well-constructed visual aids on content understandability for this population in light of established barriers to care, including language barriers and limited medical health literacy

## References

- Berkman, N. D., Sheridan, S. L., Donahue, K. E., Halpern, D. J., & Crotty, K. (2011). Low Health Literacy and Health Outcomes: An Updated Systematic Review. *Annals of Internal Medicine*, 155(2), 97. doi:10.7326/0003-4819-155-2-201107190-00005
- Centers for Disease Control and Prevention. *Division of HIV/AIDS Prevention Strategic Plan 2017 - 2020* (Rep.). (2018, April). Retrieved January, 2019, from Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention (CDC), <https://www.cdc.gov/hiv/pdf/dhap/cdc-hiv-dhap-external-strategic-plan.pdf>
- Centers for Disease Control and Prevention (2018, January 29). *Social Determinants of Health: What Affects Health*. Retrieved from <https://www.cdc.gov/socialdeterminants/research/index.htm>
- Centers for Disease Control. [Diagnoses of HIV infection in the United States and dependent areas, 2016](#). *HIV Surveillance Report* 2017;28. Transmission categories in CDC's surveillance system are *male injection drug use, male-to-male sexual contact and injection drug use, and female injection drug use*. Retrieved from <https://www.cdc.gov/hiv/group/hiv-idu.html>
- United States, Executive Office of the President, Office of Management and Budget (OMB), Office of Information and Regulatory Affairs, Office of Management and Budget. (1997). *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*(210th ed., Vol. 62, p. 58782). Retrieved January, 2019, from <https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>
- U.S. Department of Health and Human Services. 2000. *Healthy People 2010*. Washington, DC: U.S. Government Printing Office. Originally developed for Ratzan SC, Parker RM. 2000. Introduction. In *National Library of Medicine Current Bibliographies in Medicine: Health Literacy*. Selden CR, Zorn M, Ratzan SC, Parker RM, Editors. NLM Pub. No. CBM 2000-1. Bethesda, MD: National Institutes of Health, U.S. Department of Health and Human Services
- Edward, J., Morris, S., Mataoui, F., Granberry, P., Williams, M. V., & Torres, I. (2018). The impact of health and health insurance literacy on access to care for Hispanic/Latino communities. *Public Health Nursing*, 35(3), 176-183. doi:10.1111/phn.12385
- Espinoza, L., Hall, H. I., Selik, R. M., & Hu, X. (2008). Characteristics of HIV Infection Among Hispanics, United States 2003-2006. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 49(1), 94-101. doi:10.1097/qai.0b013e3181820129
- Freiman M. P. (1998). The demand for healthcare among racial/ethnic subpopulations. *Health services research*, 33(4 Pt 1), 867–890. Retrieved February, 2019, from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1070291/>

*HIV Surveillance Report* (Rep. No. 29). (2018). Retrieved January, 2019, from Centers for Disease Control and Prevention website:  
<https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-2017-vol-29.pdf>

Kalichman, S. C., & Rompa, D. (2000). Functional health literacy is associated with health status and health-related knowledge in people living with HIV-AIDS. *Journal of Acquired Immune Deficiency Syndromes*, 25(4), 337-344. Retrieved from  
<http://ezproxy.gsu.edu/login?url=https://search-proquest-com.ezproxy.gsu.edu/docview/233631995?accountid=11226>

Marks, G., Millett, G. A., Bingham, T., Bond, L., Lauby, J., Liao, A., . . . Stueve, A. (2008). Understanding Differences in HIV Sexual Transmission among Latino and Black Men who have Sex with Men: The Brothers y Hermanos Study. *AIDS and Behavior*, 13(4), 682-690.  
doi:10.1007/s10461-008-9380-6

Ortega, A. N., Fang, H., Perez, V. H., Rizzo, J. A., Carter-Pokras, O., Wallace, S. P., & Gelberg, L. (2007). Health Care Access, Use of Services, and Experiences Among Undocumented Mexicans and Other Latinos. *Archives of Internal Medicine*, 167(21), 2354.  
doi:10.1001/archinte.167.21.2354

Parker R, Baker D, Williams M, Nurss J. The test of functional health literacy in adults (TOFHLA): a new instrument for measuring patients' literacy skills. *J Gen Intern Med*. 1995;10:537-541.

Rimer, B. K., Conaway, M., Lyna, P., Glassman, B., Yarnall, K. S., Lipkus, I., & Barber, L. (1999). The impact of tailored interventions on a community health center population. *Patient Education and Counseling*, 37(2), 125-140. doi:10.1016/s0738-3991(98)00122-0

Rodriguez MA, Bustamante AV, Ang A. Perceived quality of care, receipt of preventive care, and usual source of health care among undocumented and other Latinos. *J Gen Intern Med*. 2009;24(Suppl 3): 508–513.

Ryan, C., Huebner, D., Diaz, R. M., & Sanchez, J. (2009). Family Rejection as a Predictor of Negative Health Outcomes in White and Latino Lesbian, Gay, and Bisexual Young Adults. *Pediatrics*, 123(1), 346-352. doi:10.1542/peds.2007-3524

Shoemaker SJ, Wolf MS, Brach C. Patient Education Materials Assessment Tool for Printable Materials (PEMAT- P). (Prepared by Abt Associates under Contract No. HHS290200900012I, TO 4). Rockville, MD: Agency for Healthcare Research and Quality; October 2013. AHRQ Publication No. 14-0002-EF.

U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. National action plan to improve health literacy. 2010. Available from [http://www.health.gov/communication/hlactionplan/pdf/Health\\_Literacy\\_Action\\_Plan.pdf](http://www.health.gov/communication/hlactionplan/pdf/Health_Literacy_Action_Plan.pdf) [accessed 23.08.13]

## Appendix A: Reference list of materials gathered for the study

Centers for Disease Control and Prevention: Division of HIV/AIDS Prevention, National Center for HIV/AIDS. (2018). *HIV 101*[Brochure].

Centers for Disease Control and Prevention: Division of HIV/AIDS Prevention, National Center for HIV/AIDS. (march). *PrEP 101*[Brochure].

Department of Health and Human Services. (n.d.). *Your family's hopes and dreams are worth living for* [Brochure]. 7, 2014: [Postcard].

Krantzier, N. (2012). *HIV Facts* [Pamphlet]. Education, Training and Research.org. [Pamphlet].

Centers for Disease Control and Prevention. (2014). *Love him, Love yourself more*[Pamphlet].

Centers for Disease Control and Prevention, National Center for HIV Viral Hepatitis STD and TB Prevention. [Postcard]

Oakehbush, M. (2004). *HIV & STD*[Pamphlet]. Education, Training and Resources.org. [Pamphlet].

Centers for Disease Control and Prevention .(2018). *We're doing it: Testing for HIV* [Pamphlet].

Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention. [Postcard].

Centers for Disease Control and Prevention .(2018). *Lo Estamos Haciendo: La Prueba del*

*VIH*. [Pamphlet]. Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention. [Postcard].

## Author's Statement Page

In presenting this thesis as a partial fulfillment of the requirements for an advanced degree from Georgia State University, I agree that the Library of the University shall make it available for inspection and circulation in accordance with its regulations governing materials of this type. I agree that permission to quote from, to copy from, or to publish this thesis may be granted by the author or, in his/her absence, by the professor under whose direction it was written, or in his/her absence, by the Associate Dean, College of Health and Human Sciences. Such quoting, copying, or publishing must be solely for scholarly purposes and will not involve potential financial gain. It is understood that any copying from or publication of this dissertation which involves potential financial gain will not be allowed without written permission of the author.

Rebecca L Butler

Signature of Author

Notice to Borrowers Page

All theses deposited in the Georgia State University Library must be used in accordance with the stipulations prescribed by the author in the preceding statement.

The author of this thesis is: Rebecca Liliana Butler

Street Address: 808 Brookridge Drive NE

City, State, and Zip Code: Atlanta, GA 30306

The Chair of the committee for this thesis is:

Professor's Name: Sheryl Strasser, PhD, MPH, MSW, MCHES

Department: Institute of Public Health

College: Health and Human Sciences

Georgia State University  
P.O. Box 3995  
Atlanta, Georgia 30302-3995

Users of this thesis who not regularly enrolled as students at Georgia State University are required to attest acceptance of the preceding stipulation by signing below. Libraries borrowing this thesis for the use of their patrons are required to see that each user records here the information requested.

<b>NAME OF USER</b>	<b>ADDRESS</b>	<b>DATE</b>	<b>TYPE OF USE (EXAMINATION ONLY OR COPY)</b>

