

2011

Use of Granger Causality Analysis and Artificial Spike Trains to Examine Pause Coding in Purkinje Cell Spike Activity Related to Rhythmic Licking

Selva K. Maran

Ying Cao

Mukesh Dhamala

Georgia State University, mdhamala@gsu.edu

Follow this and additional works at: http://scholarworks.gsu.edu/neurosci_facpub

 Part of the [Neuroscience and Neurobiology Commons](#)

Recommended Citation

Maran *et al.*: Use of Granger causality analysis and artificial spike trains to examine pause coding in Purkinje cell spike activity related to rhythmic licking. *BMC Neuroscience* 2011, 12(Suppl 1):P143. doi: 10.1186/1471-2202-12-S1-P143

This Presentation is brought to you for free and open access by the Neuroscience Institute at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Neuroscience Institute Faculty Publications by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact scholarworks@gsu.edu.

POSTER PRESENTATION

Open Access


Use of Granger causality analysis and artificial spike trains to examine pause coding in Purkinje cell spike activity related to rhythmic licking

Selva K Maran^{1*}, Ying Cao², Mukesh Dhamala³, Detlef Heck², Dieter Jaeger¹

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Cerebellar Purkinje cells in mice show strong activity modulation related to rhythmic licking behavior [1]. Here we examine whether this modulation may preferentially be related to long inter-spike intervals (ISIs), i.e. pauses in spike activity. A preferential use of spike pauses for event coding of Purkinje cells has previously

been suggested [2,3]. We analyzed the lick modulation of Purkinje cell spike trains by several methods. First, we used peri-lick time histograms constructed separately by spikes initiating ISIs of different duration (Fig. 1B,C). We found that short and long ISIs showed lick modulation, but licks of intermediate duration did



* Correspondence: sseland@emory.edu

¹Department of Biology, Emory University, Atlanta, Georgia, 30322, USA
Full list of author information is available at the end of the article

not. Next we conducted a wavelet-based frequency resolved Granger causality analysis [4] to determine whether ISIs of different duration were caused by licks and/or were causal in the control of lick timing. We found there was a peak at 6 Hz from lick to spike in 12 cells, and a smaller peak also at 6 Hz from spike to lick in 5 out of those 12 cells. The peaks for plots (scaled to a possible maximum of 1.0) ranged from 0.1 to 0.8 for lick to spike causality and from 0.1 to 0.3 for spike to lick causality. Again, we found that short (0-10 ms) and long (40 ms and above) were preferentially involved in lick coding. To better understand the ISI duration dependence of lick coding we constructed controlled artificial spike trains from gamma ISI distributions [5], each matching slow spontaneous and lick-triggered spike rate fluctuations. The surrogate data show the same results as the biological ones in terms of preferential short and long ISI involvement in lick modulation. However, causality from spike data to licks was not observed in the artificial data. From these results we infer that any relation between specific ISI durations and lick modulation is due to the statistical properties of rate modulated spike trains, and can be mimicked by a simple gamma - distributed process. We also found that even the spike of single Purkinje cells can be predictive of lick events, suggesting Purkinje cell activity is involved in the control of lick timing.

Author details

¹Department of Biology, Emory University, Atlanta, Georgia, 30322, USA.

²Department of Anatomy and Neurobiology, UTHSC, Memphis, Tennessee, 38163, USA. ³Department of Physics and Astronomy, GSU, Atlanta, Georgia, 30303, USA.

Published: 18 July 2011

References

1. Bryant JL, Boughter JD, Gong S, LeDoux MS, Heck DH: **Cerebellar cortical output encodes temporal aspects of rhythmic licking movements and is necessary for normal licking frequency.** *Eur J Neurosci* 2010, **32**:41-52.
2. Steuber V, Mittmann W, Hoebeek FE, Silver RA, De Zeeuw CI, Häusser M, De Schutter E: **Cerebellar LTD and pattern recognition by Purkinje cells.** *Neuron* 2007, **54**:121-136.
3. Shin SL, Hoebeek FE, Schonewille M, De Zeeuw CI, Aertsen A, De Schutter E: **Regular patterns in cerebellar Purkinje cell simple spike trains.** *PLoS One* 2007, **2**:485.
4. Dhamala M, Rangarajan G, Ding M: **Estimating Granger causality from fourier and wavelet transforms of time series data.** *Phys Rev Lett* 2008, **100**:018701.
5. Shimokawa T, Shinomoto S: **Estimating instantaneous irregularity of neuronal firing.** *Neural Comput* 2009, **21**:1931-1951.

doi:10.1186/1471-2202-12-S1-P143

Cite this article as: Maran et al.: Use of Granger causality analysis and artificial spike trains to examine pause coding in Purkinje cell spike activity related to rhythmic licking. *BMC Neuroscience* 2011 **12**(Suppl 1):P143.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

